

Table of Contents Section II – Soil and Site Information

	Issue Date	Date of Last Review	Responsible Staff
<i>Use and Explanation of Soil Interpretations</i>			
<i>Explanation of Key Phrases Used in Soil Interpretations</i>			
Soils Legends	1/02	1/02	SOI
<i>* Acreage and Proportionate Extent of the Soils</i>			
Soil Descriptions - Nontechnical	1/02	1/93	SOI
<i>Use and Explanation of Nontechnical Descriptions</i>			
<i>*Nontechnical Soils Description Report</i>			
Soil Descriptions - Technical	1/02	1/02	SOI
<i>*Map Unit Description Report</i>			
Cropland Interpretations - Technical	1/02	1/02	SOI
<i>*Prime Farmland Report</i>			
<i>* Kansas Soil Rating for Plant Growth Index</i>			
<i>*Soil Properties for Conservation Planning</i>			
Rangeland, Grazed Forestland, Native Pastureland Interpretations	1/02	1/02	SOI
<i>*Rangeland Productivity Report</i>			
<i>*Range Site Descriptions</i>			
Forestland Interpretations	1/93	1/93	SOI
<i>Use and Explanation of Forestland Interpretations</i>			
<i>*Woodland Management and Productivity</i>			
Nonagricultural Interpretations	1/02	1/02	SOI
<i>*Building Site Development Report</i>			
<i>*Construction Materials Report</i>			
Recreation Interpretations	1/02	1/02	SOI
<i>*Recreational Interpretations</i>			
Wildlife Interpretations	1/02	1/02	SOI
<i>*Wildlife Interpretations Report</i>			
Pastureland and Hayland Interpretations	1/02	1/02	SOI
<i>*Yields Per Acre of Pasture and Hayland</i>			

	Issue Date	Date of Last Review	Responsible Staff
Mined Land Interpretations <i>Use and Explanation of Mined Land Interpretations</i>	1/93	1/93	SOI
Windbreak Interpretations <i>*Conservation Tree and Shrub Management Report</i>	1/02	1/02	SOI
Engineering Interpretations <i>*Engineering Index Properties</i> <i>*Physical Properties of the Soils</i> <i>*Chemical Properties of the Soils</i> <i>*Water Features</i> <i>*Soil Features</i> <i>*Water Management Report</i>	1/02	1/02	SOI
Waste Disposal Interpretations <i>*Sanitary Facilities Report</i> <i>*Agricultural Waste Management Report</i>	1/02	1/02	SOI
Water Quantity and Quality Interpretations <i>Use and Explanation of Water Quantity and Quality Interpretations</i> <i>*Appendix A – Soils Potential For Surface Loss and Leaching</i> <i>*Appendix B – Pesticide Selected Properties Database</i> <i>*Appendix C – Herbicide Selected Properties Database</i> <i>*Soil-Pesticide Interaction Screening Procedure Worksheet (Blank)</i> <i>*WIN-PST SPISP II Soil Sensitivity to Pesticide Loss Rating Report</i>	1/02	1/02	SOI
Hydric Soil Interpretations <i>Use and Explanation of Hydric Soil Interpretations</i> <i>*Hydric Soils List</i>	1/02	1/02	SOI
HEL Interpretations <i>Use and Explanation of Highly Erodible Land Interpretations</i> <i>*Highly Erodible Lands Report</i> <i>*LS and Supporting Data for 1990 Frozen HEL List</i> <i>*CRP 20 Soil Supporting Data for 1990 Frozen HEL List</i>	7/95	1/00	SOI

**County specific computer generated reports.*

ACREAGE AND PROPORTIONATE EXTENT OF THE SOILS

Barber County, Kansas: Published

Map symbol	Soil name	Acres	Percent
033CP	Clairemont Loam, Channeled-----	1,144	0.2
033OA	Obaro Silty Clay Loam, 5 To 12 Percent Slopes-----	508	*
033OB	Obaro-Rock Outcrop Complex, 10 To 30 Percent Slopes-----	2,092	0.3
033QR	Quinlan-Woodward Loams, 6 To 15 Percent Slopes-----	2,673	0.4
033QT	Quinlan-Woodward Loams, 15 To 30 Percent Slopes-----	527	*
033YE	Yahola Fine Sandy Loam, Occasionally Flooded-----	41	*
077BP	Woodward-Port Complex, 0 To 20 Percent Slopes-----	32	*
077FU	Farnum Loam, 3 To 6 Percent Slopes-----	4	*
077GE	Gerlane Fine Sandy Loam, Occasionally Flooded-----	373	*
077MC	Minco Silt Loam, 0 To 1 Percent Slopes-----	34	*
077MN	Minco Silt Loam, 1 To 3 Percent Slopes-----	1,268	0.2
077MO	Minco Silt Loam, 3 To 6 Percent Slopes-----	5	*
077QU	Quinlan Loam, 3 To 6 Percent Slopes-----	8	*
077SE	Shellabarger Fine Sandy Loam, 1 To 3 Percent Slopes-----	4	*
077WE	Woodward-Quinlan Loams, 1 To 3 Percent Slopes-----	382	*
095AC	Albion Sandy Loam, 3 To 6 Percent Slopes-----	15	*
095DA	Dillwyn-Plevna Complex, Occasionally Flooded-----	254	*
095FA	Farnum Sandy Loam, 0 To 2 Percent Slopes-----	309	*
095PB	Pratt Loamy Fine Sand, 1 To 5 Percent Slopes-----	159	*
095SB	Shellabarger Sandy Loam, 1 To 3 Percent Slopes-----	1,034	0.1
095WA	Waldeck Fine Sandy Loam, Occasionally Flooded-----	20	*
097LN	Lincoln Sandy Loam, Occasionally Flooded-----	32	*
097QW	Quinlan-Woodward Loams, 6 To 25 Percent Slopes-----	628	*
097WA	Waldeck Loam, Occasionally Flooded-----	18	*
151AB	Albion Sandy Loam, 1 To 4 Percent Slopes-----	94	*
151AO	Albion Sandy Loam, 3 To 7 Percent Slopes, Eroded-----	18	*
151AS	Albion And Shellabarger Soils, 7 To 15 Percent Slopes-----	12,371	1.7
151BC	Blanket Silty Clay Loam, 1 To 4 Percent Slopes, Eroded-----	186	*
151CA	Carwile Fine Sandy Loam, 0 To 1 Percent Slopes-----	30	*
151CK	Case-Clark Complex, 7 To 15 Percent Slopes-----	42	*
151CM	Clark Clay Loam, 1 To 4 Percent Slopes-----	386	*
151CN	Clark Fine Sandy Loam, 1 To 3 Percent Slopes-----	17	*
151CO	Clark-Ost Clay Loams, 0 To 1 Percent Slopes-----	9	*
151FA	Farnum Clay Loam, 3 To 6 Percent Slopes, Eroded-----	33	*
151KP	Kanza-Plevna Complex, Frequently Flooded-----	363	*
151OS	Ost Clay Loam, 1 To 4 Percent Slopes-----	750	0.1
151PM	Pratt Loamy Fine Sand, 3 To 8 Percent Slopes-----	458	*
151PN	Pratt Loamy Fine Sand, 8 To 12 Percent Slopes-----	281	*
151PO	Pratt-Carwile Complex, 0 To 5 Percent Slopes-----	105	*
151SA	Albion-Kaski Complex, 0 To 15 Percent Slopes-----	103	*
151SE	Shellabarger Fine Sandy Loam, 1 To 4 Percent Slopes-----	28	*
1439	Crisfield Sandy Loam, Rarely Flooded-----	140	*
Ad	Albion-Shellabarger Sandy Loams, 2 To 4 Percent Slopes-----	1,423	0.2
Ae	Albion And Shellabarger Soils, 4 To 15 Percent Slopes-----	86,922	11.9
AED	Arents, Earthen Dam-----	16	*
AN	Albion-Shellabarger Sandy Loams, 4 To 15 Percent Slopes-----	27,056	3.7
As	Clairemont Soils, Saline, Channeled-----	2,344	0.3
At	Attica Loamy Fine Sand, 1 To 5 Percent Slopes-----	7,139	1.0
Ba	Blanket Silt Loam, 0 To 1 Percent Slopes-----	3,996	0.5
Bb	Blanket Silt Loam, 1 To 3 Percent Slopes-----	9,892	1.4
Bc	Blanket Silty Clay Loam, 1 To 3 Percent Slopes, Eroded-----	1,767	0.2
Bf	Quinlan-Clairemont Complex, 0 To 50 Percent Slopes-----	10,003	1.4
Ca	Canadian Fine Sandy Loam, Rarely Flooded-----	4,934	0.7
Cc	Case-Clark Clay Loams, 2 To 6 Percent Slopes-----	2,901	0.4
Cd	Clairemont Silt Loam, Occasionally Flooded-----	11,600	1.6
Cf	Clairemont Soils, Channeled-----	26,653	3.7
Ck	Clark Clay Loam, 0 To 2 Percent Slopes-----	3,809	0.5
Fa	Farnum Fine Sandy Loam, 0 To 1 Percent Slopes-----	2,614	0.4
Fm	Farnum Loam, 0 To 1 Percent Slopes-----	4,140	0.6
Fr	Farnum Loam, 1 To 3 Percent Slopes-----	14,247	2.0
Fu	Farnum Clay Loam, 1 To 3 Percent Slopes, Eroded-----	3,503	0.5
Ga	Grant Silt Loam, 0 To 1 Percent Slopes-----	1,578	0.2
Gb	Grant Silt Loam, 1 To 3 Percent Slopes-----	38,032	5.2
Gc	Grant Silt Loam, 3 To 6 Percent Slopes-----	15,380	2.1
GRP	Gravel Pits-----	80	*
INT	Aquolls-----	167	*
Ka	Kanza Soils, Frequently Flooded-----	3,503	0.5
Kf	Kingfisher Silt Loam, 1 To 3 Percent Slopes-----	4,799	0.7
Kv	Kingfisher-Vernon Complex, 1 To 3 Percent Slopes-----	1,195	0.2
Kz	Kingfisher-Vernon Complex, 3 To 6 Percent Slopes-----	13,527	1.9
Ln	Lincoln Soils, Frequently Flooded-----	23,798	3.3
LNN	Lincoln Loamy Sand, Occasionally Flooded-----	2,351	0.3
Ma	Mangum Clay, Occasionally Flooded-----	2,677	0.4
Mg	Mangum-Drummond Complex, Rarely Flooded-----	2,084	0.3
Mn	Minco Silt Loam, 0 To 2 Percent Slopes-----	908	0.1
Na	Naron Fine Sandy Loam, 0 To 1 Percent Slopes-----	3,628	0.5
Nb	Naron Fine Sandy Loam, 1 To 3 Percent Slopes-----	11,907	1.6
Os	Ost Clay Loam, 0 To 1 Percent Slopes-----	1,710	0.2
Ot	Ost Clay Loam, 1 To 3 Percent Slopes-----	7,477	1.0
Pa	Pond Creek Silt Loam, 0 To 1 Percent Slopes-----	11,339	1.6
Pd	Pond Creek Silt Loam, 1 To 3 Percent Slopes-----	19,127	2.6
Pb	Dale Silt Loam, Rarely Flooded-----	22,137	3.0
Pk	Buttermilk Silt Loam, Rarely Flooded-----	736	0.1
Ps	Pratt Loamy Fine Sand, 5 To 10 Percent Slopes-----	2,248	0.3
Pt	Pratt-Tivoli Loamy Fine Sands, 5 To 15 Percent Slopes-----	11,536	1.6
Qn	Quinlan Loam, 1 To 3 Percent Slopes-----	3,548	0.5
Qw	Quinlan-Woodward Complex, 5 To 15 Percent Slopes-----	60,545	8.3
Rb	Knoco-Shale Outcrop Complex, 15 To 80 Percent Slopes-----	34,225	4.7
Sb	Shellabarger Sandy Loam, 3 To 6 Percent Slopes-----	12,381	1.7
SBB	Shellabarger Fine Sandy Loam, 0 To 1 Percent Slopes-----	101	*
Sc	Shellabarger Sandy Loam, 3 To 6 Percent Slopes, Eroded-----	3,894	0.5
Tv	Tivoli Fine Sand, 5 To 20 Percent Slopes-----	2,394	0.3

ACREAGE AND PROPORTIONATE EXTENT OF THE SOILS--Continued

Barber County, Kansas: Published

Map symbol	Soil name	Acres	Percent
Vn	Vernon Clay Loam, 3 To 5 Percent Slopes-----	2,172	0.3
Vr	Vernon Clay Loam, 5 To 15 Percent Slopes-----	81,563	11.2
Vs	Vernon-Knoco Complex, 1 To 5 Percent Slopes-----	20,163	2.8
W	Water-----	3,923	0.5
Wa	Waldeck Sandy Loam, Occasionally Flooded-----	1,889	0.3
Wo	Woodward-Quinlan Loams, 0 To 3 Percent Slopes-----	5,067	0.7
Ws	Woodward-Quinlan Loams, 3 To 6 Percent Slopes-----	32,273	4.4
Ya	Yahola Sandy Loam, Occasionally Flooded-----	8,373	1.2
Ze	Zenda Clay Loam, Occasionally Flooded-----	1,059	0.1
	Total-----	727,431	100.0

* Less than 0.1 percent.

Nontechnical Soil Descriptions
Barber County, Kansas

Nontechnical soil descriptions describe soil properties or management considerations specific to a soil map unit or group of map units, shown in the NonTechnical Descriptions report. These descriptions are written in terminology that Non-technical users of soil survey information can understand. Nontechnical soil descriptions are a powerful tool for creating reports. These high quality, easy to read reports can be generated by conservation planners and other NRCS employees for distribution to land users. Soil map unit descriptions and National Soil Information System records are the basis for these descriptions.

033CP Clairemont Loam, Channeled

Clairemont soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level flood plain. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Lowland (pe20-25) range site. It is in the nonirrigated land capability classification 6w.

0330A Obaro Silty Clay Loam, 5 To 12 Percent Slopes

Obaro soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hillslope on upland. The runoff class is high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

0330B Obaro-Rock Outcrop Complex, 10 To 30 Percent Slopes

Obaro soil makes up 75 percent of the map unit. This map unit is in the Central Rolling Red Prairies Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately steep hillslope on upland. The runoff class is high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

033QR Quinlan-Woodward Loams, 6 To 15 Percent Slopes

Quinlan soil makes up 55 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep hillslope on upland. The runoff class is medium. The parent material consists of residuum. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Shallow Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

Woodward soil makes up 45 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep hillslope on upland. The runoff class is medium. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

033QT Quinlan-Woodward Loams, 15 To 30 Percent Slopes

Quinlan soil makes up 55 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately steep to steep hillslope on upland. The runoff class is very high. The parent material consists of residuum. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Shallow Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 7e.

Woodward soil makes up 45 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately steep hillslope on upland. The runoff class is very high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

033YE Yahola Fine Sandy Loam, Occasionally Flooded

Yahola soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level flood plain. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland (pe20-25) range site. It is in the nonirrigated land capability classification 2w.

077BP Woodward-Port Complex, 0 To 20 Percent Slopes

Woodward soil makes up 65 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately steep hillslope on upland. The runoff class is high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is somewhat excessively drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 6e.

Port soil makes up 35 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping terrace on river valley. The runoff class is low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is frequently flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Lowland (pe24-32) range site. It is in the nonirrigated land capability classification 5w.

077FU Farnum Loam, 3 To 6 Percent Slopes

Farnum soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping paleoterrace on river valley. The runoff class is medium. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 3e.

077GE Gerlane Fine Sandy Loam, Occasionally Flooded

Gerlane soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level terrace. The runoff class is very low. The parent material consists of alluvium. This soil is moderately well drained. The slowest permeability is moderately slow. It has a high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 48 inches. This soil is in the Subirrigated (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

077MC Minco Silt Loam, 0 To 1 Percent Slopes

Minco soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level hillslope on upland. The runoff class is very low. The parent material consists of eolian deposits. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 1.

077MN Minco Silt Loam, 1 To 3 Percent Slopes

Minco soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is very low. The parent material consists of eolian deposits. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

077MO Minco Silt Loam, 3 To 6 Percent Slopes

Minco soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of eolian deposits. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 3e.

077QU Quinlan Loam, 3 To 6 Percent Slopes

Quinlan soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of residuum. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Shallow Prairie (pe24-32) range site. It is in the nonirrigated land capability classification 4e.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

077SE Shellabarger Fine Sandy Loam, 1 To 3 Percent Slopes

Shellabarger soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

077WE Woodward-Quinlan Loams, 1 To 3 Percent Slopes

Woodward soil makes up 50 percent of the map unit. This map unit is in the Central Rolling Red Plains Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

Quinlan soil makes up 50 percent of the map unit. This map unit is in the Central Rolling Red Plains Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of residuum. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Shallow Prairie (pe24-32) range site. It is in the nonirrigated land capability classification 3e.

095AC Albion Sandy Loam, 3 To 6 Percent Slopes

Albion soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. It is in the nonirrigated land capability classification 4e.

095DA Dillwyn-Plevna Complex, Occasionally Flooded

Dillwyn soil makes up 60 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level interdune on paleoterrace on river valley, dune on paleoterrace on river valley. The runoff class is negligible. The parent material consists of sandy eolian deposits. This soil is somewhat poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 24 inches. This soil is in the Subirrigated (pe24-32) range site. It is in the nonirrigated land capability classification 4w.

Plevna soil makes up 40 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 12 inches. This soil is in the Subirrigated (pe24-32) range site. It is in the nonirrigated land capability classification 5w.

095FA Farnum Sandy Loam, 0 To 2 Percent Slopes

Farnum soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping paleoterrace on river valley. The runoff class is very low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 3e.

095PB Pratt Loamy Fine Sand, 1 To 5 Percent Slopes

Pratt soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on paleoterrace on river valley. The runoff class is low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe24-32) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

095SB Shellabarger Sandy Loam, 1 To 3 Percent Slopes

Shellabarger soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

095WA Waldeck Fine Sandy Loam, Occasionally Flooded

Waldeck soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. This soil is in the Subirrigated (pe24-32) range site. It is in the nonirrigated land capability classification 3w.

097LN Lincoln Sandy Loam, Occasionally Flooded

Lincoln soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 66 inches. This soil is in the Sandy Lowland (pe20-25) range site. It is in the nonirrigated land capability classification 6w.

097QW Quinlan-Woodward Loams, 6 To 25 Percent Slopes

Quinlan soil makes up 55 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping to steep upland. The runoff class is medium. The parent material consists of loamy residuum weathered from calcareous sandstone. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

Woodward soil makes up 45 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep upland. The runoff class is medium. The parent material consists of coarse-silty residuum weathered from calcareous sandstone. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

097WA Waldeck Loam, Occasionally Flooded

Waldeck soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain, river valley. The runoff class is very low. The parent material consists of coarse-loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. This soil is in the Subirrigated (pe20-25) range site. It is in the nonirrigated land capability classification 3w.

151AB Albion Sandy Loam, 1 To 4 Percent Slopes

Albion soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. It is in the nonirrigated land capability classification 3e.

151AO Albion Sandy Loam, 3 To 7 Percent Slopes, Eroded

Albion soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. It is in the nonirrigated land capability classification 4e.

151AS Albion And Shellabarger Soils, 7 To 15 Percent Slopes

Albion soil makes up 50 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep paleoterrace on river valley. The runoff class is medium. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. It is in the nonirrigated land capability classification 6e.

Shellabarger soil makes up 50 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep paleoterrace on river valley. The runoff class is medium. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. It is in the nonirrigated land capability classification 6e.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

151BC Blanket Silty Clay Loam, 1 To 4 Percent Slopes, Eroded

Blanket soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping paleoterrace on river valley. The runoff class is medium. The parent material consists of clayey alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe21-28) range site. It is in the nonirrigated land capability classification 3e.

151CA Carwile Fine Sandy Loam, 0 To 1 Percent Slopes

Carwile soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level depression. The runoff class is medium. The parent material consists of loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. This soil is in the Sandy (pe21-28) range site. It is in the nonirrigated land capability classification 2w.

151CK Case-Clark Complex, 7 To 15 Percent Slopes

Case soil makes up 60 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep paleoterrace on river valley. The runoff class is high. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Limy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 6e.

Clark soil makes up 40 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a strongly sloping paleoterrace on river valley. The runoff class is high. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 45 percent calcium carbonate. This soil is in the Limy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 6e.

151CM Clark Clay Loam, 1 To 4 Percent Slopes

Clark soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 45 percent calcium carbonate. This soil is in the Limy Upland (pe21-28) range site. It is in the nonirrigated land capability classification 3e.

151CN Clark Fine Sandy Loam, 1 To 3 Percent Slopes

Clark soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 45 percent calcium carbonate. This soil is in the Limy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 3e.

151CO Clark-Ost Clay Loams, 0 To 1 Percent Slopes

Clark soil makes up 70 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 45 percent calcium carbonate. This soil is in the Limy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 2c.

Ost soil makes up 30 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 34 percent calcium carbonate. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 2c.

151FA Farnum Clay Loam, 3 To 6 Percent Slopes, Eroded

Farnum soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping paleoterrace on river valley. The runoff class is medium. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe21-28) range site. It is in the nonirrigated land capability classification 4e.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

151KP Kanza-Plevna Complex, Frequently Flooded

Kanza soil makes up 50 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 18 inches. It is in the nonirrigated land capability classification 5w.

Plevna soil makes up 50 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 12 inches. This soil is in the Subirrigated (pe21-28) range site. It is in the nonirrigated land capability classification 5w.

151OS Ost Clay Loam, 1 To 4 Percent Slopes

Ost soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping paleoterrace on river valley. The runoff class is medium. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 34 percent calcium carbonate. This soil is in the Loamy Upland (pe21-28) range site. It is in the nonirrigated land capability classification 2e.

151PM Pratt Loamy Fine Sand, 3 To 8 Percent Slopes

Pratt soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe21-28) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

151PN Pratt Loamy Fine Sand, 8 To 12 Percent Slopes

Pratt soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a strongly sloping dune on paleoterrace on river valley. The runoff class is low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe21-28) range site. It is in the nonirrigated land capability classification 6e.

151PO Pratt-Carwile Complex, 0 To 5 Percent Slopes

Pratt soil makes up 60 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe21-28) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

Carwile soil makes up 40 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level depression on paleoterrace on river valley. The runoff class is very low. The parent material consists of alluvium. This soil is somewhat poorly drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. This soil is in the Sandy (pe21-28) range site. It is in the nonirrigated land capability classification 2w.

151SA Albion-Kaski Complex, 0 To 15 Percent Slopes

Albion soil makes up 70 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep paleoterrace on river valley. The runoff class is medium. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe21-28) range site. It is in the nonirrigated land capability classification 6e.

Kaski soil makes up 30 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Lowland (pe21-28) range site. It is in the nonirrigated land capability classification 2w.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

151SE Shellabarger Fine Sandy Loam, 1 To 4 Percent Slopes

Shellabarger soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe21-28) range site. It is in the nonirrigated land capability classification 2e.

1439 Crisfield Sandy Loam, Rarely Flooded

Crisfield soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level to gently sloping terrace on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is moderately well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 40 inches. This soil is in the Sandy Terrace (pe24-32) range site. It is in the nonirrigated land capability classification 3s.

Ad Albion-Shellabarger Sandy Loams, 2 To 4 Percent Slopes

Albion soil makes up 60 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe20-25) range site. It is in the nonirrigated land capability classification 4e.

Shellabarger soil makes up 40 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe20-25) range site. It is in the nonirrigated land capability classification 3e.

Ae Albion And Shellabarger Soils, 4 To 15 Percent Slopes

Albion soil makes up 55 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

Shellabarger soil makes up 45 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

AN Albion-Shellabarger Sandy Loams, 4 To 15 Percent Slopes

Albion soil makes up 65 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

Shellabarger soil makes up 35 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

As Clairemont Soils, Saline, Channeled

Clairemont soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level flood plain. The runoff class is negligible. The parent material consists of alluvium. This soil is somewhat poorly drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil contains a moderately saline horizon. This soil is in the Saline Lowland (pe20-25) range site. It is in the nonirrigated land capability classification 6s.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

At Attica Loamy Fine Sand, 1 To 5 Percent Slopes

Attica soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. It is in the nonirrigated land capability classification 3e.

Ba Blanket Silt Loam, 0 To 1 Percent Slopes

Blanket soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level paleoterrace on river valley. The runoff class is very low. The parent material consists of clayey alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 2c.

Bb Blanket Silt Loam, 1 To 3 Percent Slopes

Blanket soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on river valley. The runoff class is medium. The parent material consists of clayey alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 2e.

Bc Blanket Silty Clay Loam, 1 To 3 Percent Slopes, Eroded

Blanket soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on river valley. The runoff class is medium. The parent material consists of clayey alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 3e.

Bf Quinlan-Clairemont Complex, 0 To 50 Percent Slopes

Quinlan soil makes up 60 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping to very steep hillslope on upland. The runoff class is very high. The parent material consists of residuum. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is excessively drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 7e.

Clairemont soil makes up 40 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level flood plain. The runoff class is very low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Lowland (pe20-25) range site. It is in the nonirrigated land capability classification 5w.

Ca Canadian Fine Sandy Loam, Rarely Flooded

Canadian soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level flood plain, river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Terrace (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

Cc Case-Clark Clay Loams, 2 To 6 Percent Slopes

Case soil makes up 50 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping paleoterrace on river valley. The runoff class is low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 25 percent calcium carbonate. This soil is in the Limy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 4e.

Clark soil makes up 50 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 45 percent calcium carbonate. This soil is in the Limy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 4e.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

Cd Clairemont Silt Loam, Occasionally Flooded

Clairemont soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level flood plain. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Lowland (pe20-25) range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Cf Clairemont Soils, Channeled

Clairemont soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain. The runoff class is very low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Lowland (pe20-25) range site. It is in the nonirrigated land capability classification 5w.

Ck Clark Clay Loam, 0 To 2 Percent Slopes

Clark soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 45 percent calcium carbonate. This soil is in the Limy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 3e.

Fa Farnum Fine Sandy Loam, 0 To 1 Percent Slopes

Farnum soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level paleoterrace on river valley. The runoff class is very low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 2e.

Fm Farnum Loam, 0 To 1 Percent Slopes

Farnum soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level paleoterrace on river valley. The runoff class is very low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 2c.

Fr Farnum Loam, 1 To 3 Percent Slopes

Farnum soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on river valley. The runoff class is low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Fu Farnum Clay Loam, 1 To 3 Percent Slopes, Eroded

Farnum soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on river valley. The runoff class is low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 3e.

Ga Grant Silt Loam, 0 To 1 Percent Slopes

Grant soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level terrace on upland. The runoff class is very low. The parent material consists of residuum. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 1.

Gb Grant Silt Loam, 1 To 3 Percent Slopes

Grant soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a gently sloping terrace on upland. The runoff class is low. The parent material consists of residuum. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

Gc Grant Silt Loam, 3 To 6 Percent Slopes

Grant soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a moderately sloping terrace on upland. The runoff class is medium. The parent material consists of residuum. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 3e.

INT Aquolls

Aquolls soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level depression on terrace on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is very poorly drained. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is occasional ponded. The top of the seasonal high water table is at 0 inches. It is in the nonirrigated land capability classification 5w.

Ka Kanza Soils, Frequently Flooded

Kanza soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 18 inches. This soil is in the Subirrigated (pe24-32) range site. It is in the nonirrigated land capability classification 5w.

Kf Kingfisher Silt Loam, 1 To 3 Percent Slopes

Kingfisher soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately slow. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 2e.

Kv Kingfisher-Vernon Complex, 1 To 3 Percent Slopes

Kingfisher soil makes up 70 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately slow. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 2e.

Vernon soil makes up 30 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Red Clay Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 3e.

Kz Kingfisher-Vernon Complex, 3 To 6 Percent Slopes

Kingfisher soil makes up 60 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderately slow. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 3e.

Vernon soil makes up 40 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Red Clay Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 4e.

Ln Lincoln Soils, Frequently Flooded

Lincoln soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 66 inches. This soil is in the Sandy Lowland (pe20-25) range site. It is in the nonirrigated land capability classification 6w.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

LNN Lincoln Loamy Sand, Occasionally Flooded

Lincoln soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 66 inches. This soil is in the Sandy Lowland (pe20-25) range site. It is in the nonirrigated land capability classification 6w.

Ma Mangum Clay, Occasionally Flooded

Mangum soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level flood plain. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is very slow. It has a high available water capacity and a high shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil contains a slightly saline horizon. This soil is in the Clay Lowland (pe20-25) range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Mg Mangum-Drummond Complex, Rarely Flooded

Mangum soil makes up 65 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level flood plain. The runoff class is very low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is very slow. It has a high available water capacity and a high shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil contains a slightly saline horizon. This soil is in the Clay Lowland (pe20-25) range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 3s.

Drummond soil makes up 35 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level terrace on river valley. The runoff class is very low. The parent material consists of clayey and/or loamy alluvium. This soil is moderately well drained. The slowest permeability is very slow. It has a low available water capacity and a high shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 48 inches. This soil contains a slightly saline horizon. This soil is in the Saline Lowland (pe20-25) range site. It is in the nonirrigated land capability classification 5s.

Mn Minco Silt Loam, 0 To 2 Percent Slopes

Minco soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level to gently sloping hillslope on upland. The runoff class is very low. The parent material consists of eolian deposits. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

Na Naron Fine Sandy Loam, 0 To 1 Percent Slopes

Naron soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level dune on paleoterrace on river valley. The runoff class is negligible. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 2e.

Nb Naron Fine Sandy Loam, 1 To 3 Percent Slopes

Naron soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 3e.

Os Ost Clay Loam, 0 To 1 Percent Slopes

Ost soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 34 percent calcium carbonate. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 2c.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

Ot Ost Clay Loam, 1 To 3 Percent Slopes

Ost soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 34 percent calcium carbonate. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

Pa Pond Creek Silt Loam, 0 To 1 Percent Slopes

Pond Creek soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level terrace. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 1.

Pd Pond Creek Silt Loam, 1 To 3 Percent Slopes

Pond Creek soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a gently sloping terrace. The runoff class is very low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

Ph Dale Silt Loam, Rarely Flooded

Dale soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level terrace on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Terrace (pe24-32) range site. It is in the nonirrigated land capability classification 1.

Pk Buttermilk Silt Loam, Rarely Flooded

Buttermilk soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level river valley, flood plain. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 63 inches. This soil contains a moderately saline horizon. This soil is in the Saline Lowland (pe24-32) range site. It is in the nonirrigated land capability classification 3s.

Ps Pratt Loamy Fine Sand, 5 To 10 Percent Slopes

Pratt soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe24-32) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

Pt Pratt-Tivoli Loamy Fine Sands, 5 To 15 Percent Slopes

Pratt soil makes up 70 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep dune on paleoterrace on river valley. The runoff class is low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

Tivoli soil makes up 30 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep dune on paleoterrace on river valley. The runoff class is low. The parent material consists of sandy eolian deposits. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe20-25) range site. It is in the nonirrigated land capability classification 7e.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

Qn Quinlan Loam, 1 To 3 Percent Slopes

Quinlan soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of residuum. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Shallow Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 3e.

Qw Quinlan-Woodward Complex, 5 To 15 Percent Slopes

Quinlan soil makes up 60 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep hillslope on upland. The runoff class is high. The parent material consists of residuum. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Shallow Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

Woodward soil makes up 40 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep hillslope on upland. The runoff class is high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

Rb Knoco-Shale Outcrop Complex, 15 To 80 Percent Slopes

Knoco soil makes up 70 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately steep to very steep hillslope on upland. The runoff class is very high. <parent material is missing> The soil is 4 to 19 inches deep to bedrock (paralithic). This soil is excessively drained. The slowest permeability is very slow. It has a very low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. It has a horizon that is moderately sodic. This soil is in the Red Shale (pe20-25) range site. It is in the nonirrigated land capability classification 7s.

Sb Shellabarger Sandy Loam, 3 To 6 Percent Slopes

Shellabarger soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe20-25) range site. It is in the nonirrigated land capability classification 3e.

SBB Shellabarger Fine Sandy Loam, 0 To 1 Percent Slopes

Shellabarger soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level paleoterrace on river valley. The runoff class is negligible. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe24-32) range site. It is in the nonirrigated land capability classification 2e.

Sc Shellabarger Sandy Loam, 3 To 6 Percent Slopes, Eroded

Shellabarger soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping paleoterrace on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe20-25) range site. It is in the nonirrigated land capability classification 3e.

Tv Tivoli Fine Sand, 5 To 20 Percent Slopes

Tivoli soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep dune on paleoterrace on river valley. The runoff class is low. The parent material consists of sandy eolian deposits. This soil is excessively drained. The slowest permeability is rapid. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Choppy Sands (pe20-25) range site. It is in the nonirrigated land capability classification 7e.

Nontechnical Soil Descriptions--Continued
Barber County, Kansas

Vn Vernon Clay Loam, 3 To 5 Percent Slopes

Vernon soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is low. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Red Clay Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 4e.

Vr Vernon Clay Loam, 5 To 15 Percent Slopes

Vernon soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep hillslope on upland. The runoff class is high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Red Clay Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 6e.

Vs Vernon-Knoco Complex, 1 To 5 Percent Slopes

Vernon soil makes up 50 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping hillslope on upland. The runoff class is low. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Red Clay Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 4e.

Knoco soil makes up 50 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of dense, noncemented residuum weathered from claystone. The soil is 4 to 19 inches deep to bedrock (paralithic). This soil is excessively drained. The slowest permeability is very slow. It has a very low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. It has a horizon that is moderately sodic. This soil is in the Red Shale (pe20-25) range site. It is in the nonirrigated land capability classification 7s.

Wa Waldeck Sandy Loam, Occasionally Flooded

Waldeck soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Prairies Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is very low. The parent material consists of alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. This soil is in the Subirrigated (pe24-32) range site. It is in the nonirrigated land capability classification 3w.

Wo Woodward-Quinlan Loams, 0 To 3 Percent Slopes

Woodward soil makes up 70 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Shallow Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 2e.

Quinlan soil makes up 30 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping hillslope on upland. The runoff class is low. The parent material consists of residuum. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 3e.

Ws Woodward-Quinlan Loams, 3 To 6 Percent Slopes

Woodward soil makes up 65 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Shallow Prairie (pe20-25) range site. It is in the nonirrigated land capability classification 3e.

NONTECHNICAL SOIL DESCRIPTIONS--Continued
Barber County, Kansas

Quinlan soil makes up 35 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of residuum. The soil is 10 to 20 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe20-25) range site. It is in the nonirrigated land capability classification 4e.

Ya Yahola Sandy Loam, Occasionally Flooded

Yahola soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain. The runoff class is very low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Subirrigated (pe20-25) range site. It is in the nonirrigated land capability classification 2w.

Ze Zenda Clay Loam, Occasionally Flooded

Zenda soil makes up 100 percent of the map unit. This map unit is in the Central Rolling Red Plains Major Land Resource Area. This soil occurs on a nearly level dune on paleoterrace on river valley. The runoff class is negligible. The parent material consists of sandy eolian deposits. This soil is somewhat poorly drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. This soil contains a very slightly saline horizon. This soil is in the Subirrigated (pe20-25) range site. It is in the nonirrigated land capability classification 2w.

**033CP—Clairemont loam,
channeled****Map Unit Composition**

Clairemont: 100 percent

Component Descriptions**Clairemont**

MLRA: 78 - Central Rolling Red Plains

Landform: Flood plain

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Frequent

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Loamy Lowland (pe20-25)

Land capability (nonirrigated): 6w

Typical Profile:

H1—0 to 15 inches; loam

H2—15 to 60 inches; silt loam

Minor Components**Unnamed Wet Soils**

Phase: Loamy, Drainageway

**033OA—Obaro silty clay loam, 5 to
12 percent slopes****Map Unit Composition**

Obaro: 100 percent

Component Descriptions**Obaro**

MLRA: 78 - Central Rolling Red Plains

Landform: Hillslope on upland

Parent material: Residuum

Slope: 5 to 12 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Low (About 6.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: High

Ecological site: Loamy Upland (pe20-25)

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 35 inches; silty clay loam

Cr—35 to 35 inches; weathered bedrock

**033OB—Obaro-Rock outcrop
complex, 10 to 30 percent slopes****Map Unit Composition**

Obaro: 75 percent

Rock outcrop: 25 percent

Component Descriptions**Obaro**

MLRA: 78 - Central Rolling Red Plains, 80A - Central Rolling Red Prairies

Landform: Hillslope on upland

Parent material: Residuum

Slope: 12 to 15 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Low (About 6.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: High

Ecological site: Loamy Upland (pe20-25)

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 35 inches; silty clay loam

Cr—35 to 35 inches; weathered bedrock

Rock outcrop

MLRA: 78 - Central Rolling Red Plains, 80A - Central Rolling Red Prairies

Slope: 10 to 30 percent
Drainage class: Excessively drained
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Land capability (nonirrigated): 8

033QR—Quinlan-Woodward loams, 6 to 15 percent slopes

Map Unit Composition

Quinlan: 55 percent
 Woodward: 45 percent

Component Descriptions

Quinlan

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 6 to 15 percent
Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very low (About 2.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Shallow Prairie (pe20-25)
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 14 inches; loam
 Cr—14 to 14 inches; weathered bedrock

Woodward

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 6 to 15 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Low (About 5.1 inches)

Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Loamy Upland (pe20-25)
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 30 inches; loam
 Cr—30 to 30 inches; weathered bedrock

033QT—Quinlan-Woodward loams, 15 to 30 percent slopes

Map Unit Composition

Quinlan: 55 percent
 Woodward: 45 percent

Component Descriptions

Quinlan

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 15 to 30 percent
Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very low (About 2.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Shallow Prairie (pe20-25)
Land capability (nonirrigated): 7e

Typical Profile:

H1—0 to 14 inches; loam
 Cr—14 to 14 inches; weathered bedrock

Woodward

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 15 to 20 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Low (About 5.1 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Loamy Upland (pe20-25)
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 30 inches; loam
 Cr—30 to 30 inches; weathered bedrock

033YE—Yahola fine sandy loam, occasionally flooded

Map Unit Composition

Yahola: 100 percent

Component Descriptions

Yahola

MLRA: 78 - Central Rolling Red Plains
Landform: Flood plain
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: Moderate (About 8.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Occasional
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Sandy Lowland (pe20-25)
Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 12 inches; fine sandy loam
 H2—12 to 40 inches; fine sandy loam
 H3—40 to 60 inches; stratified loam to loamy fine sand

Minor Components

Unnamed Wet Soils

Phase: Sandy, Drainageway

077BP—Woodward-Port complex, 0 to 20 percent slopes

Map Unit Composition

Woodward: 65 percent
 Port: 35 percent

Component Descriptions

Woodward

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 1 to 20 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Somewhat excessively drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Low (About 4.1 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: High
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 24 inches; silt loam
 Cr—24 to 24 inches; weathered bedrock

Port

MLRA: 78 - Central Rolling Red Plains
Landform: Terrace on river valley
Parent material: Alluvium
Slope: 0 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.8 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Lowland (pe24-32)
Land capability (nonirrigated): 5w

Typical Profile:

H1—0 to 27 inches; silt loam
H2—27 to 60 inches; silty clay loam

Minor Components**Unnamed Wet Soils**

Phase: Sandy, Depression

Unnamed Wet Soils

Phase: Sandy, Drainageway

077FU—Farnum loam, 3 to 6 percent slopes**Map Unit Composition**

Farnum: 100 percent

Component Descriptions**Farnum**

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Alluvium
Slope: 3 to 6 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.1 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 11 inches; loam
H2—11 to 41 inches; clay loam
H3—41 to 60 inches; clay loam

077GE—Gerlane fine sandy loam, occasionally flooded**Map Unit Composition**

Gerlane: 100 percent

Component Descriptions**Gerlane**

MLRA: 80A - Central Rolling Red Prairies
Landform: Terrace
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Moderately well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Moderate (About 8.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Occasional
Depth to seasonal water saturation: About 24 to 72 inches
Runoff class: Very low
Ecological site: Subirrigated (pe24-32)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 17 inches; fine sandy loam
H2—17 to 40 inches; fine sandy loam
H3—40 to 48 inches; loamy sand
H4—48 to 60 inches; clay loam

Minor Components**Unnamed Wet Soils**

Phase: Sandy, Drainageway

077MC—Minco silt loam, 0 to 1 percent slopes**Map Unit Composition**

Minco: 100 percent

Component Descriptions**Minco**

MLRA: 80A - Central Rolling Red Prairies
Landform: Hillslope on upland
Parent material: Eolian deposits
Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.1 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Loamy Upland (pe24-32)

Land capability (nonirrigated): 1

Typical Profile:

H1—0 to 42 inches; silt loam

H2—42 to 60 inches; silt loam

077MN—Minco silt loam, 1 to 3 percent slopes

Map Unit Composition

Minco: 100 percent

Component Descriptions

Minco

MLRA: 80A - Central Rolling Red Prairies

Landform: Hillslope on upland

Parent material: Eolian deposits

Slope: 1 to 3 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.1 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Loamy Upland (pe24-32)

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 42 inches; silt loam

H2—42 to 60 inches; silt loam

077MO—Minco silt loam, 3 to 6 percent slopes

Map Unit Composition

Minco: 100 percent

Component Descriptions

Minco

MLRA: 80A - Central Rolling Red Prairies

Landform: Hillslope on upland

Parent material: Eolian deposits

Slope: 3 to 6 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.1 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Loamy Upland (pe24-32)

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 42 inches; silt loam

H2—42 to 60 inches; silt loam

077QU—Quinlan loam, 3 to 6 percent slopes

Map Unit Composition

Quinlan: 100 percent

Component Descriptions

Quinlan

MLRA: 78 - Central Rolling Red Plains

Landform: Hillslope on upland

Parent material: Residuum

Slope: 3 to 6 percent

Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Very low (About 1.7 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Shallow Prairie (pe24-32)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 9 inches; loam
Cr—9 to 9 inches; weathered bedrock

077SE—Shellabarger fine sandy loam, 1 to 3 percent slopes

Map Unit Composition

Shellabarger: 100 percent

Component Descriptions

Shellabarger

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Moderate (About 8.8 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy (pe24-32)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 13 inches; fine sandy loam
H2—13 to 38 inches; fine sandy loam
H3—38 to 60 inches; coarse sandy loam

077WE—Woodward-Quinlan loams, 1 to 3 percent slopes

Map Unit Composition

Woodward: 50 percent
Quinlan: 50 percent

Component Descriptions

Woodward

MLRA: 78 - Central Rolling Red Plains,80A -
Central Rolling Red Prairies
Landform: Hillslope on upland

Parent material: Residuum
Slope: 1 to 3 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Low (About 4.1 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 24 inches; loam
Cr—24 to 24 inches; weathered bedrock

Quinlan

MLRA: 78 - Central Rolling Red Plains,80A -
Central Rolling Red Prairies
Landform: Hillslope on upland
Parent material: Residuum
Slope: 1 to 3 percent
Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very low (About 1.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Shallow Prairie (pe24-32)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 9 inches; loam
Cr—9 to 9 inches; weathered bedrock

Minor Components

Unnamed Wet Soils

Phase: Loamy, Drainageway

095AC—Albion sandy loam, 3 to 6 percent slopes

Map Unit Composition

Albion: 100 percent

Component Descriptions

Albion

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 3 to 6 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Low (About 5.8 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sandy (pe24-32)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 8 inches; sandy loam

H2—8 to 16 inches; sandy loam

H3—16 to 26 inches; loamy sand

H4—26 to 60 inches; sand

095DA—Dillwyn-Plevna complex, occasionally flooded

Map Unit Composition

Dillwyn: 60 percent

Plevna: 40 percent

Component Descriptions

Dillwyn

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley, interdune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Low (About 4.9 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: About 12 to 36 inches

Runoff class: Negligible

Ecological site: Subirrigated (pe24-32)

Land capability (nonirrigated): 4w

Typical Profile:

H1—0 to 8 inches; loamy fine sand

H2—8 to 60 inches; loamy fine sand

Plevna

MLRA: 79 - Great Bend Sand Plains

Landform: Flood plain on river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Poorly drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 6.5 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Frequent

Depth to seasonal water saturation: About 0 to 24 inches

Runoff class: Negligible

Ecological site: Subirrigated (pe24-32)

Land capability (nonirrigated): 5w

Typical Profile:

H1—0 to 11 inches; fine sandy loam

H2—11 to 36 inches; fine sandy loam

H3—36 to 60 inches; sand

Minor Components

Unnamed Wet Soils

Phase: Sandy, Depression

095FA—Farnum sandy loam, 0 to 2 percent slopes

Map Unit Composition

Farnum: 100 percent

Component Descriptions

Farnum

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 9.7 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low
Ecological site: Sandy (pe24-32)
Land capability (irrigated): 2e
Land capability (nonirrigated): 3e

Typical Profile:
 H1—0 to 16 inches; sandy loam
 H2—16 to 50 inches; clay loam
 H3—50 to 60 inches; clay loam

Minor Components
Carwile

Unnamed Wet Soils
Phase: Loamy, Depression

Unnamed Wet Soils
Phase: Loamy, Drainageway

095PB—Pratt loamy fine sand, 1 to 5 percent slopes

Map Unit Composition

Pratt: 100 percent

Component Descriptions

Pratt

MLRA: 79 - Great Bend Sand Plains
Landform: Dune on paleoterrace on river valley
Parent material: Sandy eolian deposits
Slope: 3 to 8 percent
Drainage class: Well drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Moderate (About 6.4 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Sands (pe24-32)
Land capability (irrigated): 3e
Land capability (nonirrigated): 4e

Typical Profile:
 H1—0 to 12 inches; loamy fine sand
 H2—12 to 36 inches; loamy fine sand
 H3—36 to 60 inches; loamy fine sand

Minor Components
Carwile

Unnamed Wet Soils
Phase: Sandy, Depression

095SB—Shellabarger sandy loam, 1 to 3 percent slopes

Map Unit Composition

Shellabarger: 100 percent

Component Descriptions

Shellabarger

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 9.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy (pe24-32)
Land capability (nonirrigated): 2e

Typical Profile:
 H1—0 to 10 inches; sandy loam
 H2—10 to 45 inches; sandy clay loam
 H3—45 to 60 inches; coarse sandy loam

Minor Components
Unnamed Wet Soils
Phase: Loamy, Depression

095WA—Waldeck fine sandy loam, occasionally flooded

Map Unit Composition

Waldeck: 100 percent

Component Descriptions

Waldeck

MLRA: 80A - Central Rolling Red Prairies

Landform: Flood plain on river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 6.9 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: About 24 to 48 inches

Runoff class: Negligible

Ecological site: Subirrigated (pe24-32)

Land capability (nonirrigated): 3w

Typical Profile:

H1—0 to 12 inches; fine sandy loam

H2—12 to 36 inches; sandy loam

H3—36 to 60 inches; sand

Minor Components

Plevna

Unnamed Wet Soils

Phase: Sandy, Depression

Unnamed Wet Soils

Phase: Sandy, Drainageway

097LN—Lincoln sandy loam, occasionally flooded

Map Unit Composition

Lincoln: 100 percent

Component Descriptions

Lincoln

MLRA: 80A - Central Rolling Red Prairies

Landform: Flood plain on river valley

Parent material: Alluvium

Slope: 0 to 2 percent

Drainage class: Somewhat excessively drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Low (About 3.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: About 60 to 72 inches

Runoff class: Negligible

Ecological site: Sandy Lowland (pe20-25)

Land capability (nonirrigated): 6w

Typical Profile:

H1—0 to 8 inches; sandy loam

H2—8 to 60 inches; stratified fine sand to loamy fine sand

Minor Components

Plevna

Unnamed Wet Soils

Phase: Sandy, Drainageway

097QW—Quinlan-Woodward loams, 6 to 25 percent slopes

Map Unit Composition

Quinlan: 55 percent

Woodward: 45 percent

Component Descriptions

Quinlan

MLRA: 78 - Central Rolling Red Plains

Landform: Upland

Parent material: Loamy residuum weathered from calcareous sandstone

Slope: 6 to 25 percent

Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Very low (About 2.8 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Shallow Prairie (pe20-25)

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 15 inches; loam

Cr—15 to 15 inches; weathered bedrock

Woodward

MLRA: 78 - Central Rolling Red Plains

Landform: Upland

Parent material: Coarse-silty residuum
weathered from calcareous sandstone

Slope: 6 to 20 percent

Depth to restrictive feature: 20 to 40 inches to
bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60
in/hr)

Available water capacity: Low (About 4.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6
feet

Runoff class: Medium

Ecological site: Loamy Upland (pe20-25)

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 27 inches; loam

Cr—27 to 27 inches; weathered bedrock

097WA—Waldeck loam, occasionally flooded

Map Unit Composition

Waldeck: 100 percent

Component Descriptions

Waldeck

MLRA: 80A - Central Rolling Red Prairies

Landform: Flood plain, river valley

Parent material: Coarse-loamy alluvium

Slope: 0 to 2 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderately rapid (About
2.00 in/hr)

Available water capacity: Moderate (About 7.8
inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: About 24 to
48 inches

Runoff class: Very low

Ecological site: Subirrigated (pe20-25)

Land capability (nonirrigated): 3w

Typical Profile:

H1—0 to 14 inches; loam

H2—14 to 41 inches; fine sandy loam

H3—41 to 60 inches; fine sand

Minor Components

Unnamed Wet Soils

Phase: Sandy, Drainageway

151AB—Albion sandy loam, 1 to 4 percent slopes

Map Unit Composition

Albion: 100 percent

Component Descriptions

Albion

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 1 to 4 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About
2.00 in/hr)

Available water capacity: Moderate (About 6.0
inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6
feet

Runoff class: Very low

Ecological site: Sandy (pe24-32)

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 8 inches; sandy loam

H2—8 to 18 inches; sandy loam

H3—18 to 29 inches; coarse sandy loam

H4—29 to 60 inches; gravelly sand

Minor Components

Unnamed Wet Soils

Phase: Sandy, Drainageway

151AO—Albion sandy loam, 3 to 7 percent slopes, eroded

Map Unit Composition

Albion: 100 percent

Component Descriptions

Albion

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 3 to 7 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 6.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sandy (pe24-32)

Land capability (nonirrigated): 4e

Typical Profile:

- H1—0 to 8 inches; sandy loam
- H2—8 to 18 inches; sandy loam
- H3—18 to 29 inches; coarse sandy loam
- H4—29 to 60 inches; gravelly sand

151AS—Albion and Shellabarger Soils, 7 to 15 percent slopes

Map Unit Composition

Albion: 50 percent

Shellabarger: 50 percent

Component Descriptions

Albion

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 7 to 15 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 6.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Sandy (pe24-32)

Land capability (nonirrigated): 6e

Typical Profile:

- H1—0 to 8 inches; sandy loam
- H2—8 to 18 inches; sandy loam
- H3—18 to 29 inches; coarse sandy loam
- H4—29 to 60 inches; gravelly sand

Shellabarger

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 7 to 15 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Moderate (About 8.5 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Sandy (pe24-32)

Land capability (nonirrigated): 6e

Typical Profile:

- H1—0 to 11 inches; fine sandy loam
- H2—11 to 34 inches; sandy clay loam
- H3—34 to 60 inches; coarse sandy loam

151BC—Blanket silty clay loam, 1 to 4 percent slopes, eroded

Map Unit Composition

Blanket: 100 percent

Component Descriptions

Blanket

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Clayey alluvium

Slope: 1 to 4 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 9.2 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Loamy Upland (pe21-28)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 13 inches; silty clay loam
 H2—13 to 46 inches; silty clay
 H3—46 to 60 inches; silty clay loam

151CA—Carwile fine sandy loam, 0 to 1 percent slopes

Map Unit Composition

Carwile: 100 percent

Component Descriptions

Carwile

MLRA: 79 - Great Bend Sand Plains
Landform: Depression
Parent material: Loamy alluvium
Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: High (About 9.4 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: About 0 to 0 inches
Runoff class: Medium
Ecological site: Sandy (pe21-28)
Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 12 inches; fine sandy loam
 H2—12 to 20 inches; sandy clay loam
 H3—20 to 33 inches; clay
 H4—33 to 60 inches; sandy clay loam

151CK—Case-Clark complex, 7 to 15 percent slopes

Map Unit Composition

Case: 60 percent
 Clark: 40 percent

Component Descriptions

Case

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Alluvium
Slope: 7 to 15 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.2 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: High
Ecological site: Limy Upland (pe24-32)
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 6 inches; clay loam
 H2—6 to 60 inches; clay loam

Clark

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 7 to 12 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.3 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: High
Ecological site: Limy Upland (pe24-32)
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 8 inches; clay loam
 H2—8 to 60 inches; clay loam

151CM—Clark clay loam, 1 to 4 percent slopes

Map Unit Composition

Clark: 100 percent

Component Descriptions

Clark

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 1 to 4 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 10.3 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Limy Upland (pe21-28)

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 8 inches; clay loam

H2—8 to 60 inches; clay loam

Minor Components

Unnamed Wet Soils

Phase: Loamy, Depression

Unnamed Wet Soils

Phase: Loamy, Drainageway

151CN—Clark fine sandy loam, 1 to 3 percent slopes

Map Unit Composition

Clark: 100 percent

Component Descriptions

Clark

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 1 to 3 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 10.0 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Limy Upland (pe24-32)

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 8 inches; fine sandy loam

H2—8 to 60 inches; clay loam

151CO—Clark-Ost clay loams, 0 to 1 percent slopes

Map Unit Composition

Clark: 70 percent

Ost: 30 percent

Component Descriptions

Clark

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 10.3 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Limy Upland (pe24-32)

Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 8 inches; clay loam

H2—8 to 60 inches; clay loam

Ost

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 10.3 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 9 inches; clay loam
 H2—9 to 14 inches; clay loam
 H3—14 to 23 inches; clay loam
 H4—23 to 60 inches; clay loam

151FA—Farnum clay loam, 3 to 6 percent slopes, eroded

Map Unit Composition

Farnum: 100 percent

Component Descriptions

Farnum

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Alluvium
Slope: 3 to 6 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 9.9 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Loamy Upland (pe21-28)
Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 7 inches; clay loam
 H2—7 to 37 inches; clay loam
 H3—37 to 60 inches; fine sandy loam

151KP—Kanza-Plevna complex, frequently flooded

Map Unit Composition

Kanza: 50 percent
 Plevna: 50 percent

Component Descriptions

Kanza

MLRA: 79 - Great Bend Sand Plains
Landform: Flood plain on river valley
Parent material: Alluvium
Slope: 0 to 2 percent
Drainage class: Poorly drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 3.8 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: About 0 to 36 inches
Runoff class: Negligible
Land capability (nonirrigated): 5w

Typical Profile:

H1—0 to 11 inches; loamy fine sand
 H2—11 to 40 inches; fine sand

Plevna

MLRA: 79 - Great Bend Sand Plains
Landform: Flood plain on river valley
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Poorly drained
Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: Moderate (About 6.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: About 0 to 24 inches
Runoff class: Negligible
Ecological site: Subirrigated (pe21-28)
Land capability (nonirrigated): 5w

Typical Profile:

H1—0 to 10 inches; fine sandy loam
 H2—10 to 40 inches; fine sandy loam
 H3—40 to 60 inches; fine sand

151OS—Ost clay loam, 1 to 4 percent slopes

Map Unit Composition

Ost: 100 percent

Component Descriptions

Ost

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 1 to 4 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 10.3 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Loamy Upland (pe21-28)

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 9 inches; clay loam

H2—9 to 14 inches; clay loam

H3—14 to 23 inches; clay loam

H4—23 to 60 inches; clay loam

Minor Components

Unnamed Wet Soils

Phase: Loamy, Depression

Unnamed Wet Soils

Phase: Loamy, Drainageway

151PM—Pratt loamy fine sand, 3 to 8 percent slopes

Map Unit Composition

Pratt: 100 percent

Component Descriptions

Pratt

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 3 to 8 percent

Drainage class: Well drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Moderate (About 6.4 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sands (pe21-28)

Land capability (irrigated): 3e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 10 inches; loamy fine sand

H2—10 to 40 inches; loamy fine sand

H3—40 to 60 inches; loamy fine sand

Minor Components

Carwile

Unnamed Wet Soils

Phase: Sandy, Depression

151PN—Pratt loamy fine sand, 8 to 12 percent slopes

Map Unit Composition

Pratt: 100 percent

Component Descriptions

Pratt

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 8 to 12 percent

Drainage class: Well drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Moderate (About 6.4 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sands (pe21-28)

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 10 inches; loamy fine sand
 H2—10 to 40 inches; loamy fine sand
 H3—40 to 60 inches; loamy fine sand

151PO—Pratt-Carwile complex, 0 to 5 percent slopes

Map Unit Composition

Pratt: 60 percent
 Carwile: 40 percent

Component Descriptions

Pratt

MLRA: 79 - Great Bend Sand Plains
Landform: Dune on paleoterrace on river valley
Parent material: Sandy eolian deposits
Slope: 3 to 8 percent
Drainage class: Well drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Moderate (About 6.4 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sands (pe21-28)
Land capability (irrigated): 3e
Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 10 inches; loamy fine sand
 H2—10 to 40 inches; loamy fine sand
 H3—40 to 60 inches; loamy fine sand

Carwile

MLRA: 79 - Great Bend Sand Plains
Landform: Depression on paleoterrace on river valley
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: High (About 9.4 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: About 0 to 0 inches
Runoff class: Very low
Ecological site: Sandy (pe21-28)

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 12 inches; fine sandy loam
 H2—12 to 20 inches; sandy clay loam
 H3—20 to 33 inches; clay
 H4—33 to 60 inches; sandy clay loam

Minor Components

Unnamed Wet Soils

Phase: Sandy, Depression

151SA—Albion-Kaski complex, 0 to 15 percent slopes

Map Unit Composition

Albion: 70 percent
 Kaski: 30 percent

Component Descriptions

Albion

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 7 to 15 percent
Drainage class: Well drained
Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: Moderate (About 6.0 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Sandy (pe21-28)
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 8 inches; fine sandy loam
 H2—8 to 18 inches; sandy loam
 H3—18 to 29 inches; coarse sandy loam
 H4—29 to 60 inches; gravelly sand

Kaski

MLRA: 79 - Great Bend Sand Plains
Landform: Flood plain on river valley
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 10.5 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: Occasional
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Lowland (pe21-28)
Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 26 inches; loam
 H2—26 to 40 inches; loam
 H3—40 to 60 inches; sandy loam

151SE—Shellabarger fine sandy loam, 1 to 4 percent slopes

Map Unit Composition

Shellabarger: 100 percent

Component Descriptions

Shellabarger

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 1 to 4 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Moderate (About 8.5 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy (pe21-28)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 11 inches; fine sandy loam
 H2—11 to 34 inches; sandy clay loam
 H3—34 to 60 inches; coarse sandy loam

Minor Components

Unnamed Wet Soils

Phase: Sandy, Drainageway

1439—Crisfield sandy loam, rarely flooded

Map Unit Composition

Crisfield: 100 percent

Component Descriptions

Crisfield

MLRA: 80A - Central Rolling Red Prairies
Landform: Terrace on river valley
Parent material: Alluvium
Slope: 0 to 2 percent
Drainage class: Moderately well drained
Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: Low (About 4.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Rare
Depth to seasonal water saturation: About 40 to 73 inches
Runoff class: Negligible
Ecological site: Sandy Terrace (pe24-32)
Land capability (nonirrigated): 3s

Typical Profile:

H1—0 to 12 inches; sandy loam
 H2—12 to 24 inches; sandy loam
 H3—24 to 80 inches; coarse sand

Minor Components

Unnamed Wet Soils

Phase: Loamy, Depression

Ad—Albion-Shellabarger sandy loams, 2 to 4 percent slopes

Map Unit Composition

Albion: 60 percent
 Shellabarger: 40 percent

Component Descriptions

Albion

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 2 to 4 percent
Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: Low (About 5.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy (pe20-25)
Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 8 inches; sandy loam
 H2—8 to 16 inches; sandy loam
 H3—16 to 27 inches; loamy sand
 H4—27 to 60 inches; sand

Shellabarger

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 2 to 4 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 9.4 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy (pe20-25)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 14 inches; sandy loam
 H2—14 to 48 inches; sandy clay loam
 H3—48 to 60 inches; coarse sandy loam

Ae—Albion and Shellabarger Soils, 4 to 15 percent slopes

Map Unit Composition

Albion: 55 percent
 Shellabarger: 45 percent

Component Descriptions

Albion

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium

Slope: 4 to 15 percent
Drainage class: Well drained
Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: Low (About 5.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Sandy (pe20-25)
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 8 inches; sandy loam
 H2—8 to 16 inches; sandy loam
 H3—16 to 27 inches; loamy sand
 H4—27 to 60 inches; sand

Shellabarger

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 4 to 15 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 9.4 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Sandy (pe20-25)
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 14 inches; sandy loam
 H2—14 to 48 inches; sandy clay loam
 H3—48 to 60 inches; coarse sandy loam

Minor Components

Unnamed Wet Soils

Phase: Sandy, Drainageway

AED—Arents, Earthen Dam

AN—Albion-Shellabarger sandy loams, 4 to 15 percent slopes

Map Unit Composition

Albion: 65 percent

Shellabarger: 35 percent

Component Descriptions

Albion

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 4 to 15 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Low (About 5.9 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sandy (pe20-25)

Land capability (nonirrigated): 6e

Typical Profile:

- H1—0 to 8 inches; sandy loam
- H2—8 to 16 inches; sandy loam
- H3—16 to 28 inches; loamy sand
- H4—28 to 60 inches; sand

Shellabarger

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 4 to 15 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Moderate (About 8.8 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sandy (pe20-25)

Land capability (nonirrigated): 6e

Typical Profile:

- H1—0 to 11 inches; sandy loam
- H2—11 to 38 inches; sandy clay loam
- H3—38 to 60 inches; coarse sandy loam

As—Clairemont Soils, Saline, channeled

Map Unit Composition

Clairemont: 100 percent

Component Descriptions

Clairemont

MLRA: 78 - Central Rolling Red Plains

Landform: Flood plain

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Moderate (About 7.7 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Frequent

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Saline Lowland (pe20-25)

Land capability (nonirrigated): 6s

Typical Profile:

- H1—0 to 8 inches; silt loam
- H2—8 to 60 inches; loam

Minor Components

Unnamed Wet Soils

Phase: Sandy, Depression

Unnamed Wet Soils

Phase: Sandy, Drainageway

At—Attica loamy fine sand, 1 to 5 percent slopes

Map Unit Composition

Attica: 100 percent

Component Descriptions

Attica

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Eolian deposits

Slope: 1 to 5 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 7.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy (pe24-32)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 12 inches; loamy fine sand
 H2—12 to 30 inches; fine sandy loam
 H3—30 to 60 inches; loamy fine sand

Minor Components

Unnamed Wet Soils

Phase: Sandy, Depression

Ba—Blanket silt loam, 0 to 1 percent slopes

Map Unit Composition

Blanket: 100 percent

Component Descriptions

Blanket

MLRA: 80A - Central Rolling Red Prairies
Landform: Paleoterrace on river valley
Parent material: Clayey alluvium
Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: High (About 9.1 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Loamy Upland (pe20-25)
Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 9 inches; silt loam
 H2—9 to 30 inches; silty clay
 H3—30 to 60 inches; silty clay loam

Bb—Blanket silt loam, 1 to 3 percent slopes

Map Unit Composition

Blanket: 100 percent

Component Descriptions

Blanket

MLRA: 80A - Central Rolling Red Prairies
Landform: Paleoterrace on river valley
Parent material: Clayey alluvium
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: High (About 9.1 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Loamy Upland (pe20-25)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 9 inches; silt loam
 H2—9 to 30 inches; silty clay
 H3—30 to 60 inches; silty clay loam

Bc—Blanket silty clay loam, 1 to 3 percent slopes, eroded

Map Unit Composition

Blanket: 100 percent

Component Descriptions

Blanket

MLRA: 80A - Central Rolling Red Prairies
Landform: Paleoterrace on river valley
Parent material: Clayey alluvium
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 9.1 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Loamy Upland (pe20-25)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 9 inches; silty clay loam
 H2—9 to 30 inches; silty clay
 H3—30 to 60 inches; silty clay loam

Bf—Quinlan-Clairemont complex, 0 to 50 percent slopes

Map Unit Composition

Quinlan: 60 percent
 Clairemont: 40 percent

Component Descriptions

Quinlan

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 3 to 50 percent
Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)
Drainage class: Excessively drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very low (About 2.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Shallow Prairie (pe20-25)
Land capability (nonirrigated): 7e

Typical Profile:

H1—0 to 14 inches; loam
 Cr—14 to 14 inches;

Clairemont

MLRA: 78 - Central Rolling Red Plains
Landform: Flood plain

Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Loamy Lowland (pe20-25)
Land capability (nonirrigated): 5w

Typical Profile:

H1—0 to 14 inches; silt loam
 H2—14 to 60 inches; silt loam

Minor Components

Unnamed Wet Soils

Phase: Loamy, Drainageway

Ca—Canadian fine sandy loam, rarely flooded

Map Unit Composition

Canadian: 100 percent

Component Descriptions

Canadian

MLRA: 80A - Central Rolling Red Prairies
Landform: River valley, flood plain
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderately rapid (About 1.98 in/hr)
Available water capacity: Moderate (About 8.4 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Rare
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Sandy Terrace (pe24-32)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 18 inches; fine sandy loam
 H2—18 to 48 inches; fine sandy loam
 H3—48 to 60 inches; fine sandy loam

Cc—Case-Clark clay loams, 2 to 6 percent slopes**Map Unit Composition**

Case: 50 percent

Clark: 50 percent

Component Descriptions**Case***MLRA:* 80A - Central Rolling Red Prairies*Landform:* Paleoterrace on river valley*Parent material:* Alluvium*Slope:* 2 to 6 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 10.2 inches)*Shrink-swell potential:* Moderate (About 4.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Low*Ecological site:* Limy Upland (pe24-32)*Land capability (nonirrigated):* 4e*Typical Profile:*

H1—0 to 6 inches; clay loam

H2—6 to 60 inches; clay loam

Clark*MLRA:* 80A - Central Rolling Red Prairies*Landform:* Paleoterrace on river valley*Parent material:* Loamy alluvium*Slope:* 2 to 6 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 10.3 inches)*Shrink-swell potential:* Moderate (About 4.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Low*Ecological site:* Limy Upland (pe24-32)*Land capability (nonirrigated):* 4e*Typical Profile:*

H1—0 to 10 inches; clay loam

H2—10 to 60 inches; clay loam

Cd—Clairemont silt loam, occasionally flooded**Map Unit Composition**

Clairemont: 100 percent

Component Descriptions**Clairemont***MLRA:* 78 - Central Rolling Red Plains*Landform:* Flood plain*Parent material:* Alluvium*Slope:* 0 to 1 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 11.2 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Occasional*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Loamy Lowland (pe20-25)*Land capability (irrigated):* 2w*Land capability (nonirrigated):* 2w*Typical Profile:*

H1—0 to 14 inches; silt loam

H2—14 to 60 inches; silt loam

Minor Components**Unnamed Wet Soils***Phase:* Loamy, Drainageway**Cf—Clairemont soils, channeled****Map Unit Composition**

Clairemont: 100 percent

Component Descriptions**Clairemont***MLRA:* 78 - Central Rolling Red Plains*Landform:* Flood plain*Parent material:* Alluvium*Slope:* 0 to 2 percent*Drainage class:* Well drained

Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Loamy Lowland (pe20-25)
Land capability (nonirrigated): 5w

Typical Profile:
 H1—0 to 14 inches; silt loam
 H2—14 to 60 inches; silt loam

Minor Components
Unnamed Wet Soils
Phase: Loamy, Drainageway

Ck—Clark clay loam, 0 to 2 percent slopes

Map Unit Composition

Clark: 100 percent

Component Descriptions

Clark
MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 0 to 2 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.3 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Limy Upland (pe24-32)
Land capability (nonirrigated): 3e

Typical Profile:
 H1—0 to 10 inches; clay loam
 H2—10 to 60 inches; clay loam

Fa—Farnum fine sandy loam, 0 to 1 percent slopes

Map Unit Composition

Farnum: 100 percent

Component Descriptions

Farnum
MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 9.9 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy (pe24-32)
Land capability (irrigated): 1
Land capability (nonirrigated): 2e

Typical Profile:
 H1—0 to 9 inches; fine sandy loam
 H2—9 to 60 inches; clay loam

Minor Components
Unnamed Wet Soils
Phase: Loamy, Depression

Fm—Farnum loam, 0 to 1 percent slopes

Map Unit Composition

Farnum: 100 percent

Component Descriptions

Farnum
MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.4 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Loamy Upland (pe24-32)
Land capability (irrigated): 1
Land capability (nonirrigated): 2c

Typical Profile:
 H1—0 to 9 inches; loam
 H2—9 to 60 inches; clay loam

Minor Components
Unnamed Wet Soils
Phase: Loamy, Depression

Fr—Farnum loam, 1 to 3 percent slopes

Map Unit Composition

Farnum: 100 percent

Component Descriptions

Farnum
MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Alluvium
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.4 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe24-32)
Land capability (irrigated): 2e
Land capability (nonirrigated): 2e

Typical Profile:
 H1—0 to 9 inches; loam

H2—9 to 60 inches; clay loam

Fu—Farnum clay loam, 1 to 3 percent slopes, eroded

Map Unit Composition

Farnum: 100 percent

Component Descriptions

Farnum
MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Alluvium
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.4 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 3e

Typical Profile:
 H1—0 to 9 inches; clay loam
 H2—9 to 60 inches; clay loam

Ga—Grant silt loam, 0 to 1 percent slopes

Map Unit Composition

Grant: 100 percent

Component Descriptions

Grant
MLRA: 80A - Central Rolling Red Prairies
Landform: Terrace on upland
Parent material: Residuum
Slope: 0 to 1 percent
Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 1

Typical Profile:

H1—0 to 13 inches; silt loam
 H2—13 to 50 inches; silt loam
 H3—50 to 60 inches; silt loam

Gb—Grant silt loam, 1 to 3 percent slopes

Map Unit Composition

Grant: 100 percent

Component Descriptions

Grant

MLRA: 80A - Central Rolling Red Prairies
Landform: Terrace on upland
Parent material: Residuum
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 13 inches; silt loam
 H2—13 to 50 inches; silt loam
 H3—50 to 60 inches; silt loam

Gc—Grant silt loam, 3 to 6 percent slopes

Map Unit Composition

Grant: 100 percent

Component Descriptions

Grant

MLRA: 80A - Central Rolling Red Prairies
Landform: Terrace on upland
Parent material: Residuum
Slope: 3 to 6 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 13 inches; silt loam
 H2—13 to 50 inches; silt loam
 H3—50 to 60 inches; silt loam

GRP—Gravel Pits

INT—Aquolls

Ka—Kanza Soils, frequently flooded

Map Unit Composition

Kanza: 100 percent

Component Descriptions

Kanza

MLRA: 78 - Central Rolling Red Plains

Landform: Flood plain on river valley
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Poorly drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 4.5 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: About 0 to 36 inches
Runoff class: Negligible
Ecological site: Subirrigated (pe24-32)
Land capability (nonirrigated): 5w

Typical Profile:

H1—0 to 7 inches; loamy fine sand
 H2—7 to 48 inches; loamy fine sand

Kf—Kingfisher silt loam, 1 to 3 percent slopes

Map Unit Composition

Kingfisher: 100 percent

Component Descriptions

Kingfisher

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 1 to 3 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Low (About 5.5 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe20-25)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 9 inches; silt loam
 H2—9 to 17 inches; silty clay loam
 H3—17 to 28 inches; silty clay loam
 Cr—28 to 28 inches; weathered bedrock

Kv—Kingfisher-Vernon complex, 1 to 3 percent slopes

Map Unit Composition

Kingfisher: 70 percent
 Vernon: 30 percent

Component Descriptions

Kingfisher

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 1 to 3 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Low (About 5.5 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe20-25)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 9 inches; silt loam
 H2—9 to 17 inches; silty clay loam
 H3—17 to 28 inches; silty clay loam
 Cr—28 to 28 inches; weathered bedrock

Vernon

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 1 to 3 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: Low (About 3.5 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Red Clay Prairie (pe20-25)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 5 inches; clay loam
 H2—5 to 22 inches; silty clay
 H3—22 to 60 inches; weathered bedrock

Kz—Kingfisher-Vernon complex, 3 to 6 percent slopes

Map Unit Composition

Kingfisher: 60 percent
 Vernon: 40 percent

Component Descriptions

Kingfisher

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 3 to 6 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Low (About 5.5 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Loamy Upland (pe20-25)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 9 inches; silt loam
 H2—9 to 17 inches; silty clay loam
 H3—17 to 28 inches; silty clay loam
 Cr—28 to 28 inches; weathered bedrock

Vernon

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 3 to 6 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: Low (About 3.5 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Red Clay Prairie (pe20-25)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 5 inches; clay loam
 H2—5 to 22 inches; silty clay
 H3—22 to 60 inches; weathered bedrock

Ln—Lincoln Soils, frequently flooded

Map Unit Composition

Lincoln: 100 percent

Component Descriptions

Lincoln

MLRA: 78 - Central Rolling Red Plains
Landform: Flood plain on river valley
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Somewhat excessively drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 3.4 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: About 60 to 72 inches
Runoff class: Negligible
Ecological site: Sandy Lowland (pe20-25)
Land capability (nonirrigated): 6w

Typical Profile:

H1—0 to 6 inches; fine sandy loam
 H2—6 to 60 inches; stratified fine sand to clay loam

Minor Components

Kanza

Unnamed Wet Soils

Phase: Sandy, Depression

Unnamed Wet Soils

Phase: Sandy, Drainageway

LNN—Lincoln loamy sand, occasionally flooded

Map Unit Composition

Lincoln: 100 percent

Component Descriptions

Lincoln

MLRA: 80A - Central Rolling Red Prairies

Landform: Flood plain on river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat excessively drained

Slowest permeability: Rapid (About 6.00 in/hr)

Available water capacity: Low (About 3.3 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: About 60 to 72 inches

Runoff class: Negligible

Ecological site: Sandy Lowland (pe20-25)

Land capability (nonirrigated): 6w

Typical Profile:

H1—0 to 10 inches; loamy sand

H2—10 to 60 inches; sand

Minor Components

Kanza

Ma—Mangum clay, occasionally flooded

Map Unit Composition

Mangum: 100 percent

Component Descriptions

Mangum

MLRA: 78 - Central Rolling Red Plains

Landform: Flood plain

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: High (About 9.6 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Clay Lowland (pe20-25)

Land capability (irrigated): 3w

Land capability (nonirrigated): 3w

Typical Profile:

H1—0 to 10 inches; clay

H2—10 to 50 inches; clay

H3—50 to 60 inches; clay loam

Minor Components

Unnamed Wet Soils

Phase: Clayey, Depression

Mg—Mangum-Drummond complex, rarely flooded

Map Unit Composition

Mangum: 65 percent

Drummond: 35 percent

Component Descriptions

Mangum

MLRA: 78 - Central Rolling Red Plains

Landform: Flood plain

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: High (About 9.6 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Clay Lowland (pe20-25)

Land capability (irrigated): 2s

Land capability (nonirrigated): 3s

Typical Profile:

H1—0 to 10 inches; clay

H2—10 to 50 inches; clay

H3—50 to 60 inches; clay loam

Drummond

MLRA: 78 - Central Rolling Red Plains

Landform: Terrace on river valley
Parent material: Clayey and/or loamy alluvium
Slope: 0 to 1 percent
Drainage class: Moderately well drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Low (About 4.2 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: Rare
Depth to seasonal water saturation: About 24 to 72 inches
Runoff class: Very low
Ecological site: Saline Lowland (pe20-25)
Land capability (nonirrigated): 5s

Typical Profile:

H1—0 to 8 inches; silty clay loam
 H2—8 to 30 inches; clay loam
 H3—30 to 60 inches; variable

Minor Components

Unnamed Wet Soils

Phase: Clayey, Depression

Mn—Minco silt loam, 0 to 2 percent slopes

Map Unit Composition

Minco: 100 percent

Component Descriptions

Minco

MLRA: 80A - Central Rolling Red Prairies
Landform: Hillslope on upland
Parent material: Eolian deposits
Slope: 0 to 2 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 27 inches; silt loam
 H2—27 to 60 inches; silt loam

Na—Naron fine sandy loam, 0 to 1 percent slopes

Map Unit Composition

Naron: 100 percent

Component Descriptions

Naron

MLRA: 79 - Great Bend Sand Plains
Landform: Dune on paleoterrace on river valley
Parent material: Loamy eolian deposits
Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 9.0 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Sandy (pe24-32)
Land capability (irrigated): 1
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 12 inches; fine sandy loam
 H2—12 to 37 inches; fine sandy loam
 H3—37 to 60 inches; fine sandy loam

Minor Components

Unnamed Wet Soils

Phase: Loamy, Depression

Nb—Naron fine sandy loam, 1 to 3 percent slopes

Map Unit Composition

Naron: 100 percent

Component Descriptions

Naron

MLRA: 79 - Great Bend Sand Plains
Landform: Dune on paleoterrace on river valley
Parent material: Loamy eolian deposits
Slope: 1 to 3 percent

Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 9.0 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy (pe24-32)
Land capability (irrigated): 2e
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 12 inches; fine sandy loam
 H2—12 to 37 inches; fine sandy loam
 H3—37 to 60 inches; fine sandy loam

Minor Components**Unnamed Wet Soils**

Phase: Loamy, Depression

Unnamed Wet Soils

Phase: Loamy, Drainageway

Os—Ost clay loam, 0 to 1 percent slopes**Map Unit Composition**

Ost: 100 percent

Component Descriptions**Ost**

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: High (About 10.3 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 8 inches; clay loam
 H2—8 to 22 inches; clay loam
 H3—22 to 60 inches; clay loam

Minor Components**Carwile****Ot—Ost clay loam, 1 to 3 percent slopes****Map Unit Composition**

Ost: 100 percent

Component Descriptions**Ost**

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: High (About 10.3 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe24-32)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 8 inches; clay loam
 H2—8 to 22 inches; clay loam
 H3—22 to 60 inches; clay loam

Pa—Pond Creek silt loam, 0 to 1 percent slopes**Map Unit Composition**

Pond Creek: 100 percent

Component Descriptions

Pond Creek

MLRA: 80A - Central Rolling Red Prairies

Landform: Terrace

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 11.1 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Loamy Upland (pe24-32)

Land capability (nonirrigated): 1

Typical Profile:

H1—0 to 11 inches; silt loam

H2—11 to 60 inches; silty clay loam

Pd—Pond Creek silt loam, 1 to 3 percent slopes

Map Unit Composition

Pond Creek: 100 percent

Component Descriptions

Pond Creek

MLRA: 80A - Central Rolling Red Prairies

Landform: Terrace

Parent material: Alluvium

Slope: 1 to 3 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: High (About 11.1 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Loamy Upland (pe24-32)

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 11 inches; silt loam

H2—11 to 60 inches; silty clay loam

Minor Components

Unnamed Wet Soils

Phase: Loamy, Drainageway

Ph—Dale silt loam, rarely flooded

Map Unit Composition

Dale: 100 percent

Component Descriptions

Dale

MLRA: 80A - Central Rolling Red Prairies

Landform: Terrace on river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.8 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Loamy Terrace (pe24-32)

Land capability (nonirrigated): 1

Typical Profile:

H1—0 to 20 inches; silt loam

H2—20 to 60 inches; silt loam

Minor Components

Unnamed Wet Soils

Phase: Loamy, Depression

Pk—Buttermilk silt loam, rarely flooded

Map Unit Composition

Buttermilk: 100 percent

Component Descriptions

Buttermilk

MLRA: 80A - Central Rolling Red Prairies

Landform: Flood plain, river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 9.1 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: About 54 to 72 inches

Runoff class: Negligible

Ecological site: Saline Lowland (pe24-32)

Land capability (nonirrigated): 3s

Typical Profile:

H1—0 to 20 inches; silt loam

H2—20 to 32 inches; silt loam

H3—32 to 60 inches; silt loam

Minor Components

Unnamed Wet Soils

Phase: Loamy, Depression

Unnamed Wet Soils

Phase: Loamy, Drainageway

Ps—Pratt loamy fine sand, 5 to 10 percent slopes

Map Unit Composition

Pratt: 100 percent

Component Descriptions

Pratt

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 5 to 10 percent

Drainage class: Well drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Moderate (About 6.3 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sands (pe24-32)

Land capability (irrigated): 3e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 7 inches; loamy fine sand

H2—7 to 30 inches; loamy fine sand

H3—30 to 60 inches; loamy fine sand

Minor Components

Carwile

Unnamed Wet Soils

Phase: Sandy, Depression

Pt—Pratt-Tivoli loamy fine sands, 5 to 15 percent slopes

Map Unit Composition

Pratt: 70 percent

Tivoli: 30 percent

Component Descriptions

Pratt

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 5 to 15 percent

Drainage class: Well drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Moderate (About 6.3 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sands (pe20-25)

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 7 inches; loamy fine sand

H2—7 to 30 inches; loamy fine sand

H3—30 to 60 inches; loamy fine sand

Tivoli

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 5 to 15 percent
Drainage class: Excessively drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 3.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Sands (pe20-25)
Land capability (nonirrigated): 7e

Typical Profile:

H1—0 to 5 inches; loamy fine sand
 H2—5 to 60 inches; fine sand

Minor Components
Unnamed Wet Soils
Phase: Sandy, Depression

Qn—Quinlan loam, 1 to 3 percent slopes

Map Unit Composition

Quinlan: 100 percent

Component Descriptions

Quinlan
MLRA: 80A - Central Rolling Red Prairies
Landform: Hillslope on upland
Parent material: Residuum
Slope: 1 to 3 percent
Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very low (About 2.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Shallow Prairie (pe20-25)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 14 inches; loam
 Cr—14 to 14 inches; weathered bedrock

Minor Components
Unnamed Wet Soils
Phase: Loamy, Drainageway

Qw—Quinlan-Woodward complex, 5 to 15 percent slopes

Map Unit Composition

Quinlan: 60 percent
 Woodward: 40 percent

Component Descriptions

Quinlan
MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 5 to 15 percent
Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very low (About 2.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: High
Ecological site: Shallow Prairie (pe20-25)
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 14 inches; loam
 Cr—14 to 14 inches; weathered bedrock

Woodward
MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 5 to 15 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Moderate (About 6.8 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet
Runoff class: High
Ecological site: Loamy Upland (pe20-25)
Land capability (nonirrigated): 6e

Typical Profile:
 H1—0 to 40 inches; loam
 Cr—40 to 40 inches; weathered bedrock

Minor Components
Unnamed Wet Soils
Phase: Loamy, Drainageway

Rb—Knoco-Shale Outcrop complex, 15 to 80 percent slopes

Map Unit Composition

Knoco: 70 percent
 Rock outcrop: 30 percent

Component Descriptions

Knoco
MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Slope: 15 to 50 percent
Depth to restrictive feature: 4 to 19 inches to bedrock (paralithic)
Drainage class: Excessively drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Very low (About 1.5 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Red Shale (pe20-25)
Land capability (nonirrigated): 7s

Typical Profile:
 H1—0 to 9 inches; clay
 H2—9 to 19 inches;
 H3—19 to 60 inches;

Rock outcrop
MLRA: 78 - Central Rolling Red Plains
Slope: 15 to 80 percent
Depth to restrictive feature: 0 inches to bedrock (paralithic)
Drainage class: Excessively drained

Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Land capability (nonirrigated): 8s

Minor Components
Unnamed Wet Soils
Phase: Loamy, Drainageway

Sb—Shellabarger sandy loam, 3 to 6 percent slopes

Map Unit Composition

Shellabarger: 100 percent

Component Descriptions

Shellabarger
MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 3 to 6 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 9.4 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Sandy (pe20-25)
Land capability (nonirrigated): 3e

Typical Profile:
 H1—0 to 14 inches; sandy loam
 H2—14 to 48 inches; sandy clay loam
 H3—48 to 60 inches; coarse sandy loam

SBB—Shellabarger fine sandy loam, 0 to 1 percent slopes

Map Unit Composition

Shellabarger: 100 percent

Component Descriptions

Shellabarger

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Moderate (About 8.5 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Sandy (pe24-32)

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 11 inches; fine sandy loam

H2—11 to 34 inches; sandy clay loam

H3—34 to 60 inches; coarse sandy loam

Sc—Shellabarger sandy loam, 3 to 6 percent slopes, eroded

Map Unit Composition

Shellabarger: 100 percent

Component Descriptions

Shellabarger

MLRA: 79 - Great Bend Sand Plains

Landform: Paleoterrace on river valley

Parent material: Loamy alluvium

Slope: 3 to 6 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 9.4 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sandy (pe20-25)

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 14 inches; sandy loam

H2—14 to 48 inches; sandy clay loam

H3—48 to 60 inches; coarse sandy loam

Minor Components

Unnamed Wet Soils

Phase: Loamy, Drainageway

Tv—Tivoli fine sand, 5 to 20 percent slopes

Map Unit Composition

Tivoli: 100 percent

Component Descriptions

Tivoli

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 5 to 20 percent

Drainage class: Excessively drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Very low (About 3.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Choppy Sands (pe20-25)

Land capability (nonirrigated): 7e

Typical Profile:

H1—0 to 5 inches; fine sand

H2—5 to 60 inches; fine sand

Minor Components

Unnamed Wet Soils

Phase: Sandy, Depression

Vn—Vernon clay loam, 3 to 5 percent slopes

Map Unit Composition

Vernon: 100 percent

Component Descriptions

Vernon

MLRA: 78 - Central Rolling Red Plains

Landform: Hillslope on upland

Parent material: Residuum

Slope: 3 to 5 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: Low (About 3.5 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Red Clay Prairie (pe20-25)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 5 inches; clay loam

H2—5 to 22 inches; clay

H3—22 to 60 inches; weathered bedrock

Vr—Vernon clay loam, 5 to 15 percent slopes

Map Unit Composition

Vernon: 100 percent

Component Descriptions

Vernon

MLRA: 78 - Central Rolling Red Plains

Landform: Hillslope on upland

Parent material: Residuum

Slope: 5 to 15 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: Low (About 3.5 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: High

Ecological site: Red Clay Prairie (pe20-25)

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 5 inches; clay loam

H2—5 to 22 inches; clay

H3—22 to 60 inches; weathered bedrock

Minor Components

Unnamed Wet Soils

Phase: Clayey, Drainageway

Vs—Vernon-Knoco complex, 1 to 5 percent slopes

Map Unit Composition

Vernon: 50 percent

Knoco: 50 percent

Component Descriptions

Vernon

MLRA: 78 - Central Rolling Red Plains

Landform: Hillslope on upland

Parent material: Residuum

Slope: 1 to 5 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: Low (About 3.5 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Red Clay Prairie (pe20-25)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 5 inches; clay loam

H2—5 to 22 inches; silty clay

H3—22 to 60 inches; weathered bedrock

Knoco

MLRA: 78 - Central Rolling Red Plains

Landform: Hillslope on upland

Parent material: Dense, noncemented residuum weathered from claystone

Slope: 1 to 5 percent

Depth to restrictive feature: 4 to 19 inches to bedrock (paralithic)

Drainage class: Excessively drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Very low (About 1.5 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Red Shale (pe20-25)
Land capability (nonirrigated): 7s

Typical Profile:
 H1—0 to 9 inches; clay
 H2—9 to 19 inches; clay
 H3—19 to 60 inches;

W—Water

Wa—Waldeck sandy loam, occasionally flooded

Map Unit Composition

Waldeck: 100 percent

Component Descriptions

Waldeck

MLRA: 80A - Central Rolling Red Prairies
Landform: Flood plain on river valley
Parent material: Alluvium
Slope: 0 to 2 percent
Drainage class: Somewhat poorly drained
Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: Moderate (About 6.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Occasional
Depth to seasonal water saturation: About 24 to 48 inches
Runoff class: Very low
Ecological site: Subirrigated (pe24-32)
Land capability (nonirrigated): 3w

Typical Profile:
 H1—0 to 12 inches; sandy loam
 H2—12 to 36 inches; sandy loam
 H3—36 to 60 inches; sand

Minor Components

Kanza

Wo—Woodward-Quinlan loams, 0 to 3 percent slopes

Map Unit Composition

Woodward: 70 percent
 Quinlan: 30 percent

Component Descriptions

Woodward

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 1 to 3 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Moderate (About 6.8 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Shallow Prairie (pe20-25)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 40 inches; loam
 Cr—40 to 40 inches; weathered bedrock

Quinlan

MLRA: 78 - Central Rolling Red Plains
Landform: Hillslope on upland
Parent material: Residuum
Slope: 0 to 3 percent
Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very low (About 2.7 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe20-25)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 14 inches; loam
Cr—14 to 14 inches; weathered bedrock

Minor Components**Unnamed Wet Soils**

Phase: Loamy, Drainageway

Ws—Woodward-Quinlan loams, 3 to 6 percent slopes

Map Unit Composition

Woodward: 65 percent

Quinlan: 35 percent

Component Descriptions

Woodward

MLRA: 78 - Central Rolling Red Plains

Landform: Hillslope on upland

Parent material: Residuum

Slope: 3 to 6 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Moderate (About 6.8 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Shallow Prairie (pe20-25)

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 40 inches; loam
Cr—40 to 40 inches; weathered bedrock

Quinlan

MLRA: 78 - Central Rolling Red Plains

Landform: Hillslope on upland

Parent material: Residuum

Slope: 3 to 6 percent

Depth to restrictive feature: 10 to 20 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Very low (About 2.7 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Loamy Upland (pe20-25)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 14 inches; loam
Cr—14 to 14 inches; weathered bedrock

Minor Components**Unnamed Wet Soils**

Phase: Loamy, Drainageway

Ya—Yahola sandy loam, occasionally flooded

Map Unit Composition

Yahola: 100 percent

Component Descriptions

Yahola

MLRA: 78 - Central Rolling Red Plains

Landform: Flood plain

Parent material: Alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: High (About 9.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Subirrigated (pe20-25)

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 14 inches; sandy loam
H2—14 to 60 inches; fine sandy loam
H3—60 to 66 inches; stratified loam to loamy fine sand

Minor Components**Unnamed Wet Soils***Phase:* Loamy, Drainageway**Unnamed Wet Soils***Phase:* Loamy, Depression**Ze—Zenda clay loam, occasionally flooded****Map Unit Composition**

Zenda: 100 percent

Component Descriptions**Zenda***MLRA:* 78 - Central Rolling Red Plains*Landform:* Dune on paleoterrace on river valley*Parent material:* Sandy eolian deposits*Slope:* 0 to 1 percent*Drainage class:* Somewhat poorly drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 10.6 inches)*Shrink-swell potential:* Moderate (About 4.5 LEP)*Flooding hazard:* Occasional*Depth to seasonal water saturation:* About 24 to 48 inches*Runoff class:* Negligible*Ecological site:* Subirrigated (pe20-25)*Land capability (nonirrigated):* 2w*Typical Profile:*

H1—0 to 20 inches; clay loam

H2—20 to 60 inches; clay loam

Minor Components**Unnamed Wet Soils***Phase:* Loamy, Drainageway

Farmland Classification
Barber County, Kansas : Published

Prime farmland is one of several kinds of important farmland defined by the U.S. Department of Agriculture. It is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil qualities, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. It is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

A recent trend in land use in some parts of the survey area has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

The map units in the survey area that are considered prime farmland are listed in the following table. This list does not constitute a recommendation for a particular land use. On some soils included in the list, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures. The extent of each listed map unit is shown in the "Acres and Proportionate Extent of Soils" table. The location is shown on the detailed soil maps. The soil qualities that affect use and management are described in other tables in this document."

Map symbol	Mapunit name	Farmland Classification
033YE	Yahola fine sandy loam, occasionally flooded	All areas are prime farmland
077FU	Farnum loam, 3 to 6 percent slopes	All areas are prime farmland
077GE	Gerlane fine sandy loam, occasionally flooded	All areas are prime farmland
077MC	Minco silt loam, 0 to 1 percent slopes	All areas are prime farmland
077MN	Minco silt loam, 1 to 3 percent slopes	All areas are prime farmland
077MO	Minco silt loam, 3 to 6 percent slopes	All areas are prime farmland
077SE	Shellabarger fine sandy loam, 1 to 3 percent slopes	All areas are prime farmland
095FA	Farnum sandy loam, 0 to 2 percent slopes	All areas are prime farmland
095SB	Shellabarger sandy loam, 1 to 3 percent slopes	All areas are prime farmland
095WA	Waldeck fine sandy loam, occasionally flooded	All areas are prime farmland
097WA	Waldeck loam, occasionally flooded	All areas are prime farmland
151BC	Blanket silty clay loam, 1 to 4 percent slopes, eroded	All areas are prime farmland
151CM	Clark clay loam, 1 to 4 percent slopes	All areas are prime farmland
151CN	Clark fine sandy loam, 1 to 3 percent slopes	All areas are prime farmland
151CO	Clark-ost clay loams, 0 to 1 percent slopes	All areas are prime farmland
151FA	Farnum clay loam, 3 to 6 percent slopes, eroded	All areas are prime farmland
151OS	Ost clay loam, 1 to 4 percent slopes	All areas are prime farmland
151SE	Shellabarger fine sandy loam, 1 to 4 percent slopes	All areas are prime farmland
Ba	Blanket silt loam, 0 to 1 percent slopes	All areas are prime farmland
Bb	Blanket silt loam, 1 to 3 percent slopes	All areas are prime farmland
Bc	Blanket silty clay loam, 1 to 3 percent slopes, eroded	All areas are prime farmland
Ca	Canadian fine sandy loam, rarely flooded	All areas are prime farmland
Cc	Case-clark clay loams, 2 to 6 percent slopes	All areas are prime farmland
Cd	Clairemont silt loam, occasionally flooded	All areas are prime farmland
Ck	Clark clay loam, 0 to 2 percent slopes	All areas are prime farmland
Fa	Farnum fine sandy loam, 0 to 1 percent slopes	All areas are prime farmland
Fm	Farnum loam, 0 to 1 percent slopes	All areas are prime farmland
Fr	Farnum loam, 1 to 3 percent slopes	All areas are prime farmland
Fu	Farnum clay loam, 1 to 3 percent slopes, eroded	All areas are prime farmland
Ga	Grant silt loam, 0 to 1 percent slopes	All areas are prime farmland
Gb	Grant silt loam, 1 to 3 percent slopes	All areas are prime farmland
Gc	Grant silt loam, 3 to 6 percent slopes	All areas are prime farmland
Kf	Kingfisher silt loam, 1 to 3 percent slopes	All areas are prime farmland
Mn	Minco silt loam, 0 to 2 percent slopes	All areas are prime farmland
Na	Naron fine sandy loam, 0 to 1 percent slopes	All areas are prime farmland
Nb	Naron fine sandy loam, 1 to 3 percent slopes	All areas are prime farmland
Os	Ost clay loam, 0 to 1 percent slopes	All areas are prime farmland
Ot	Ost clay loam, 1 to 3 percent slopes	All areas are prime farmland
Pa	Pond creek silt loam, 0 to 1 percent slopes	All areas are prime farmland
Pd	Pond creek silt loam, 1 to 3 percent slopes	All areas are prime farmland
Ph	Dale silt loam, rarely flooded	All areas are prime farmland
Sb	Shellabarger sandy loam, 3 to 6 percent slopes	All areas are prime farmland
SBB	Shellabarger fine sandy loam, 0 to 1 percent slopes	All areas are prime farmland
Sc	Shellabarger sandy loam, 3 to 6 percent slopes, eroded	All areas are prime farmland
Wa	Waldeck sandy loam, occasionally flooded	All areas are prime farmland
Ze	Zenda clay loam, occasionally flooded	All areas are prime farmland
151AB	Albion sandy loam, 1 to 4 percent slopes	Prime farmland if irrigated
Ad	Albion-shellabarger sandy loams, 2 to 4 percent slopes	Prime farmland if irrigated

The "Soil Rating for Plant Growth, modified 1998" (SRPG) is a relative rating of the capacity of a soil to produce a specific plant under a defined management system. The index is determined from yield data on a few benchmark soils and is used to calculate yields, the net returns from crops, land assessment values, and taxes and to perform risk analysis when land management decisions are made. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

Map symbol	Soil name	Crop Index
033CP	Clairemont Loam, Channeled-----	41
033OA	Obaro Silty Clay Loam, 5 To 12 Percent Slopes-----	43
033OB	Obaro-Rock Outcrop Complex, 10 To 30 Percent Slopes-----	34
033QR	Quinlan-Woodward Loams, 6 To 15 Percent Slopes-----	19
033QT	Quinlan-Woodward Loams, 15 To 30 Percent Slopes-----	14
033YE	Yahola Fine Sandy Loam, Occasionally Flooded-----	44
077BP	Woodward-Port Complex, 0 To 20 Percent Slopes-----	39
077FU	Farnum Loam, 3 To 6 Percent Slopes-----	71
077GE	Gerlane Fine Sandy Loam, Occasionally Flooded-----	58
077MC	Minco Silt Loam, 0 To 1 Percent Slopes-----	69
077MN	Minco Silt Loam, 1 To 3 Percent Slopes-----	67
077MO	Minco Silt Loam, 3 To 6 Percent Slopes-----	65
077QU	Quinlan Loam, 3 To 6 Percent Slopes-----	4
077SE	Shellabarger Fine Sandy Loam, 1 To 3 Percent Slopes-----	69
077WE	Woodward-Quinlan Loams, 1 To 3 Percent Slopes-----	16
095AC	Albion Sandy Loam, 3 To 6 Percent Slopes-----	41
095DA	Dillwyn-Plevna Complex, Occasionally Flooded-----	31
095FA	Farnum Sandy Loam, 0 To 2 Percent Slopes-----	71
095PB	Pratt Loamy Fine Sand, 1 To 5 Percent Slopes-----	38
095SB	Shellabarger Sandy Loam, 1 To 3 Percent Slopes-----	71
095WA	Waldeck Fine Sandy Loam, Occasionally Flooded-----	42
097LN	Lincoln Sandy Loam, Occasionally Flooded-----	23
097QW	Quinlan-Woodward Loams, 6 To 25 Percent Slopes-----	16
097WA	Waldeck Loam, Occasionally Flooded-----	43
1439	Crisfield Sandy Loam, Rarely Flooded-----	30
151AB	Albion Sandy Loam, 1 To 4 Percent Slopes-----	42
151AO	Albion Sandy Loam, 3 To 7 Percent Slopes, Eroded-----	41
151AS	Albion And Shellabarger Soils, 7 To 15 Percent Slopes-----	46
151BC	Blanket Silty Clay Loam, 1 To 4 Percent Slopes, Eroded-----	64
151CA	Carwile Fine Sandy Loam, 0 To 1 Percent Slopes-----	21
151CK	Case-Clark Complex, 7 To 15 Percent Slopes-----	35
151CM	Clark Clay Loam, 1 To 4 Percent Slopes-----	33
151CN	Clark Fine Sandy Loam, 1 To 3 Percent Slopes-----	34
151CO	Clark-Ost Clay Loams, 0 To 1 Percent Slopes-----	35
151FA	Farnum Clay Loam, 3 To 6 Percent Slopes, Eroded-----	67
151KP	Kanza-Plevna Complex, Frequently Flooded-----	27
151OS	Ost Clay Loam, 1 To 4 Percent Slopes-----	38
151PM	Pratt Loamy Fine Sand, 3 To 8 Percent Slopes-----	37
151PN	Pratt Loamy Fine Sand, 8 To 12 Percent Slopes-----	34
151PO	Pratt-Carwile Complex, 0 To 5 Percent Slopes-----	30
151SA	Albion-Kaski Complex, 0 To 15 Percent Slopes-----	46
151SE	Shellabarger Fine Sandy Loam, 1 To 4 Percent Slopes-----	64
AED	Arents, Earthen Dam-----	0
AN	Albion-Shellabarger Sandy Loams, 4 To 15 Percent Slopes-----	43
Ad	Albion-Shellabarger Sandy Loams, 2 To 4 Percent Slopes-----	51
Ae	Albion And Shellabarger Soils, 4 To 15 Percent Slopes-----	47
As	Clairemont Soils, Saline, Channeled-----	29
At	Attica Loamy Fine Sand, 1 To 5 Percent Slopes-----	49
Ba	Blanket Silt Loam, 0 To 1 Percent Slopes-----	66
Bb	Blanket Silt Loam, 1 To 3 Percent Slopes-----	65
Bc	Blanket Silty Clay Loam, 1 To 3 Percent Slopes, Eroded-----	64
Bf	Quinlan-Clairemont Complex, 0 To 50 Percent Slopes-----	18
Ca	Canadian Fine Sandy Loam, Rarely Flooded-----	54
Cc	Case-Clark Clay Loams, 2 To 6 Percent Slopes-----	39
Cd	Clairemont Silt Loam, Occasionally Flooded-----	52
Cf	Clairemont Soils, Channeled-----	41
Ck	Clark Clay Loam, 0 To 2 Percent Slopes-----	34
Fa	Farnum Fine Sandy Loam, 0 To 1 Percent Slopes-----	68
Fm	Farnum Loam, 0 To 1 Percent Slopes-----	72
Fr	Farnum Loam, 1 To 3 Percent Slopes-----	70
Fu	Farnum Clay Loam, 1 To 3 Percent Slopes, Eroded-----	70
GRP	Gravel Pits-----	0
Ga	Grant Silt Loam, 0 To 1 Percent Slopes-----	69
Gb	Grant Silt Loam, 1 To 3 Percent Slopes-----	68
Gc	Grant Silt Loam, 3 To 6 Percent Slopes-----	66
INT	Aquolls-----	12
Ka	Kanza Soils, Frequently Flooded-----	26
Kf	Kingfisher Silt Loam, 1 To 3 Percent Slopes-----	41
Kv	Kingfisher-Vernon Complex, 1 To 3 Percent Slopes-----	34
Kz	Kingfisher-Vernon Complex, 3 To 6 Percent Slopes-----	31
LNN	Lincoln Loamy Sand, Occasionally Flooded-----	22
Ln	Lincoln Soils, Frequently Flooded-----	17
Ma	Mangum Clay, Occasionally Flooded-----	45
Mg	Mangum-Drummond Complex, Rarely Flooded-----	41
Mn	Minco Silt Loam, 0 To 2 Percent Slopes-----	65
Na	Naron Fine Sandy Loam, 0 To 1 Percent Slopes-----	67
Nb	Naron Fine Sandy Loam, 1 To 3 Percent Slopes-----	66
Os	Ost Clay Loam, 0 To 1 Percent Slopes-----	50
Ot	Ost Clay Loam, 1 To 3 Percent Slopes-----	50
Pa	Pond Creek Silt Loam, 0 To 1 Percent Slopes-----	73
Pd	Pond Creek Silt Loam, 1 To 3 Percent Slopes-----	72

The "Soil Rating for Plant Growth, modified 1998" (SRPG) is a relative rating of the capacity of a soil to produce a specific plant under a defined management system. The index is determined from yield data on a few benchmark soils and is used to calculate yields, the net returns from crops, land assessment values, and taxes and to perform risk analysis when land management decisions are made. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

Map symbol	Soil name	Crop Index
Ph	Dale Silt Loam, Rarely Flooded-----	64
Pk	Buttermilk Silt Loam, Rarely Flooded-----	44
Ps	Pratt Loamy Fine Sand, 5 To 10 Percent Slopes-----	35
Pt	Pratt-Tivoli Loamy Fine Sands, 5 To 15 Percent Slopes-----	29
Qn	Quinlan Loam, 1 To 3 Percent Slopes-----	8
Qw	Quinlan-Woodward Complex, 5 To 15 Percent Slopes-----	25
Rb	Knoco-Shale Outcrop Complex, 15 To 80 Percent Slopes-----	1
SBB	Shellabarger Fine Sandy Loam, 0 To 1 Percent Slopes-----	66
Sb	Shellabarger Sandy Loam, 3 To 6 Percent Slopes-----	65
Sc	Shellabarger Sandy Loam, 3 To 6 Percent Slopes, Eroded-----	65
Tv	Tivoli Fine Sand, 5 To 20 Percent Slopes-----	18
Vn	Vernon Clay Loam, 3 To 5 Percent Slopes-----	17
Vr	Vernon Clay Loam, 5 To 15 Percent Slopes-----	15
Vs	Vernon-Knoco Complex, 1 To 5 Percent Slopes-----	16
W	Water-----	0
Wa	Waldeck Sandy Loam, Occasionally Flooded-----	40
Wo	Woodward-Quinlan Loams, 0 To 3 Percent Slopes-----	44
Ws	Woodward-Quinlan Loams, 3 To 6 Percent Slopes-----	40
Ya	Yahola Sandy Loam, Occasionally Flooded-----	44
Ze	Zenda Clay Loam, Occasionally Flooded-----	58

Barber County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
033CP:CLAIREMONT	100	N/A	6w	Not prime farmland	B	Loamy Lowland (pe20-25)	1K	.43	.43	5	4L	86
033OA:OBARO-----	100	N/A	6e	Not prime farmland	B	Loamy Upland (pe20-25)		.43	.43	3	4L	86
033OB:OBARO-----	75	N/A	6e	Not prime farmland	B	Loamy Upland (pe20-25)		.43	.43	3	4L	86
033OB:ROCK OUTCROP-----	25	N/A	8	Not prime farmland	D	Unspecified		---	---	-	---	---
033QR:QUINLAN---	55	N/A	6e	Not prime farmland	C	Shallow Prairie (pe20-25)	10	.37	.37	2	4L	86
033QR:WOODWARD--	45	N/A	6e	Not prime farmland	B	Loamy Upland (pe20-25)	8	.37	.37	3	4L	86
033QT:QUINLAN---	55	N/A	7e	Not prime farmland	C	Shallow Prairie (pe20-25)	10	.37	.37	2	4L	86
033QT:WOODWARD--	45	N/A	6e	Not prime farmland	B	Loamy Upland (pe20-25)	8	.37	.37	3	4L	86
033YE:YAHOLA----	100	N/A	2w	All areas are prime farmland	B	Sandy Lowland (pe20-25)	1K	.20	.20	5	3	86
077BP:WOODWARD--	65	N/A	6e	Not prime farmland	B	Loamy Upland (pe24-32)	8	.37	.37	3	4L	86
077BP:PORT-----	35	N/A	5w	Not prime farmland	B	Loamy Lowland (pe24-32)	1	.37	.37	5	6	48
077FU:FARNUM----	100	N/A	3e	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.28	.28	5	6	48
077GE:GERLANE---	100	N/A	2e	All areas are prime farmland	B	Subirrigated (pe24-32)	1	.20	.20	5	3	86
077MC:MINCO-----	100	N/A	1	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.37	.37	5	5	56
077MN:MINCO-----	100	N/A	2e	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.37	.37	5	5	56
077MO:MINCO-----	100	N/A	3e	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.37	.37	5	5	56
077QU:QUINLAN---	100	N/A	4e	Not prime farmland	C	Shallow Prairie (pe24-32)	10	.37	.37	2	4L	86
077SE:SHELLABARG ER-----	100	N/A	2e	All areas are prime farmland	B	Sandy (pe24-32)	5	.20	.20	5	3	86

Barber County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
077WE:WOODWARD--	50	N/A	2e	Not prime farmland	B	Loamy Upland (pe24-32)	8	.37	.37	3	4L	86
077WE:QUINLAN---	50	N/A	3e	Not prime farmland	C	Shallow Prairie (pe24-32)	10	.37	.37	2	4L	86
095AC:ALBION----	100	N/A	4e	Not prime farmland	B	Sandy (pe24-32)	6G	.20	.20	4	3	86
095DA:DILLWYN---	60	N/A	4w	Not prime farmland	A	Subirrigated (pe24-32)	1	.17	.17	5	2	134
095DA:PLEVNA----	40	N/A	5w	Not prime farmland	D	Subirrigated (pe24-32)	2	.20	.20	5	3	86
095FA:FARNUM----	100	2e-	3e	All areas are prime farmland	B	Sandy (pe24-32)	3	.20	.20	5	3	86
095PB:PRATT-----	100	3e-	4e	Not prime farmland	A	Sands (pe24-32)	7	.17	.17	5	2	134
095SB:SHELLABARG ER-----	100	N/A	2e	All areas are prime farmland	B	Sandy (pe24-32)	5	.20	.20	5	3	86
095WA:WALDECK---	100	N/A	3w	All areas are prime farmland	C	Subirrigated (pe24-32)	1K	.20	.20	4	3	86
097LN:LINCOLN---	100	N/A	6w	Not prime farmland	A	Sandy Lowland (pe20-25)	1K	.20	.20	5	3	86
097QW:QUINLAN---	55	N/A	6e	Not prime farmland	C	Shallow Prairie (pe20-25)	10	.37	.37	2	4L	86
097QW:WOODWARD--	45	N/A	6e	Not prime farmland	B	Loamy Upland (pe20-25)	8	.37	.37	3	4L	86
097WA:WALDECK---	100	N/A	3w	All areas are prime farmland	C	Subirrigated (pe20-25)	1K	.28	.28	4	5	56
1439:CRISFIELD--	100	N/A	3s	Not prime farmland	B	Sandy Terrace (pe24-32)	5	.15	.15	5	3	86
151AB:ALBION----	100	N/A	3e	Prime farmland if irrigated	B	Sandy (pe24-32)	6G	.20	.20	4	3	86
151AO:ALBION----	100	N/A	4e	Not prime farmland	B	Sandy (pe24-32)	6G	.20	.20	4	3	86
151AS:ALBION----	50	N/A	6e	Not prime farmland	B	Sandy (pe24-32)	6G	.20	.20	4	3	86
151AS:SHELLABARG ER-----	50	N/A	6e	Not prime farmland	B	Sandy (pe24-32)	5	.20	.20	5	3	86
151BC:BLANKET---	100	N/A	3e	All areas are prime farmland	C	Loamy Upland (pe21-28)	4C	.37	.37	5	6	48

Barber County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
151CA:CARWILE---	100	N/A	2w	Not prime farmland	D	Sandy (pe21-28)	1	.24	.24	5	3	86
151CK:CASE-----	60	N/A	6e	Not prime farmland	B	Limy Upland (pe24-32)	8	.32	.32	5	4L	86
151CK:CLARK-----	40	N/A	6e	Not prime farmland	B	Limy Upland (pe24-32)	8	.28	.28	5	4L	86
151CM:CLARK-----	100	N/A	3e	All areas are prime farmland	B	Limy Upland (pe21-28)	8	.28	.28	5	4L	86
151CN:CLARK-----	100	N/A	3e	All areas are prime farmland	B	Limy Upland (pe24-32)	8	.20	.20	5	3	86
151CO:CLARK-----	70	N/A	2c	All areas are prime farmland	B	Limy Upland (pe24-32)	8	.28	.28	5	4L	86
151CO:OST-----	30	N/A	2c	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.32	.32	5	6	48
151FA:FARNUM----	100	N/A	4e	All areas are prime farmland	B	Loamy Upland (pe21-28)	3	.28	.28	5	6	48
151KP:KANZA-----	50	N/A	5w	Not prime farmland	D	Unspecified	2	.17	.17	5	2	134
151KP:PLEVNA----	50	N/A	5w	Not prime farmland	D	Subirrigated (pe21-28)	2	.20	.20	5	3	86
151OS:OST-----	100	N/A	2e	All areas are prime farmland	B	Loamy Upland (pe21-28)	3	.32	.32	5	6	48
151PM:PRATT-----	100	3e-	4e	Not prime farmland	A	Sands (pe21-28)	7	.17	.17	5	2	134
151PN:PRATT-----	100	N/A	6e	Not prime farmland	A	Sands (pe21-28)	7	.17	.17	5	2	134
151PO:PRATT-----	60	3e-	4e	Not prime farmland	A	Sands (pe21-28)	7	.17	.17	5	2	134
151PO:CARWILE---	40	N/A	2w	Not prime farmland	D	Sandy (pe21-28)	1	.24	.24	5	3	86
151SA:ALBION----	70	N/A	6e	Not prime farmland	B	Sandy (pe21-28)	6G	.20	.20	4	3	86
151SA:KASKI-----	30	N/A	2w	Not prime farmland	B	Loamy Lowland (pe21-28)	1K	.28	.28	5	6	48
151SE:SHELLABARG ER-----	100	N/A	2e	All areas are prime farmland	B	Sandy (pe21-28)	5	.20	.20	5	3	86

Barber County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
AED:ARENTS, EARTHEN DAM----	100	N/A	8	Not prime farmland		Unspecified		---	---	-	---	---
AN:ALBION-----	65	N/A	6e	Not prime farmland	B	Sandy (pe20-25)	6G	.20	.20	4	3	86
AN:SHELLABARGER--	35	N/A	6e	Not prime farmland	B	Sandy (pe20-25)	5	.20	.20	5	3	86
Ad:ALBION-----	60	N/A	4e	Prime farmland if irrigated	B	Sandy (pe20-25)	6G	.20	.24	4	3	86
Ad:SHELLABARGER--	40	N/A	3e	Prime farmland if irrigated	B	Sandy (pe20-25)	5	.20	.20	5	3	86
Ae:ALBION-----	55	N/A	6e	Not prime farmland	B	Sandy (pe20-25)	6G	.20	.20	4	3	86
Ae:SHELLABARGER--	45	N/A	6e	Not prime farmland	B	Sandy (pe20-25)	5	.20	.20	5	3	86
As:CLAIREMONT---	100	N/A	6s	Not prime farmland	B	Saline Lowland (pe20-25)	1K	.43	.43	5	4L	86
At:ATTICA-----	100	N/A	3e	Not prime farmland	B	Sandy (pe24-32)	5	.17	.17	5	2	134
Ba:BLANKET-----	100	N/A	2c	All areas are prime farmland	C	Loamy Upland (pe20-25)	4C	.37	.37	5	5	56
Bb:BLANKET-----	100	N/A	2e	All areas are prime farmland	C	Loamy Upland (pe20-25)	4C	.37	.37	5	5	56
Bc:BLANKET-----	100	N/A	3e	All areas are prime farmland	C	Loamy Upland (pe20-25)	4C	.37	.37	5	6	48
Bf:QUINLAN-----	60	N/A	7e	Not prime farmland	C	Shallow Prairie (pe20-25)	10	.37	.37	2	4L	86
Bf:CLAIREMONT---	40	N/A	5w	Not prime farmland	B	Loamy Lowland (pe20-25)	1K	.43	.43	5	4L	86
Ca:CANADIAN-----	100	N/A	2e	All areas are prime farmland	B	Sandy Terrace (pe24-32)	1	.20	.20	5	3	86
Cc:CASE-----	50	N/A	4e	All areas are prime farmland	B	Limy Upland (pe24-32)	8	.32	.32	5	4L	86
Cc:CLARK-----	50	N/A	4e	All areas are prime farmland	B	Limy Upland (pe24-32)	8	.28	.28	5	4L	86
Cd:CLAIREMONT---	100	2w-	2w	All areas are prime farmland	B	Loamy Lowland (pe20-25)	1K	.43	.43	5	4L	86

Barber County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Cf:CLAIREMONT---	100	N/A	5w	Not prime farmland	B	Loamy Lowland (pe20-25)	1K	.43	.43	5	4L	86
Ck:CLARK-----	100	N/A	3e	All areas are prime farmland	B	Limy Upland (pe24-32)	8	.28	.28	5	4L	86
Fa:FARNUM-----	100	1-	2e	All areas are prime farmland	B	Sandy (pe24-32)	3	.20	.20	5	3	86
Fm:FARNUM-----	100	1-	2c	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.28	.28	5	6	48
Fr:FARNUM-----	100	2e-	2e	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.28	.28	5	6	48
Fu:FARNUM-----	100	N/A	3e	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.28	.28	5	6	48
GRP:GRAVEL PITS-	100	N/A	N/A	Not prime farmland		Unspecified		---	---	-	---	---
Ga:GRANT-----	100	N/A	1	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.37	.37	4	5	56
Gb:GRANT-----	100	N/A	2e	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.37	.37	4	5	56
Gc:GRANT-----	100	N/A	3e	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.37	.37	4	5	56
INT:AQUOLLS-----	100	N/A	5w	Not prime farmland	C	Unspecified		---	---	-	---	0
Ka:KANZA-----	100	N/A	5w	Not prime farmland	D	Subirrigated (pe24-32)	2	.17	.17	5	2	134
Kf:KINGFISHER---	100	N/A	2e	All areas are prime farmland	B	Loamy Upland (pe20-25)	6D	.37	.37	3	5	56
Kv:KINGFISHER---	70	N/A	2e	Not prime farmland	B	Loamy Upland (pe20-25)	6D	.37	.37	3	5	56
Kv:VERNON-----	30	N/A	3e	Not prime farmland	D	Red Clay Prairie (pe20-25)		.37	.37	3	4	86
Kz:KINGFISHER---	60	N/A	3e	Not prime farmland	B	Loamy Upland (pe20-25)	6D	.37	.37	3	5	56
Kz:VERNON-----	40	N/A	4e	Not prime farmland	D	Red Clay Prairie (pe20-25)		.37	.37	3	4	86
LNN:LINCOLN-----	100	N/A	6w	Not prime farmland	A	Sandy Lowland (pe20-25)	1K	.17	.17	5	2	134

Barber County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Ln:LINCOLN-----	100	N/A	6w	Not prime farmland	A	Sandy Lowland (pe20-25)	1K	.20	.20	5	3	86
Ma:MANGUM-----	100	3w-	3w	Not prime farmland	D	Clay Lowland (pe20-25)	1K	.32	.32	5	4	86
Mg:MANGUM-----	65	2s-	3s	Not prime farmland	D	Clay Lowland (pe20-25)	1K	.32	.32	5	4	86
Mg:DRUMMOND-----	35	N/A	5s	Not prime farmland	D	Saline Lowland (pe20-25)	9W	.43	.43	2	4L	86
Mn:MINCO-----	100	N/A	2e	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.37	.37	5	5	56
Na:NARON-----	100	1-	2e	All areas are prime farmland	B	Sandy (pe24-32)	5	.20	.20	5	3	86
Nb:NARON-----	100	2e-	3e	All areas are prime farmland	B	Sandy (pe24-32)	5	.20	.20	5	3	86
Os:OST-----	100	N/A	2c	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.32	.32	5	6	48
Ot:OST-----	100	N/A	2e	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.32	.32	5	6	48
Pa:POND CREEK---	100	N/A	1	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.37	.37	5	6	48
Pd:POND CREEK---	100	N/A	2e	All areas are prime farmland	B	Loamy Upland (pe24-32)	3	.37	.37	5	6	48
Ph:DALE-----	100	N/A	1	All areas are prime farmland	B	Loamy Terrace (pe24-32)	1	.37	.37	5	5	56
Pk:BUTTERMILK---	100	N/A	3s	Not prime farmland	B	Saline Lowland (pe24-32)	9L	.32	.32	5	5	56
Ps:PRATT-----	100	3e-	4e	Not prime farmland	A	Sands (pe24-32)	7	.17	.17	5	2	134
Pt:PRATT-----	70	N/A	6e	Not prime farmland	A	Sands (pe20-25)	7	.17	.17	5	2	134
Pt:TIVOLI-----	30	N/A	7e	Not prime farmland	A	Sands (pe20-25)	7	.17	.17	5	2	134
Qn:QUINLAN-----	100	N/A	3e	Not prime farmland	C	Shallow Prairie (pe20-25)	10	.37	.37	2	4L	86
Qw:QUINLAN-----	60	N/A	6e	Not prime farmland	C	Shallow Prairie (pe20-25)	10	.37	.37	2	4L	86

Barber County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Qw:WOODWARD----	40	N/A	6e	Not prime farmland	B	Loamy Upland (pe20-25)	8	.37	.37	3	4L	86
Rb:KNOCO-----	70	N/A	7s	Not prime farmland	D	Red Shale (pe20- 25)		.37	.37	2	4	86
Rb:ROCK OUTCROP-	30	N/A	8s	Not prime farmland	D	Unspecified		---	---	-	---	0
SBB:SHELLABARGER	100	N/A	2e	All areas are prime farmland	B	Sandy (pe24-32)	5	.20	.20	5	3	86
Sb:SHELLABARGER-	100	N/A	3e	All areas are prime farmland	B	Sandy (pe20-25)	5	.20	.20	5	3	86
Sc:SHELLABARGER-	100	N/A	3e	All areas are prime farmland	B	Sandy (pe20-25)	5	.20	.20	5	3	86
Tv:TIVOLI-----	100	N/A	7e	Not prime farmland	A	Choppy Sands (pe20-25)	7	.17	.17	5	1	250
Vn:VERNON-----	100	N/A	4e	Not prime farmland	D	Red Clay Prairie (pe20-25)	4C	.37	.37	3	4	86
Vr:VERNON-----	100	N/A	6e	Not prime farmland	D	Red Clay Prairie (pe20-25)	4C	.37	.37	4	4	86
Vs:VERNON-----	50	N/A	4e	Not prime farmland	D	Red Clay Prairie (pe20-25)	4C	.37	.37	4	4	86
Vs:KNOCO-----	50	N/A	7s	Not prime farmland	D	Red Shale (pe20- 25)		.37	.37	2	4	86
W:WATER-----	100	N/A	N/A			Unspecified		---	---	-	---	---
Wa:WALDECK-----	100	N/A	3w	All areas are prime farmland	C	Subirrigated (pe24-32)	1K	.20	.20	4	3	86
Wo:WOODWARD----	70	N/A	2e	Not prime farmland	B	Shallow Prairie (pe20-25)	8	.37	.37	3	4L	86
Wo:QUINLAN-----	30	N/A	3e	Not prime farmland	C	Loamy Upland (pe20-25)	10	.37	.37	2	4L	86
Ws:WOODWARD----	65	N/A	3e	Not prime farmland	B	Shallow Prairie (pe20-25)	8	.37	.37	3	4L	86
Ws:QUINLAN-----	35	N/A	4e	Not prime farmland	C	Loamy Upland (pe20-25)	10	.37	.37	2	4L	86
Ya:YAHOLA-----	100	N/A	2w	Not prime farmland	B	Subirrigated (pe20-25)	1K	.20	.20	5	3	86
Ze:ZENDA-----	100	N/A	2w	All areas are prime farmland	C	Subirrigated (pe20-25)	1	.28	.28	5	6	48

RANGELAND PRODUCTIVITY
Barber County, Kansas

Use and Explanation of Rangeland, Grazed Forest Land, Native Pastureland Interpretations

Information in this subsection can be used to plan the use and management of soils for rangeland, grazed forest land, and native pasture. Different kinds of soils vary in their capacity to produce native grasses and other plants suitable for grazing. Information in this subsection provides groupings of similar soils and estimates of potential forage production, which can be used to determine livestock stocking rates.

Rangeland. Range is land on which the native vegetation (climax or natural potential plant community) is predominantly grasses, grasslike plants, forbs, and shrubs suitable for grazing and browsing. Range includes natural grasslands, savannas, many wetlands, some deserts, tundra, and certain shrub and forb communities. Rangeland receives no regular or frequent cultural treatment. The composition and production of the plant community are determined by soil, climate, topography, overstory canopy, and grazing management.

Grazed Forest Land. Includes land on which the understory includes, as an integral part of the forest plant community, plants that can be grazed without significantly impairing other forest values.

Native Pasture. Includes land on which the native vegetation (climax or natural potential plant community) is forest but which is used and managed primarily for production of native plants for forage. Native pasture includes cut-over forest land and forest land cleared and now managed for native or naturalized forage plants.

Rangeland

In areas that have similar climate and topography, differences in the kind and amount of vegetation produced on rangeland are closely related to the kind of soil. Effective management based on the relationship between the soils and vegetation and water.

The Rangeland, Grazed Forest land, Native Pastureland Interpretations shows, for each soil that supports rangeland vegetation, the ecological site and the potential annual production of vegetation in favorable, normal, unfavorable years. An explanation of the column headings in this table follows.

An ecological site is the product of all the environmental factors responsible for its development. It has characteristic soils that have developed over time throughout the soil development process; a characteristic hydrology, particularly infiltration and runoff, that has developed over time; and a characteristic plant community (kind and amount of vegetation). The hydrology of a site is influenced by development of the soil and plant community. The vegetation, soils, and hydrology are all interrelated. Each is influenced by the others and influences the development of the others. The plant community on an ecological site is typified by an association of species that differs from that of other ecological sites in the kind and/or proportion of species or in total production. Descriptions of ecological sites are provided in the Field Office Technical Guide, which is available in local offices of the Natural Resources Conservation Service.

Total dry-weight production is the amount of vegetation that can be expected to grow annually on well managed rangeland that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, average, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are about average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Yields are adjusted to a common percent of air-dry moisture content.

Range management requires a knowledge of the kinds of soil and of the potential natural plant community. It also requires an evaluation of the present range similarity index and rangeland trend. Range similarity index is determined by comparing the present plant community with the potential natural plant community on a particular rangeland ecological site. The more closely the existing community resembles the potential community, the higher the range similarity index. Rangeland trend is defined as the direction of change in an existing plant community relative to the potential natural plant community. Further information about the range similarity index and rangeland trend is available in chapter 4 of the National Range and Pasture Handbook, which is available in local offices of the Natural Resources Conservation Service. The objective in range management is to control grazing so that the plants growing on a site are about the same in kind and amount as the potential natural plant community for that site. Such management generally results in the optimum production of vegetation, control of undesirable brush species, conservation of water, and control of erosion. Sometimes, however, an area with a range similarity index somewhat below the potential meets grazing needs, provides wildlife habitat, and protects soil and water resources.

RANGELAND PRODUCTIVITY--Continued
Barber County, Kansas

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
033CP: Clairemont-----	Loamy Lowland (pe20-25)	3,400	2,600	1,800
033OA: Obaro-----	Loamy Upland (pe20-25)	2,400	1,800	1,200
033OB: Obaro-----	Loamy Upland (pe20-25)	2,400	1,800	1,200
Rock Outcrop-----	---	---	---	---
033QR: Quinlan-----	Shallow Prairie (pe20-25)	2,500	1,800	1,300
Woodward-----	Loamy Upland (pe20-25)	4,000	2,800	2,000
033QT: Quinlan-----	Shallow Prairie (pe20-25)	1,800	1,350	1,000
Woodward-----	Loamy Upland (pe20-25)	4,000	2,800	2,000
033YE: Yahola-----	Sandy Lowland (pe20-25)	7,000	4,900	3,500
077BP: Woodward-----	Loamy Upland (pe24-32)	4,000	2,800	2,000
Port-----	Loamy Lowland (pe24-32)	8,500	6,100	4,500
077FU: Farnum-----	Loamy Upland (pe24-32)	5,500	4,000	2,500
077GE: Gerlane-----	Subirrigated (pe24-32)	9,000	8,000	7,000
077MC: Minco-----	Loamy Upland (pe24-32)	5,500	3,850	2,750
077MN: Minco-----	Loamy Upland (pe24-32)	5,500	3,850	2,750
077MO: Minco-----	Loamy Upland (pe24-32)	5,500	3,850	2,750
077QU: Quinlan-----	Shallow Prairie (pe24-32)	2,500	1,800	1,300
077SE: Shellabarger-----	Sandy (pe24-32)	4,500	3,200	2,000
077WE: Quinlan-----	Shallow Prairie (pe24-32)	2,500	1,800	1,300
Woodward-----	Loamy Upland (pe24-32)	4,000	2,800	2,000
095AC: Albion-----	Sandy (pe24-32)	4,000	3,000	2,000
095DA: Dillwyn-----	Subirrigated (pe24-32)	9,000	8,000	7,000
Plevna-----	Subirrigated (pe24-32)	9,000	8,000	7,000
095FA: Farnum-----	Sandy (pe24-32)	5,000	3,500	2,500
095PB: Pratt-----	Sands (pe24-32)	4,500	3,500	2,500
095SB: Shellabarger-----	Sandy (pe24-32)	4,500	3,200	2,000
095WA: Waldeck-----	Subirrigated (pe24-32)	9,000	8,000	7,000
097LN: Lincoln-----	Sandy Lowland (pe20-25)	3,000	2,300	1,800
097QW: Quinlan-----	Shallow Prairie (pe20-25)	2,500	1,800	1,300
Woodward-----	Loamy Upland (pe20-25)	4,000	2,800	2,000
097WA: Waldeck-----	Subirrigated (pe20-25)	9,000	8,000	7,000
151AB: Albion-----	Sandy (pe24-32)	4,000	3,000	2,000
151AO: Albion-----	Sandy (pe24-32)	4,000	3,000	2,000
151AS: Albion-----	Sandy (pe24-32)	4,000	3,000	2,000
Shellabarger-----	Sandy (pe24-32)	4,500	3,200	2,000
151BC: Blanket-----	Loamy Upland (pe21-28)	6,500	5,000	3,000
151CA: Carwile-----	Sandy (pe21-28)	5,000	3,800	3,000
151CK: Case-----	Limy Upland (pe24-32)	5,000	4,000	3,000
Clark-----	Limy Upland (pe24-32)	5,000	4,000	3,000
151CM: Clark-----	Limy Upland (pe21-28)	5,000	4,000	3,000
151CN: Clark-----	Limy Upland (pe24-32)	5,000	4,000	3,000
151CO: Clark-----	Limy Upland (pe24-32)	5,000	4,000	3,000
Ost-----	Loamy Upland (pe24-32)	5,500	4,000	2,500
151FA: Farnum-----	Loamy Upland (pe21-28)	5,500	4,000	2,500
151KP: Kanza-----	---	---	---	---
Plevna-----	Subirrigated (pe21-28)	9,000	8,000	7,000
151OS: Ost-----	Loamy Upland (pe21-28)	5,500	4,000	2,500
151PM: Pratt-----	Sands (pe21-28)	4,500	3,500	2,500

RANGELAND PRODUCTIVITY--Continued
Barber County, Kansas

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
151PN:				
Pratt-----	Sands (pe21-28)	4,500	3,500	2,500
151PO:				
Pratt-----	Sands (pe21-28)	4,500	3,500	2,500
Carwile-----	Sandy (pe21-28)	5,000	3,800	3,000
151SA:				
Albion-----	Sandy (pe21-28)	4,000	3,000	2,000
Kaski-----	Loamy Lowland (pe21-28)	7,000	6,000	4,500
151SE:				
Shellabarger-----	Sandy (pe21-28)	4,500	3,200	2,000
1439:				
Crisfield-----	Sandy Terrace (pe24-32)	6,000	5,000	3,500
Ad:				
Albion-----	Sandy (pe20-25)	4,000	3,000	2,000
Shellabarger-----	Sandy (pe20-25)	4,500	3,200	2,000
Ae:				
Albion-----	Sandy (pe20-25)	4,000	3,000	2,000
Shellabarger-----	Sandy (pe20-25)	4,500	3,200	2,000
AED:				
Arents, Earthen Dam-----	---	---	---	---
AN:				
Albion-----	Sandy (pe20-25)	4,000	3,000	2,000
Shellabarger-----	Sandy (pe20-25)	4,500	3,200	2,000
As:				
Clairemont-----	Saline Lowland (pe20-25)	3,500	2,600	1,800
At:				
Attica-----	Sandy (pe24-32)	4,500	3,000	2,000
Ba:				
Blanket-----	Loamy Upland (pe20-25)	6,500	5,000	3,000
Bb:				
Blanket-----	Loamy Upland (pe20-25)	6,500	5,000	3,000
Bc:				
Blanket-----	Loamy Upland (pe20-25)	6,500	5,000	3,000
Bf:				
Quinlan-----	Shallow Prairie (pe20-25)	1,800	1,350	1,000
Clairemont-----	Loamy Lowland (pe20-25)	3,400	2,600	1,800
Ca:				
Canadian-----	Sandy Terrace (pe24-32)	8,500	6,100	4,500
Cc:				
Case-----	Limy Upland (pe24-32)	5,000	4,000	3,000
Clark-----	Limy Upland (pe24-32)	5,000	4,000	3,000
Cd:				
Clairemont-----	Loamy Lowland (pe20-25)	3,400	2,600	1,800
Cf:				
Clairemont-----	Loamy Lowland (pe20-25)	3,400	2,600	1,800
Ck:				
Clark-----	Limy Upland (pe24-32)	5,000	4,000	3,000
Fa:				
Farnum-----	Sandy (pe24-32)	5,000	3,500	2,500
Fm:				
Farnum-----	Loamy Upland (pe24-32)	5,500	4,000	2,500
Fr:				
Farnum-----	Loamy Upland (pe24-32)	5,500	4,000	2,500
Fu:				
Farnum-----	Loamy Upland (pe24-32)	5,500	4,000	2,500
Ga:				
Grant-----	Loamy Upland (pe24-32)	5,500	3,700	2,500
Gb:				
Grant-----	Loamy Upland (pe24-32)	5,500	3,700	2,500
Gc:				
Grant-----	Loamy Upland (pe24-32)	5,500	3,700	2,500
GRP:				
Gravel Pits-----	---	---	---	---
INT:				
Aquolls-----	---	---	---	---
Ka:				
Kanza-----	Subirrigated (pe24-32)	9,000	8,000	7,000
Kf:				
Kingfisher-----	Loamy Upland (pe20-25)	5,000	3,500	2,500
Kv:				
Kingfisher-----	Loamy Upland (pe20-25)	5,000	3,500	2,500
Vernon-----	Red Clay Prairie (pe20-25)	4,000	2,800	2,000
Kz:				
Kingfisher-----	Loamy Upland (pe20-25)	5,000	3,500	2,500
Vernon-----	Red Clay Prairie (pe20-25)	4,000	2,800	2,000
Ln:				
Lincoln-----	Sandy Lowland (pe20-25)	3,000	2,300	1,800
LNN:				
Lincoln-----	Sandy Lowland (pe20-25)	3,000	2,300	1,800
Ma:				
Mangum-----	Clay Lowland (pe20-25)	3,000	2,250	1,500
Mg:				
Mangum-----	Clay Lowland (pe20-25)	3,000	2,250	1,500
Drummond-----	Saline Lowland (pe20-25)	7,000	5,800	5,000
Mn:				

RANGELAND PRODUCTIVITY--Continued
Barber County, Kansas

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
Minco-----	Loamy Upland (pe24-32)	5,500	3,850	2,750
Na:				
Naron-----	Sandy (pe24-32)	4,500	3,000	2,000
Nb:				
Naron-----	Sandy (pe24-32)	4,500	3,000	2,000
Os:				
Ost-----	Loamy Upland (pe24-32)	5,500	4,000	2,500
Ot:				
Ost-----	Loamy Upland (pe24-32)	5,500	4,000	2,500
Pa:				
Pond Creek-----	Loamy Upland (pe24-32)	5,500	3,850	2,750
Pd:				
Pond Creek-----	Loamy Upland (pe24-32)	5,500	3,850	2,750
Ph:				
Dale-----	Loamy Terrace (pe24-32)	8,500	6,100	4,500
Pk:				
Buttermilk-----	Saline Lowland (pe24-32)	4,000	3,000	2,000
Ps:				
Pratt-----	Sands (pe24-32)	4,500	3,500	2,500
Pt:				
Pratt-----	Sands (pe20-25)	4,500	3,500	2,500
Tivoli-----	Sands (pe20-25)	2,000	1,400	1,000
Qn:				
Quinlan-----	Shallow Prairie (pe20-25)	2,500	1,800	1,300
Qw:				
Quinlan-----	Shallow Prairie (pe20-25)	2,500	1,800	1,300
Woodward-----	Loamy Upland (pe20-25)	4,000	2,800	2,000
Rb:				
Knoco-----	Red Shale (pe20-25)	1,400	1,000	600
Rock Outcrop-----	---	---	---	---
Sb:				
Shellabarger-----	Sandy (pe20-25)	4,500	3,200	2,000
SBB:				
Shellabarger-----	Sandy (pe24-32)	4,500	3,200	2,000
Sc:				
Shellabarger-----	Sandy (pe20-25)	4,500	3,200	2,000
Tv:				
Tivoli-----	Choppy Sands (pe20-25)	2,000	1,400	1,000
Vn:				
Vernon-----	Red Clay Prairie (pe20-25)	4,000	2,800	2,000
Vr:				
Vernon-----	Red Clay Prairie (pe20-25)	4,000	2,800	2,000
Vs:				
Knoco-----	Red Shale (pe20-25)	1,400	1,000	600
Vernon-----	Red Clay Prairie (pe20-25)	4,000	2,800	2,000
W:				
Water-----	---	---	---	---
Wa:				
Waldeck-----	Subirrigated (pe24-32)	9,000	8,000	7,000
Wo:				
Woodward-----	Shallow Prairie (pe20-25)	4,000	2,800	2,000
Quinlan-----	Loamy Upland (pe20-25)	2,500	1,800	1,300
Ws:				
Woodward-----	Shallow Prairie (pe20-25)	4,000	2,800	2,000
Quinlan-----	Loamy Upland (pe20-25)	2,500	1,800	1,300
Ya:				
Yahola-----	Subirrigated (pe20-25)	7,000	4,900	3,500
Ze:				
Zenda-----	Subirrigated (pe20-25)	9,000	8,000	7,000

BUILDING SITE DEVELOPMENT
Barber County, Kansas

Building Site Development

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. The following tables show the degree and kind of soil limitations that affect dwellings with and without basements, small commercial buildings, local roads and streets, shallow excavations, and lawns and landscaping.

The ratings in the tables are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Local roads and streets have an all-weather surface and carry automobile and light truck traffic all year. They have a subgrade of cut or fill soil material; a base of gravel, crushed rock, or soil material stabilized by lime or cement; and a surface of flexible material (asphalt), rigid material (concrete), or gravel with a binder. The ratings are based on the soil properties that affect the ease of excavation and grading and the traffic-supporting capacity. The properties that affect the ease of excavation and grading are depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, depth to a water table, ponding, flooding, the amount of large stones, and slope. The properties that affect the traffic-supporting capacity are soil strength (as inferred from the AASHTO group index number), subsidence, linear extensibility (shrink-swell potential), the potential for frost action, depth to a water table, and ponding.

Shallow excavations are trenches or holes dug to a maximum depth of 5 or 6 feet for graves, utility lines, open ditches, or other purposes. The ratings are based on the soil properties that influence the ease of digging and the resistance to sloughing. Depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, the amount of large stones, and dense layers influence the ease of digging, filling, and compacting. Depth to the seasonal high water table, flooding, and ponding may restrict the period when excavations can be made. Slope influences the ease of using machinery. Soil texture, depth to the water table, and linear extensibility (shrink-swell potential) influence the resistance to sloughing.

Lawns and landscaping require soils on which turf and ornamental trees and shrubs can be established and maintained. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer.

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
033CP: Clairemont-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
033OA: Obaro-----	100	Somewhat limited Slope	0.04	Somewhat limited Depth to soft bedrock Slope	0.10 0.04	Very limited Slope	1.00
033OB: Obaro-----	75	Somewhat limited Slope	0.96	Somewhat limited Slope Depth to soft bedrock	0.96 0.10	Very limited Slope	1.00
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
033QR: Quinlan-----	55	Somewhat limited Depth to soft bedrock Slope	1.00 0.37	Very limited Depth to soft bedrock Slope	1.00 0.37	Very limited Depth to soft bedrock Slope	1.00 1.00
Woodward-----	45	Somewhat limited Slope	0.37	Somewhat limited Depth to soft bedrock Slope	0.46 0.37	Very limited Slope	1.00
033QT: Quinlan-----	55	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Slope Depth to soft bedrock	1.00 1.00
Woodward-----	45	Very limited Slope	1.00	Very limited Slope Depth to soft bedrock	1.00 0.46	Very limited Slope	1.00
033YE: Yahola-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
077BP: Woodward-----	65	Somewhat limited Slope	0.37	Somewhat limited Depth to soft bedrock Slope	0.42 0.37	Very limited Slope	1.00
Port-----	35	Very limited Flooding	1.00	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding	1.00
077FU: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
077GE: Gerlane-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.61	Very limited Flooding	1.00
077MC: Minco-----	100	Not limited		Not limited		Not limited	
077MN: Minco-----	100	Not limited		Not limited		Not limited	
077MO: Minco-----	100	Not limited		Not limited		Somewhat limited Slope	0.12
077QU: Quinlan-----	100	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock	1.00	Somewhat limited Depth to soft bedrock Slope	1.00 0.12
077SE: Shellabarger-----	100	Not limited		Not limited		Not limited	
077WE: Quinlan-----	50	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock	1.00	Somewhat limited Depth to soft bedrock	1.00
Woodward-----	50	Not limited		Somewhat limited Depth to soft bedrock	0.90	Not limited	
095AC: Albion-----	100	Not limited		Not limited		Somewhat limited Slope	0.12

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
095DA: Dillwyn-----	60	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
Plevna-----	40	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
095FA: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
095PB: Pratt-----	100	Not limited		Not limited		Somewhat limited Slope	0.48
095SB: Shellabarger-----	100	Not limited		Not limited		Not limited	
095WA: Waldeck-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.95	Very limited Flooding	1.00
097LN: Lincoln-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.03	Very limited Flooding	1.00
097QW: Quinlan-----	55	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00
Woodward-----	45	Somewhat limited Slope	0.84	Somewhat limited Slope Depth to soft bedrock	0.84 0.71	Very limited Slope	1.00
097WA: Waldeck-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.95	Very limited Flooding	1.00
151AB: Albion-----	100	Not limited		Not limited		Not limited	
151AO: Albion-----	100	Not limited		Not limited		Somewhat limited Slope	0.12
151AS: Albion-----	50	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope	1.00
Shellabarger-----	50	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope	1.00
151BC: Blanket-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
151CA: Carwile-----	100	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00
151CK: Case-----	60	Somewhat limited Shrink-swell Slope	0.50 0.37	Somewhat limited Shrink-swell Slope	0.50 0.37	Very limited Slope Shrink-swell	1.00 0.50
Clark-----	40	Somewhat limited Shrink-swell Slope	0.50 0.16	Somewhat limited Shrink-swell Slope	0.50 0.16	Very limited Slope Shrink-swell	1.00 0.50
151CM: Clark-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
151CN: Clark-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
151CO: Clark-----	70	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Ost-----	30	Not limited		Not limited		Not limited	

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
151FA: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
151KP: Kanza-----	50	Very limited Flooding Depth to saturated zone	1.00 0.98	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.98
Plevna-----	50	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
151OS: Ost-----	100	Not limited		Not limited		Not limited	
151PM: Pratt-----	100	Not limited		Not limited		Somewhat limited Slope	0.48
151PN: Pratt-----	100	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
151PO: Pratt-----	60	Not limited		Not limited		Somewhat limited Slope	0.48
Carwile-----	40	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00
151SA: Albion-----	70	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope	1.00
Kaski-----	30	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding	1.00	Very limited Flooding Shrink-swell	1.00 0.50
151SE: Shellabarger-----	100	Not limited		Not limited		Not limited	
1439: Crisfield-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.87	Very limited Flooding	1.00
Ad: Albion-----	60	Not limited		Not limited		Not limited	
Shellabarger-----	40	Not limited		Not limited		Not limited	
Ae: Albion-----	55	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Shellabarger-----	45	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
AN: Albion-----	65	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Shellabarger-----	35	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
As: Clairemont-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
At: Attica-----	100	Not limited		Not limited		Not limited	
Ba: Blanket-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Bb: Blanket-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Bc: Blanket-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Bf: Quinlan-----	60	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00
Clairemont-----	40	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ca: Canadian-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Cc: Case-----	50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.00
Clark-----	50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.00
Cd: Clairemont-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Cf: Clairemont-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Ck: Clark-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Fa: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Fm: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Fr: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Fu: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Ga: Grant-----	100	Not limited		Not limited		Not limited	
Gb: Grant-----	100	Not limited		Not limited		Not limited	
Gc: Grant-----	100	Not limited		Not limited		Somewhat limited Slope	0.12
GRP: Gravel Pits-----	100	Not rated		Not rated		Not rated	
INT: Aguolls-----	100	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
Ka: Kanza-----	100	Very limited Flooding Depth to saturated zone	1.00 0.98	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.98
Kf: Kingfisher-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Shrink-swell	0.64 0.50	Somewhat limited Shrink-swell	0.50
Kv: Kingfisher-----	70	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Shrink-swell	0.64 0.50	Somewhat limited Shrink-swell	0.50
Vernon-----	30	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to soft bedrock	1.00 0.97	Very limited Shrink-swell	1.00
Kz: Kingfisher-----	60	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Shrink-swell	0.64 0.50	Somewhat limited Shrink-swell	0.50
Vernon-----	40	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to soft bedrock	1.00 0.97	Slope Very limited Shrink-swell Slope	0.12 1.00 0.12

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ln: Lincoln-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.03	Very limited Flooding	1.00
LNN: Lincoln-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.03	Very limited Flooding	1.00
Ma: Mangum-----	100	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Shrink-swell	1.00 1.00
Mg: Mangum-----	65	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Shrink-swell	1.00 1.00
Drummond-----	35	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.61	Very limited Flooding Shrink-swell	1.00 1.00
Mn: Minco-----	100	Not limited		Not limited		Not limited	
Na: Naron-----	100	Not limited		Not limited		Not limited	
Nb: Naron-----	100	Not limited		Not limited		Not limited	
Os: Ost-----	100	Not limited		Not limited		Not limited	
Ot: Ost-----	100	Not limited		Not limited		Not limited	
Pa: Pond Creek-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Pd: Pond Creek-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Ph: Dale-----	100	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 0.50
Pk: Buttermilk-----	100	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell Depth to saturated zone	1.00 0.50 0.08	Very limited Flooding Shrink-swell	1.00 0.50
Ps: Pratt-----	100	Somewhat limited Slope	0.00	Somewhat limited Slope	0.00	Very limited Slope	1.00
Pt: Pratt-----	70	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Tivoli-----	30	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Qn: Quinlan-----	100	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock	1.00	Somewhat limited Depth to soft bedrock	1.00
Qw: Quinlan-----	60	Somewhat limited Depth to soft bedrock Slope	1.00 0.16	Very limited Depth to soft bedrock Slope	1.00 0.16	Very limited Depth to soft bedrock Slope	1.00 1.00
Woodward-----	40	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Rb: Knoco-----	70	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 1.00	Very limited Slope Depth to soft bedrock Shrink-swell	1.00 1.00 1.00
Rock Outcrop-----	30	Shrink-swell Not rated	1.00	Not rated	1.00	Not rated	

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Sb: Shellabarger-----	100	Not limited		Not limited		Somewhat limited Slope	0.12
SBB: Shellabarger-----	100	Not limited		Not limited		Not limited	
Sc: Shellabarger-----	100	Not limited		Not limited		Somewhat limited Slope	0.12
Tv: Tivoli-----	100	Somewhat limited Slope	0.84	Somewhat limited Slope	0.84	Very limited Slope	1.00
Vn: Vernon-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to soft bedrock	1.00 0.97	Very limited Shrink-swell Slope	1.00 0.00
Vr: Vernon-----	100	Very limited Shrink-swell Slope	1.00 0.16	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 0.97 0.16	Very limited Shrink-swell Slope	1.00 1.00
Vs: Knoco-----	50	Very limited Depth to soft bedrock Shrink-swell	1.00 1.00	Very limited Shrink-swell Depth to soft bedrock	1.00 1.00	Very limited Depth to soft bedrock Shrink-swell	1.00 1.00
Vernon-----	50	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to soft bedrock	1.00 0.42	Very limited Shrink-swell	1.00
W: Water-----	100	Not rated		Not rated		Not rated	
Wa: Waldeck-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.95	Very limited Flooding	1.00
Wo: Woodward-----	70	Not limited		Not limited		Not limited	
Quinlan-----	30	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock	1.00	Somewhat limited Depth to soft bedrock	1.00
Ws: Woodward-----	65	Not limited		Not limited		Somewhat limited Slope	0.12
Quinlan-----	35	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock	1.00	Somewhat limited Depth to soft bedrock Slope	1.00 0.12
Ya: Yahola-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Ze: Zenda-----	100	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 0.95 0.50	Very limited Flooding Shrink-swell	1.00 0.50

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
033CP: Clairemont-----	100	Very limited Flooding	1.00	Somewhat limited Flooding Cutbanks cave	0.80 0.10	Very limited Flooding	1.00
033OA: Obaro-----	100	Somewhat limited Slope	0.04	Somewhat limited Cutbanks cave Depth to soft bedrock Slope	0.10 0.10 0.04	Somewhat limited Depth to bedrock Slope	0.10 0.04
033OB: Obaro-----	75	Somewhat limited Slope	0.96	Somewhat limited Slope Cutbanks cave Depth to soft bedrock	0.96 0.10 0.10	Somewhat limited Slope Depth to bedrock	0.96 0.10
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
033QR: Quinlan-----	55	Somewhat limited Depth to soft bedrock Slope	1.00 0.37	Very limited Depth to soft bedrock Slope Cutbanks cave	1.00 0.37 0.10	Very limited Depth to bedrock Droughty Slope	1.00 0.74 0.37
Woodward-----	45	Somewhat limited Slope	0.37	Somewhat limited Depth to soft bedrock Slope Cutbanks cave	0.46 0.37 0.10	Somewhat limited Depth to bedrock Slope	0.46 0.37
033QT: Quinlan-----	55	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope Cutbanks cave	1.00 1.00 0.10	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.74
Woodward-----	45	Very limited Slope	1.00	Very limited Slope Depth to soft bedrock Cutbanks cave	1.00 0.46 0.10	Very limited Slope Depth to bedrock	1.00 0.46
033YE: Yahola-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding	1.00 0.60	Somewhat limited Flooding	0.60
077BP: Woodward-----	65	Somewhat limited Slope	0.37	Somewhat limited Depth to soft bedrock Slope Cutbanks cave	0.42 0.37 0.10	Somewhat limited Depth to bedrock Slope	0.42 0.37
Port-----	35	Very limited Flooding	1.00	Somewhat limited Flooding Cutbanks cave	0.80 0.10	Very limited Flooding	1.00
077FU: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
077GE: Gerlane-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 0.61 0.60	Somewhat limited Flooding	0.60
077MC: Minco-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
077MN: Minco-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
077MO: Minco-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
077QU: Quinlan-----	100	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock Cutbanks cave	1.00 0.10	Very limited Depth to bedrock Droughty	1.00 1.00

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
077SE: Shellabarger-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
077WE: Quinlan-----	50	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock Cutbanks cave	1.00 0.10	Very limited Depth to bedrock	1.00
Woodward-----	50	Not limited		Somewhat limited Depth to soft bedrock Cutbanks cave	0.90 0.10	Droughty Somewhat limited Depth to bedrock	1.00 0.90
095AC: Albion-----	100	Not limited		Very limited Cutbanks cave	1.00	Not limited	
095DA: Dillwyn-----	60	Somewhat limited Depth to saturated zone	0.19	Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.22
Plevna-----	40	Very limited Flooding	1.00	Depth to saturated zone Very limited Depth to saturated zone Cutbanks cave	1.00 1.00 1.00	Depth to saturated zone Very limited Flooding	0.19 1.00
		Depth to saturated zone	1.00	Flooding	0.80	Depth to saturated zone	1.00
095FA: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
095PB: Pratt-----	100	Not limited		Very limited Cutbanks cave	1.00	Not limited	
095SB: Shellabarger-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
095WA: Waldeck-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 0.95 0.60	Somewhat limited Flooding	0.60
097LN: Lincoln-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding Depth to saturated zone	1.00 0.60 0.03	Somewhat limited Droughty Flooding	0.80 0.60
097QW: Quinlan-----	55	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope Cutbanks cave	1.00 1.00 0.10	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.62
Woodward-----	45	Somewhat limited Slope	0.84	Somewhat limited Slope Depth to soft bedrock Cutbanks cave	0.84 0.71 0.10	Somewhat limited Slope Depth to bedrock	0.84 0.71
097WA: Waldeck-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 0.95 0.60	Somewhat limited Flooding	0.60
151AB: Albion-----	100	Not limited		Very limited Cutbanks cave	1.00	Not limited	
151AO: Albion-----	100	Not limited		Very limited Cutbanks cave	1.00	Not limited	
151AS: Albion-----	50	Somewhat limited Slope	0.37	Very limited Cutbanks cave Slope	1.00 0.37	Somewhat limited Slope	0.37
Shellabarger-----	50	Somewhat limited Slope	0.37	Somewhat limited Slope Cutbanks cave	0.37 0.10	Somewhat limited Slope	0.37

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
151BC: Blanket-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave Too clayey	0.10 0.03	Not limited	
151CA: Carwile-----	100	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Too clayey Cutbanks cave	1.00 0.28 0.10	Very limited Depth to saturated zone	1.00
151CK: Case-----	60	Somewhat limited Shrink-swell Slope	0.50 0.37	Somewhat limited Slope Cutbanks cave	0.37 0.10	Somewhat limited Slope	0.37
Clark-----	40	Somewhat limited Shrink-swell Slope	0.50 0.16	Somewhat limited Slope Cutbanks cave	0.16 0.10	Somewhat limited Slope	0.16
151CM: Clark-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
151CN: Clark-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
151CO: Clark-----	70	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Ost-----	30	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
151FA: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
151KP: Kanza-----	50	Very limited Flooding Depth to saturated zone	1.00 0.75	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 1.00 0.80	Very limited Flooding Depth to saturated zone Droughty	1.00 0.75 0.02
Plevna-----	50	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Flooding	1.00 1.00 0.80	Very limited Flooding Depth to saturated zone	1.00 1.00
151OS: Ost-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
151PM: Pratt-----	100	Not limited		Very limited Cutbanks cave	1.00	Not limited	
151PN: Pratt-----	100	Somewhat limited Slope	0.16	Very limited Cutbanks cave Slope	1.00 0.16	Somewhat limited Slope	0.16
151PO: Pratt-----	60	Not limited		Very limited Cutbanks cave	1.00	Not limited	
Carwile-----	40	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Too clayey Cutbanks cave	1.00 0.28 0.10	Very limited Depth to saturated zone	1.00
151SA: Albion-----	70	Somewhat limited Slope	0.37	Very limited Cutbanks cave Slope	1.00 0.37	Somewhat limited Slope	0.37
Kaski-----	30	Very limited Flooding Shrink-swell	1.00 0.50	Somewhat limited Flooding Cutbanks cave	0.60 0.10	Somewhat limited Flooding	0.60
151SE: Shellabarger-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
1439: Crisfield-----	100	Somewhat limited Flooding	0.40	Very limited Cutbanks cave Depth to saturated zone	1.00 0.87	Somewhat limited Droughty	0.31

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ad: Albion-----	60	Not limited		Very limited Cutbanks cave	1.00	Not limited	
Shellabarger-----	40	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Ae: Albion-----	55	Somewhat limited Slope	0.16	Very limited Cutbanks cave Slope	1.00 0.16	Somewhat limited Slope	0.16
Shellabarger-----	45	Somewhat limited Slope	0.16	Somewhat limited Slope Cutbanks cave	0.16 0.10	Somewhat limited Slope	0.16
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
AN: Albion-----	65	Somewhat limited Slope	0.16	Very limited Cutbanks cave Slope	1.00 0.16	Somewhat limited Slope	0.16
Shellabarger-----	35	Somewhat limited Slope	0.16	Somewhat limited Slope Cutbanks cave	0.16 0.10	Somewhat limited Slope	0.16
As: Clairemont-----	100	Very limited Flooding	1.00	Somewhat limited Flooding Cutbanks cave	0.80 0.10	Very limited Flooding Salinity	1.00 1.00
At: Attica-----	100	Not limited		Very limited Cutbanks cave	1.00	Not limited	
Ba: Blanket-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave Too clayey	0.10 0.03	Not limited	
Bb: Blanket-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave Too clayey	0.10 0.03	Not limited	
Bc: Blanket-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave Too clayey	0.10 0.03	Not limited	
Bf: Quinlan-----	60	Very limited Depth to soft bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope Cutbanks cave	1.00 1.00 0.10	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.74
Clairemont-----	40	Very limited Flooding	1.00	Somewhat limited Flooding Cutbanks cave	0.80 0.10	Very limited Flooding	1.00
Ca: Canadian-----	100	Somewhat limited Flooding	0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
Cc: Case-----	50	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Clark-----	50	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Cd: Clairemont-----	100	Very limited Flooding	1.00	Somewhat limited Flooding Cutbanks cave	0.60 0.10	Somewhat limited Flooding	0.60
Cf: Clairemont-----	100	Very limited Flooding	1.00	Somewhat limited Flooding Cutbanks cave	0.80 0.10	Very limited Flooding	1.00
Ck: Clark-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Fa: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Fm: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Fr: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Fu: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Ga: Grant-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Gb: Grant-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Gc: Grant-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
GRP: Gravel Pits-----	100	Not rated		Not rated		Not rated	
INT: Aquolls-----	100	Very limited Depth to saturated zone Ponding Frost action	1.00 1.00 0.50	Very limited Depth to saturated zone Ponding Cutbanks cave	1.00 1.00 0.10	Very limited Depth to saturated zone Ponding	1.00 1.00
Ka: Kanza-----	100	Very limited Flooding Depth to saturated zone	1.00 0.75	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 1.00 0.80	Very limited Flooding Depth to saturated zone Droughty	1.00 0.75 0.04
KF: Kingfisher-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Cutbanks cave	0.64 0.10	Somewhat limited Depth to bedrock	0.65
Kv: Kingfisher-----	70	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Cutbanks cave	0.64 0.10	Somewhat limited Depth to bedrock	0.65
Vernon-----	30	Very limited Shrink-swell	1.00	Somewhat limited Depth to soft bedrock Too clayey Cutbanks cave	0.97 0.28 0.10	Somewhat limited Depth to bedrock Droughty Content of large stones	0.97 0.11 0.03
Kz: Kingfisher-----	60	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Cutbanks cave	0.64 0.10	Somewhat limited Depth to bedrock	0.65
Vernon-----	40	Very limited Shrink-swell	1.00	Somewhat limited Depth to soft bedrock Too clayey Cutbanks cave	0.97 0.28 0.10	Somewhat limited Depth to bedrock Droughty Content of large stones	0.97 0.11 0.03
Ln: Lincoln-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding Depth to saturated zone	1.00 0.80 0.03	Very limited Flooding Droughty	1.00 0.89
LNN: Lincoln-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding Depth to saturated zone	1.00 0.60 0.03	Somewhat limited Droughty Flooding	0.92 0.60
Ma: Mangum-----	100	Very limited Flooding Shrink-swell	1.00 1.00	Somewhat limited Flooding Too clayey Cutbanks cave	0.60 0.50 0.10	Very limited Too clayey Flooding	1.00 0.60

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Mg: Mangum-----	65	Very limited Shrink-swell Flooding	1.00 0.40	Somewhat limited Too clayey Cutbanks cave	0.50 0.10	Very limited Too clayey	1.00
Drummond-----	35	Very limited Shrink-swell Flooding	1.00 0.40	Somewhat limited Depth to saturated zone Too clayey Cutbanks cave	0.61 0.28 0.10	Not limited	
Mn: Minco-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Na: Naron-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Nb: Naron-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Os: Ost-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Ot: Ost-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Pa: Pond Creek-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Pd: Pond Creek-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Ph: Dale-----	100	Somewhat limited Shrink-swell Flooding	0.50 0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
Pk: Buttermilk-----	100	Somewhat limited Shrink-swell Flooding	0.50 0.40	Somewhat limited Cutbanks cave Depth to saturated zone	0.10 0.08	Somewhat limited Salinity	0.13
Ps: Pratt-----	100	Somewhat limited Slope	0.00	Very limited Cutbanks cave Slope	1.00 0.00	Somewhat limited Slope	0.00
Pt: Pratt-----	70	Somewhat limited Slope	0.16	Very limited Cutbanks cave Slope	1.00 0.16	Somewhat limited Slope	0.16
Tivoli-----	30	Somewhat limited Slope	0.16	Very limited Cutbanks cave Slope	1.00 0.16	Somewhat limited Droughty Slope	0.98 0.16
Qn: Quinlan-----	100	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock Cutbanks cave	1.00 0.10	Very limited Depth to bedrock Droughty	1.00 0.74
Qw: Quinlan-----	60	Somewhat limited Depth to soft bedrock Slope	1.00 0.16	Very limited Depth to soft bedrock Slope Cutbanks cave	1.00 0.16 0.10	Very limited Depth to bedrock Droughty Slope	1.00 0.74 0.16
Woodward-----	40	Somewhat limited Slope	0.16	Somewhat limited Slope Cutbanks cave	0.16 0.10	Somewhat limited Slope	0.16
Rb: Knoco-----	70	Very limited Depth to soft bedrock Slope Shrink-swell	1.00 1.00 1.00	Very limited Depth to soft bedrock Slope Depth to dense layer Too clayey Cutbanks cave	1.00 1.00 0.50 0.28 0.10	Very limited Depth to bedrock Droughty Slope Too clayey Sodium content	1.00 1.00 1.00 1.00 1.00
Rock Outcrop-----	30	Not rated		Not rated		Not rated	

BUILDING SITE DEVELOPMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Sb: Shellabarger-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
SBB: Shellabarger-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Sc: Shellabarger-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Tv: Tivoli-----	100	Somewhat limited Slope	0.84	Very limited Cutbanks cave Slope	1.00 0.84	Very limited Droughty Slope	1.00 0.84
Vn: Vernon-----	100	Very limited Shrink-swell	1.00	Somewhat limited Depth to soft bedrock Too clayey Cutbanks cave	0.97 0.28 0.10	Somewhat limited Depth to bedrock Droughty Content of large stones	0.97 0.11 0.03
Vr: Vernon-----	100	Very limited Shrink-swell Slope	1.00 0.16	Somewhat limited Depth to soft bedrock Too clayey Slope Cutbanks cave	0.97 0.28 0.16 0.10	Somewhat limited Depth to bedrock Slope Droughty Content of large stones	0.97 0.16 0.11 0.03
Vs: Knoco-----	50	Very limited Depth to soft bedrock Shrink-swell	1.00 1.00	Very limited Depth to soft bedrock Depth to dense layer Too clayey Cutbanks cave	1.00 0.50 0.28 0.10	Very limited Depth to bedrock Droughty Too clayey Sodium content	1.00 1.00 1.00
Vernon-----	50	Very limited Shrink-swell	1.00	Somewhat limited Depth to soft bedrock Too clayey Cutbanks cave	0.42 0.28 0.10	Somewhat limited Depth to bedrock Droughty Content of large stones	0.42 0.11 0.03
W: Water-----	100	Not rated		Not rated		Not rated	
Wa: Waldeck-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 0.95 0.60	Somewhat limited Flooding	0.60
Wo: Woodward-----	70	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Quinlan-----	30	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock Cutbanks cave	1.00 0.10	Very limited Depth to bedrock Droughty	1.00 0.74
Ws: Woodward-----	65	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
Quinlan-----	35	Somewhat limited Depth to soft bedrock	1.00	Very limited Depth to soft bedrock Cutbanks cave	1.00 0.10	Very limited Depth to bedrock Droughty	1.00 0.74
Ya: Yahola-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding	1.00 0.60	Somewhat limited Flooding	0.60
Ze: Zenda-----	100	Very limited Flooding Shrink-swell	1.00 0.50	Somewhat limited Depth to saturated zone Flooding Cutbanks cave	0.95 0.60 0.10	Somewhat limited Flooding	0.60

CONSTRUCTION MATERIALS
Barber County, Kansas

Construction Materials

The following tables give information about the soils as potential sources of gravel, sand, topsoil, reclamation material, and roadfill. Normal compaction, minor processing, and other standard construction practices are assumed.

The soils are rated good, fair, or poor as potential sources of topsoil, reclamation material, and roadfill. The features that limit the soils as sources of these materials are specified in the tables. The numerical ratings given after the specified features indicate the degree to which the features limit the soils as sources of topsoil, reclamation material, or roadfill. The lower the number, the greater the limitation.

The soils are rated as a probable or improbable source of sand and gravel. A rating of probable means that the source material is likely to be in or below the soil. The numerical ratings in these columns indicate the degree of probability. The number 0.00 indicates that the soil is an improbable source. A number between 0.00 and 1.00 indicates the degree to which the soil is a probable source of sand or gravel.

Sand and gravel are natural aggregates suitable for commercial use with a minimum of processing. They are used in many kinds of construction. Specifications for each use vary widely. In these tables, only the probability of finding material in suitable quantity is evaluated. The suitability of the material for specific purposes is not evaluated, nor are factors that affect excavation of the material. The properties used to evaluate the soil as a source of sand or gravel are gradation of grain sizes (as indicated by the Unified classification of the soil), the thickness of suitable material, and the content of rock fragments. If the lowest layer of the soil contains sand or gravel, the soil is rated as a probable source regardless of thickness. The assumption is that the sand or gravel layer below the depth of observation exceeds the minimum thickness.

Topsoil is used to cover an area so that vegetation can be established and maintained. The upper 40 inches of a soil is evaluated for use as topsoil. Also evaluated is the reclamation potential of the borrow area. The ratings are based on the soil properties that affect plant growth; the ease of excavating, loading, and spreading the material; and reclamation of the borrow area. Toxic substances, soil reaction, and the properties that are inferred from soil texture, such as available water capacity and fertility, affect plant growth. The ease of excavating, loading, and spreading is affected by rock fragments, slope, depth to a water table, soil texture, and thickness of suitable material. Reclamation of the borrow area is affected by slope, depth to a water table, rock fragments, depth to bedrock or a cemented pan, and toxic material.

The surface layer of most soils is generally preferred for topsoil because of its organic matter content. Organic matter greatly increases the absorption and retention of moisture and nutrients for plant growth.

Reclamation material is used in areas that have been drastically disturbed by surface mining or similar activities. When these areas are reclaimed, layers of soil material or unconsolidated geological material, or both, are replaced in a vertical sequence. The reconstructed soil favors plant growth. The ratings in the table do not apply to quarries and other mined areas that require an offsite source of reconstruction material. The ratings are based on the soil properties that affect erosion and stability of the surface and the productive potential of the reconstructed soil. These properties include the content of sodium, salts, and calcium carbonate; reaction; available water capacity; erodibility; texture; content of rock fragments; and content of organic matter and other features that affect fertility.

Roadfill is soil material that is excavated in one place and used in road embankments in another place. In this table, the soils are rated as a source of roadfill for low embankments, generally less than 6 feet high and less exacting in design than higher embankments.

The ratings are for the whole soil, from the surface to a depth of about 5 feet. It is assumed that soil layers will be mixed when the soil material is excavated and spread.

The ratings are based on the amount of suitable material and on soil properties that affect the ease of excavation and the performance of the material after it is in place. The thickness of the suitable material is a major consideration. The ease of excavation is affected by large stones, depth to a water table, and slope. How well the soil performs in place after it has been compacted and drained is determined by its strength (as inferred from the AASHTO classification of the soil) and linear extensibility (shrink-swell potential).

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
033CP: Clairemont-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
033OA: Obaro-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
033OB: Obaro-----	75	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rock Outcrop-----	25	Not rated		Not rated	
033QR: Quinlan-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Woodward-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
033QT: Quinlan-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Woodward-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
033YE: Yahola-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.07
077BP: Woodward-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Port-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
077FU: Farnum-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
077GE: Gerlane-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.09
077MC: Minco-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
077MN: Minco-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
077MO: Minco-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
077QU: Quinlan-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
077SE: Shellabarger-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.09
077WE: Quinlan-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Woodward-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
095AC: Albion-----	100	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.29 0.91
095DA: Dillwyn-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.18 0.18
Plevna-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
095FA: Farnum-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
095PB: Pratt-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.44
095SB: Shellabarger-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.09
095WA: Waldeck-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
097LN: Lincoln-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.22
097QW: Quinlan-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Woodward-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
097WA: Waldeck-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
151AB: Albion-----	100	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.91
151AO: Albion-----	100	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.91

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
151AS: Albion-----	50	Poor Thickest layer Bottom layer	0.00 0.00	Fair Bottom layer Thickest layer	0.01 0.09
Shellabarger-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.09
151BC: Blanket-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
151CA: Carwile-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
151CK: Case-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Clark-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
151CM: Clark-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
151CN: Clark-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.08
151CO: Clark-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ost-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
151FA: Farnum-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.06
151KP: Kanza-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.57 0.98
Plevna-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
151OS: Ost-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
151PM: Pratt-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.57
151PN: Pratt-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.57

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
151PO: Pratt-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.57
Carwile-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.02
151SA: Albion-----	70	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.91
Kaski-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.09
151SE: Shellabarger-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.09
1439: Crisfield-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.10 0.78
Ad: Albion-----	60	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.29 0.91
Shellabarger-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.09
Ae: Albion-----	55	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.29 0.91
Shellabarger-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.09
AED: Arents, Earthen Dam-	100	Not rated		Not rated	
AN: Albion-----	65	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.29 0.91
Shellabarger-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.09
As: Clairemont-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
At: Attica-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.72
Ba: Blanket-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Bb: Blanket-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Bc: Blanket-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Bf: Quinlan-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Clairemont-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ca: Canadian-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.07 0.09
Cc: Case-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Clark-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Cd: Clairemont-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Cf: Clairemont-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ck: Clark-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Fa: Farnum-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.07
Fm: Farnum-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Fr: Farnum-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Fu: Farnum-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ga: Grant-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Gb: Grant-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Gc: Grant-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
GRP: Gravel Pits-----	100	Not rated		Not rated	
INT: Aguolls-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ka: Kanza-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.57 0.65
Kf: Kingfisher-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Kv: Kingfisher-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Vernon-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Kz: Kingfisher-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Vernon-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ln: Lincoln-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.22
LNN: Lincoln-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.56 0.89
Ma: Mangum-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Mg: Mangum-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Drummond-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Mn: Minco-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Na: Naron-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.08
Nb: Naron-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.08

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Os: Ost-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ot: Ost-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Pa: Pond Creek-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Pd: Pond Creek-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ph: Dale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Pk: Buttermilk-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ps: Pratt-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.57
Pt: Pratt-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.57
Tivoli-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.57 0.99
Qn: Quinlan-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Qw: Quinlan-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Woodward-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rb: Knoco-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rock Outcrop-----	30	Not rated		Not rated	
Sb: Shellabarger-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.09
SBB: Shellabarger-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.09

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Sc: Shellabarger-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.09
Tv: Tivoli-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.99 0.99
Vn: Vernon-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Vr: Vernon-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Vs: Knoco-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Vernon-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
W: Water-----	100	Not rated		Not rated	
Wa: Waldeck-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.10
Wo: Woodward-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Quinlan-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ws: Woodward-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Quinlan-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ya: Yahola-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.07
Ze: Zenda-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
033CP: Clairemont-----	100	Poor Low content of organic matter Water erosion	0.00 0.90	Good		Good	
033OA: Obaro-----	100	Fair Low content of organic matter Water erosion Depth to bedrock Too clayey	0.50 0.90 0.90 0.98	Poor Depth to bedrock	0.00	Fair Too Clayey Depth to bedrock Slope	0.64 0.90 0.96
033OB: Obaro-----	75	Fair Low content of organic matter Water erosion Depth to bedrock Too clayey	0.50 0.90 0.90 0.98	Poor Depth to bedrock	0.00	Fair Slope Too Clayey Depth to bedrock	0.04 0.64 0.90
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
033QR: Quinlan-----	55	Poor Depth to bedrock Droughty Low content of organic matter No water erosion limitation	0.00 0.00 0.50 0.99	Poor Depth to bedrock	0.00	Poor Depth to bedrock Slope	0.00 0.63
Woodward-----	45	Fair Depth to bedrock Droughty No water erosion limitation	0.54 0.85 0.99	Poor Depth to bedrock	0.00	Fair Depth to bedrock Slope	0.54 0.63
033QT: Quinlan-----	55	Poor Depth to bedrock Droughty Low content of organic matter No water erosion limitation	0.00 0.00 0.50 0.99	Poor Depth to bedrock Slope	0.00 0.08	Poor Depth to bedrock Slope	0.00 0.00
Woodward-----	45	Fair Depth to bedrock Droughty No water erosion limitation	0.54 0.85 0.99	Poor Depth to bedrock Slope	0.00 0.82	Poor Slope Depth to bedrock	0.00 0.54
033YE: Yahola-----	100	Poor Low content of organic matter	0.00	Good		Good	
077BP: Woodward-----	65	Fair Droughty Depth to bedrock No water erosion limitation	0.29 0.58 0.99	Poor Depth to bedrock	0.00	Fair Depth to bedrock Slope	0.58 0.63
Port-----	35	Poor Low content of organic matter No water erosion limitation	0.00 0.99	Fair Shrink-swell	0.99	Good	
077FU: Farnum-----	100	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.99	Good	

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
077GE: Gerlane-----	100	Poor Low content of organic matter	0.00	Good		Good	
077MC: Minco-----	100	Fair No water erosion limitation	0.99	Good		Good	
077MN: Minco-----	100	Fair No water erosion limitation	0.99	Good		Good	
077MO: Minco-----	100	Fair No water erosion limitation	0.99	Good		Good	
077QU: Quinlan-----	100	Poor Droughty Depth to bedrock Low content of organic matter No water erosion limitation	0.00 0.00 0.50 0.99	Poor Depth to bedrock	0.00	Poor Depth to bedrock	0.00
077SE: Shellabarger-----	100	Poor Low content of organic matter Too acid	0.00 0.84	Good		Good	
077WE: Quinlan-----	50	Poor Droughty Depth to bedrock Low content of organic matter No water erosion limitation	0.00 0.00 0.50 0.99	Poor Depth to bedrock	0.00	Poor Depth to bedrock	0.00
Woodward-----	50	Fair Depth to bedrock Droughty No water erosion limitation	0.10 0.29 0.99	Poor Depth to bedrock	0.00	Fair Depth to bedrock	0.10
095AC: Albion-----	100	Poor Too sandy Low content of organic matter Too acid	0.00 0.00 0.95	Good		Poor Too sandy Rock fragments Hard to reclaim	0.00 0.00 0.68
095DA: Dillwyn-----	60	Poor Wind erosion Low content of organic matter Too sandy Droughty	0.00 0.00 0.36 0.79	Fair Depth to saturated zone	0.53	Fair Too sandy Depth to saturated zone	0.36 0.53
Plevna-----	40	Poor Low content of organic matter	0.00	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone	0.00
095FA: Farnum-----	100	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.98	Good	

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
095PB: Pratt-----	100	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.00 0.00	Good		Poor Too sandy	0.00
095SB: Shellabarger-----	100	Poor Low content of organic matter Too acid	0.00 0.84	Good		Good	
095WA: Waldeck-----	100	Poor Low content of organic matter	0.00	Good		Good	
097LN: Lincoln-----	100	Poor Low content of organic matter Droughty Too sandy	0.00 0.10 0.22	Good		Fair Too sandy	0.22
097QW: Quinlan-----	55	Poor Depth to bedrock Droughty Low content of organic matter No water erosion limitation	0.00 0.00 0.50 0.99	Poor Depth to bedrock Slope	0.00 0.98	Poor Depth to bedrock Slope	0.00 0.00
Woodward-----	45	Fair Depth to bedrock Droughty No water erosion limitation	0.29 0.62 0.99	Poor Depth to bedrock	0.00	Fair Slope Depth to bedrock	0.16 0.29
097WA: Waldeck-----	100	Poor Low content of organic matter	0.00	Good		Good	
151AB: Albion-----	100	Poor Low content of organic matter Too acid	0.00 0.95	Good		Fair Hard to reclaim Rock fragments	0.68 0.72
151AO: Albion-----	100	Poor Low content of organic matter Too acid	0.00 0.95	Good		Fair Hard to reclaim Rock fragments	0.68 0.72
151AS: Albion-----	50	Poor Low content of organic matter Too acid	0.00 0.95	Good		Fair Slope Hard to reclaim Rock fragments	0.63 0.68 0.72
Shellabarger-----	50	Poor Low content of organic matter Too acid	0.00 0.84	Good		Fair Slope	0.63
151BC: Blanket-----	100	Poor Low content of organic matter Too clayey Water erosion	0.00 0.00 0.90	Fair Shrink-swell	0.87	Poor Too Clayey	0.00

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
151CA: Carwile-----	100	Poor Low content of organic matter Too clayey Too acid No water erosion limitation	0.00 0.00 0.97 0.99	Poor Depth to saturated zone Shrink-swell	0.00 0.28	Poor Depth to saturated zone Too Clayey	0.00 0.00
151CK: Case-----	60	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.87	Fair Slope	0.63
Clark-----	40	Poor Low content of organic matter Carbonate content	0.00 0.68	Fair Shrink-swell	0.87	Fair Carbonate content Slope	0.68 0.84
151CM: Clark-----	100	Poor Low content of organic matter Carbonate content	0.00 0.68	Fair Shrink-swell	0.87	Fair Carbonate content	0.68
151CN: Clark-----	100	Poor Low content of organic matter Carbonate content	0.00 0.68	Fair Shrink-swell	0.87	Fair Carbonate content	0.68
151CO: Clark-----	70	Poor Low content of organic matter Carbonate content	0.00 0.68	Fair Shrink-swell	0.87	Fair Carbonate content	0.68
Ost-----	30	Poor Low content of organic matter Carbonate content	0.00 0.68	Good		Fair Carbonate content	0.68
151FA: Farnum-----	100	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.99	Good	
151KP: Kanza-----	50	Poor Wind erosion Low content of organic matter Too sandy Droughty Too acid	0.00 0.00 0.00 0.18 0.95	Fair Depth to saturated zone	0.14	Poor Too sandy Depth to saturated zone	0.00 0.14
Plevna-----	50	Poor Low content of organic matter	0.00	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone	0.00
151OS: Ost-----	100	Poor Low content of organic matter Carbonate content	0.00 0.68	Good		Fair Carbonate content	0.68
151PM: Pratt-----	100	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.00 0.00	Good		Poor Too sandy	0.00
151PN: Pratt-----	100	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.00 0.00	Good		Poor Too sandy Slope	0.00 0.84

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
151PO: Pratt-----	60	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.00 0.00	Good		Poor Too sandy	0.00
Carwile-----	40	Poor Low content of organic matter Too clayey Too acid No water erosion limitation	0.00 0.00 0.97 0.99	Poor Depth to saturated zone Shrink-swell	0.00 0.28	Poor Depth to saturated zone Too Clayey	0.00 0.00
151SA: Albion-----	70	Poor Low content of organic matter Too acid	0.00 0.95	Good		Fair Slope Hard to reclaim Rock fragments	0.63 0.68 0.72
Kaski-----	30	Good		Fair Shrink-swell	0.99	Good	
151SE: Shellabarger-----	100	Poor Low content of organic matter Too acid	0.00 0.84	Good		Good	
1439: Crisfield-----	100	Poor Too sandy Low content of organic matter Too acid Droughty	0.00 0.01 0.46 0.89	Good		Poor Too sandy	0.00
Ad: Albion-----	60	Poor Too sandy Low content of organic matter Too acid	0.00 0.00 0.95	Good		Poor Too sandy Rock fragments Hard to reclaim	0.00 0.00 0.68
Shellabarger-----	40	Poor Low content of organic matter Too acid	0.00 0.84	Good		Good	
Ae: Albion-----	55	Poor Too sandy Low content of organic matter Too acid	0.00 0.00 0.95	Good		Poor Too sandy Rock fragments Hard to reclaim Slope	0.00 0.00 0.68 0.84
Shellabarger-----	45	Poor Low content of organic matter Too acid	0.00 0.84	Good		Fair Slope	0.84
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
AN: Albion-----	65	Poor Low content of organic matter Too sandy Too acid	0.00 0.10 0.95	Good		Fair Too sandy Hard to reclaim Rock fragments Slope	0.10 0.68 0.72 0.84
Shellabarger-----	35	Poor Low content of organic matter Too acid	0.00 0.84	Good		Fair Slope	0.84

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
As: Clairemont-----	100	Poor Low content of organic matter Salinity Water erosion	0.00 0.88 0.90	Good		Poor Salinity	0.00
At: Attica-----	100	Poor Wind erosion Low content of organic matter Too acid	0.00 0.00 0.95	Good		Good	
Ba: Blanket-----	100	Poor Low content of organic matter Too clayey Water erosion	0.00 0.00 0.90	Fair Shrink-swell	0.87	Poor Too Clayey	0.00
Bb: Blanket-----	100	Poor Low content of organic matter Too clayey Water erosion	0.00 0.00 0.90	Fair Shrink-swell	0.87	Poor Too Clayey	0.00
Bc: Blanket-----	100	Poor Low content of organic matter Too clayey Water erosion	0.00 0.00 0.90	Fair Shrink-swell	0.87	Poor Too Clayey	0.00
Bf: Quinlan-----	60	Poor Depth to bedrock Droughty Low content of organic matter No water erosion limitation	0.00 0.00 0.50 0.99	Poor Depth to bedrock Slope	0.00 0.00	Poor Depth to bedrock Slope	0.00 0.00
Clairemont-----	40	Poor Low content of organic matter Water erosion	0.00 0.90	Good		Good	
Ca: Canadian-----	100	Poor Low content of organic matter	0.00	Good		Good	
Cc: Case-----	50	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.87	Good	
Clark-----	50	Poor Low content of organic matter Carbonate content	0.00 0.68	Fair Shrink-swell	0.87	Fair Carbonate content	0.68
Cd: Clairemont-----	100	Poor Low content of organic matter Water erosion	0.00 0.90	Good		Good	
Cf: Clairemont-----	100	Poor Low content of organic matter Water erosion	0.00 0.90	Good		Good	

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ck: Clark-----	100	Poor Low content of organic matter Carbonate content	0.00 0.68	Fair Shrink-swell	0.87	Fair Carbonate content	0.68
Fa: Farnum-----	100	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.87	Good	
Fm: Farnum-----	100	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.87	Good	
Fr: Farnum-----	100	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.87	Good	
Fu: Farnum-----	100	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.87	Good	
Ga: Grant-----	100	Poor Low content of organic matter No water erosion limitation	0.00 0.99	Good		Good	
Gb: Grant-----	100	Poor Low content of organic matter No water erosion limitation	0.00 0.99	Good		Good	
Gc: Grant-----	100	Poor Low content of organic matter No water erosion limitation	0.00 0.99	Good		Good	
GRP: Gravel Pits-----	100	Not rated		Not rated		Not rated	
INT: Aquolls-----	100	Poor Low content of organic matter	0.00	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone	0.00
Ka: Kanza-----	100	Poor Wind erosion Low content of organic matter Too sandy Droughty Too acid	0.00 0.00 0.00 0.52 0.95	Fair Depth to saturated zone	0.14	Poor Too sandy Depth to saturated zone	0.00 0.14
KF: Kingfisher-----	100	Poor Low content of organic matter Depth to bedrock Droughty Too clayey No water erosion limitation	0.00 0.35 0.96 0.98 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.87	Fair Depth to bedrock Too Clayey	0.35 0.49

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Kv: Kingfisher-----	70	Poor Low content of organic matter Depth to bedrock Droughty Too clayey No water erosion limitation	0.00 0.35 0.96 0.98 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.87	Fair Depth to bedrock Too Clayey	0.35 0.49
Vernon-----	30	Poor Low content of organic matter Too clayey Depth to bedrock Droughty No water erosion limitation	0.00 0.00 0.03 0.07 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Depth to bedrock Rock fragments	0.00 0.03 0.88
Kz: Kingfisher-----	60	Poor Low content of organic matter Depth to bedrock Droughty Too clayey No water erosion limitation	0.00 0.35 0.96 0.98 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.87	Fair Depth to bedrock Too Clayey	0.35 0.49
Vernon-----	40	Poor Low content of organic matter Too clayey Depth to bedrock Droughty No water erosion limitation	0.00 0.00 0.03 0.07 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Depth to bedrock Rock fragments	0.00 0.03 0.88
Ln: Lincoln-----	100	Poor Low content of organic matter Droughty Too sandy	0.00 0.06 0.22	Good		Fair Too sandy	0.22
LNN: Lincoln-----	100	Poor Wind erosion Low content of organic matter Too sandy Droughty	0.00 0.00 0.00 0.04	Good		Poor Too sandy	0.00
Ma: Mangum-----	100	Poor Too clayey Low content of organic matter	0.00 0.00	Fair Shrink-swell	0.23	Poor Too Clayey	0.00
Mg: Mangum-----	65	Poor Too clayey Low content of organic matter	0.00 0.00	Fair Shrink-swell	0.23	Poor Too Clayey	0.00
Drummond-----	35	Poor Low content of organic matter Too clayey Droughty Water erosion	0.00 0.00 0.36 0.37	Fair		Poor Too Clayey Salinity	0.00 0.88
Mn: Minco-----	100	Poor Low content of organic matter No water erosion limitation	0.00 0.99	Good		Good	

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Na: Naron-----	100	Poor Low content of organic matter	0.00	Good		Good	
Nb: Naron-----	100	Poor Low content of organic matter	0.00	Good		Good	
Os: Ost-----	100	Poor Low content of organic matter Carbonate content	0.00 0.68	Good		Fair Carbonate content	0.68
Ot: Ost-----	100	Poor Low content of organic matter Carbonate content	0.00 0.68	Good		Fair Carbonate content	0.68
Pa: Pond Creek-----	100	Poor Low content of organic matter Too acid No water erosion limitation	0.00 0.97 0.99	Fair Shrink-swell	0.89	Good	
Pd: Pond Creek-----	100	Poor Low content of organic matter Too acid No water erosion limitation	0.00 0.97 0.99	Fair Shrink-swell	0.89	Good	
Ph: Dale-----	100	Poor Low content of organic matter No water erosion limitation	0.00 0.99	Fair Shrink-swell	0.96	Good	
Pk: Buttermilk-----	100	Poor Low content of organic matter Salinity No water erosion limitation	0.00 0.88 0.99	Fair Shrink-swell	0.96	Fair Salinity	0.88
Ps: Pratt-----	100	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.00 0.00	Good		Poor Too sandy	0.00
Pt: Pratt-----	70	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.00 0.00	Good		Poor Too sandy Slope	0.00 0.84
Tivoli-----	30	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.00 0.01	Good		Poor Too sandy Slope	0.00 0.84

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Qn: Quinlan-----	100	Poor Depth to bedrock Droughty Low content of organic matter No water erosion limitation	0.00 0.00 0.50 0.99	Poor Depth to bedrock	0.00	Poor Depth to bedrock	0.00
Qw: Quinlan-----	60	Poor Depth to bedrock Droughty Low content of organic matter No water erosion limitation	0.00 0.00 0.50 0.99	Poor Depth to bedrock	0.00	Poor Depth to bedrock Slope	0.00 0.84
Woodward-----	40	Fair No water erosion limitation	0.99	Poor Depth to bedrock	0.00	Fair Slope	0.84
Rb: Knoco-----	70	Poor Too clayey Droughty Depth to bedrock Sodium content Carbonate content No water erosion limitation	0.00 0.00 0.00 0.00 0.99 0.99	Poor Depth to bedrock Slope Shrink-swell	0.00 0.00 0.12	Poor Too Clayey Hard to reclaim Depth to bedrock Slope	0.00 0.00 0.00 0.00
Rock Outcrop-----	30	Not rated		Not rated		Not rated	
Sb: Shellabarger-----	100	Poor Low content of organic matter Too acid	0.00 0.84	Good		Good	
SBB: Shellabarger-----	100	Poor Low content of organic matter Too acid	0.00 0.84	Good		Good	
Sc: Shellabarger-----	100	Poor Low content of organic matter Too acid	0.00 0.84	Good		Good	
Tv: Tivoli-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.00 0.00	Good		Poor Too sandy Slope	0.00 0.16
Vn: Vernon-----	100	Poor Low content of organic matter Too clayey Depth to bedrock Droughty No water erosion limitation	0.00 0.00 0.03 0.07 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Depth to bedrock Rock fragments	0.00 0.03 0.88
Vr: Vernon-----	100	Poor Low content of organic matter Too clayey Depth to bedrock Droughty No water erosion limitation	0.00 0.00 0.03 0.07 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Depth to bedrock Slope Rock fragments	0.00 0.03 0.84 0.88

CONSTRUCTION MATERIALS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Vs: Knoco-----	50	Poor Too clayey Droughty Depth to bedrock Sodium content Carbonate content No water erosion limitation	0.00 0.00 0.00 0.00 0.00 0.99 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Hard to reclaim Depth to bedrock	0.00 0.00 0.00
Vernon-----	50	Poor Low content of organic matter Too clayey Droughty Depth to bedrock No water erosion limitation	0.00 0.00 0.07 0.58 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.88	Poor Too Clayey Depth to bedrock Rock fragments	0.00 0.58 0.88
W: Water-----	100	Not rated		Not rated		Not rated	
Wa: Waldeck-----	100	Poor Low content of organic matter	0.00	Good		Good	
Wo: Woodward-----	70	Fair No water erosion limitation	0.99	Poor Depth to bedrock	0.00	Good	
Quinlan-----	30	Poor Depth to bedrock Droughty Low content of organic matter No water erosion limitation	0.00 0.00 0.50 0.99	Poor Depth to bedrock	0.00	Poor Depth to bedrock	0.00
Ws: Woodward-----	65	Fair No water erosion limitation	0.99	Poor Depth to bedrock	0.00	Good	
Quinlan-----	35	Poor Depth to bedrock Droughty Low content of organic matter No water erosion limitation	0.00 0.00 0.50 0.99	Poor Depth to bedrock	0.00	Poor Depth to bedrock	0.00
Ya: Yahola-----	100	Poor Low content of organic matter	0.00	Good		Good	
Ze: Zenda-----	100	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.87	Good	

RECREATIONAL INTERPRETATIONS
Barber County, Kansas

Recreation

The soils of the survey area are rated in the following tables according to limitations that affect their suitability for recreation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the recreational uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The ratings in the tables are based on restrictive soil features, such as wetness, slope, and texture of the surface layer. Susceptibility to flooding is considered. Not considered in the ratings, but important in evaluating a site, are the location and accessibility of the area, the size and shape of the area and its scenic quality, vegetation, access to water, potential water impoundment sites, and access to public sewer lines. The capacity of the soil to absorb septic tank effluent and the ability of the soil to support vegetation also are important. Soils that are subject to flooding are limited for recreational uses by the duration and intensity of flooding and the season when flooding occurs. In planning recreational facilities, onsite assessment of the height, duration, intensity, and frequency of flooding is essential.

The information in this table can be supplemented by other information in this survey, for example, interpretations for building site development, construction materials, sanitary facilities, and water management.

Camp areas require site preparation, such as shaping and leveling the tent and parking areas, stabilizing roads and intensively used areas, and installing sanitary facilities and utility lines. Camp areas are subject to heavy foot traffic and some vehicular traffic. The ratings are based on the soil properties that affect the ease of developing camp areas and the performance of the areas after development. Slope, stoniness, and depth to bedrock or a cemented pan are the main concerns affecting the development of camp areas.

The soil properties that affect the performance of the areas after development are those that influence trafficability and promote the growth of vegetation, especially in heavily used areas. For good trafficability, the surface of camp areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Picnic areas are subject to heavy foot traffic. Most vehicular traffic is confined to access roads and parking areas. The ratings are based on the soil properties that affect the ease of developing picnic areas and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of picnic areas. For good trafficability, the surface of picnic areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Playgrounds require soils that are nearly level, are free of stones, and can withstand intensive foot traffic. The ratings are based on the soil properties that affect the ease of developing playgrounds and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of playgrounds. For good trafficability, the surface of the playgrounds should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Paths and trails for hiking and horseback riding should require little or no slope modification through cutting and filling. The ratings are based on the soil properties that affect trafficability and erodibility. These properties are stoniness, depth to a water table, ponding, flooding, slope, and texture of the surface layer.

Golf fairways are subject to heavy foot traffic and some light vehicular traffic. Cutting or filling may be required. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer. The suitability of the soil for traps, tees, roughs, and greens is not considered in the ratings.

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
033CP: Clairemont-----	100	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
033OA: Obaro-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope Depth to bedrock	1.00 0.10
033OB: Obaro-----	75	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope Depth to bedrock	1.00 0.10
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
033QR: Quinlan-----	55	Very limited Depth to bedrock Slope	1.00 0.37	Very limited Depth to bedrock Slope	1.00 0.37	Very limited Depth to bedrock Slope	1.00 1.00
Woodward-----	45	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope Depth to bedrock	1.00 0.46
033QT: Quinlan-----	55	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Slope Depth to bedrock	1.00 1.00
Woodward-----	45	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Depth to bedrock	1.00 0.46
033YE: Yahola-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
077BP: Woodward-----	65	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope Depth to bedrock	1.00 0.42
Port-----	35	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding Slope	1.00 0.00
077FU: Farnum-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
077GE: Gerlane-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
077MC: Minco-----	100	Not limited		Not limited		Not limited	
077MN: Minco-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
077MO: Minco-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
077QU: Quinlan-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Slope	1.00 0.87
077SE: Shellabarger-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
077WE: Quinlan-----	50	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Slope	1.00 0.00
Woodward-----	50	Not limited		Not limited		Somewhat limited Slope	0.00
095AC: Albion-----	100	Not limited		Not limited		Somewhat limited Slope Gravel content	0.87 0.06
095DA: Dillwyn-----	60	Somewhat limited Depth to saturated zone Too sandy	0.39 0.37	Somewhat limited Too sandy Depth to saturated zone	0.37 0.19	Somewhat limited Depth to saturated zone Too sandy	0.39 0.37
Plevna-----	40	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 1.00
095FA: Farnum-----	100	Not limited		Not limited		Not limited	

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
095PB: Pratt-----	100	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy	0.37	Very limited Slope Too sandy	1.00 0.37
095SB: Shellabarger-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
095WA: Waldeck-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
097LN: Lincoln-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
097QW: Quinlan-----	55	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00
Woodward-----	45	Somewhat limited Slope	0.84	Somewhat limited Slope	0.84	Very limited Slope Depth to bedrock	1.00 0.71
097WA: Waldeck-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
151AB: Albion-----	100	Not limited		Not limited		Somewhat limited Slope Gravel content	0.13 0.06
151AO: Albion-----	100	Not limited		Not limited		Somewhat limited Slope Gravel content	0.87 0.06
151AS: Albion-----	50	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope Gravel content	1.00 0.06
Shellabarger-----	50	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope	1.00
151BC: Blanket-----	100	Not limited		Not limited		Somewhat limited Slope	0.13
151CA: Carwile-----	100	Very limited Depth to saturated zone Restricted permeability	1.00 0.94	Very limited Depth to saturated zone Restricted permeability	1.00 0.94	Very limited Depth to saturated zone Restricted permeability	1.00 0.94
151CK: Case-----	60	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope	1.00
Clark-----	40	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
151CM: Clark-----	100	Not limited		Not limited		Somewhat limited Slope	0.13
151CN: Clark-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
151CO: Clark-----	70	Not limited		Not limited		Not limited	
Ost-----	30	Not limited		Not limited		Not limited	
151FA: Farnum-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
151KP: Kanza-----	50	Very limited Flooding Depth to saturated zone Too sandy	1.00 0.98 0.92	Somewhat limited Too sandy Depth to saturated zone Flooding	0.92 0.75 0.40	Very limited Flooding Depth to saturated zone Too sandy	1.00 0.98 0.92
Plevna-----	50	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 1.00
151OS: Ost-----	100	Not limited		Not limited		Somewhat limited Slope	0.13
151PM: Pratt-----	100	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy	0.37	Very limited Slope Too sandy	1.00 0.37

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
151PN: Pratt-----	100	Somewhat limited Too sandy Slope	0.37 0.16	Somewhat limited Too sandy Slope	0.37 0.16	Very limited Slope Too sandy	1.00 0.37
151PO: Pratt-----	60	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy	0.37	Very limited Slope Too sandy	1.00 0.37
Carwile-----	40	Very limited Depth to saturated zone Restricted permeability	1.00 0.94	Very limited Depth to saturated zone Restricted permeability	1.00 0.94	Very limited Depth to saturated zone Restricted permeability	1.00 0.94
151SA: Albion-----	70	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope Gravel content	1.00 0.06
Kaski-----	30	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
151SE: Shellabarger-----	100	Not limited		Not limited		Somewhat limited Slope	0.13
1439: Crisfield-----	100	Very limited Flooding Too sandy	1.00 0.00	Somewhat limited Too sandy	0.00	Somewhat limited Too sandy	0.00
Ad: Albion-----	60	Not limited		Not limited		Somewhat limited Slope Gravel content	0.13 0.06
Shellabarger-----	40	Not limited		Not limited		Somewhat limited Slope	0.13
Ae: Albion-----	55	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope Gravel content	1.00 0.06
Shellabarger-----	45	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
AN: Albion-----	65	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope Gravel content	1.00 0.06
Shellabarger-----	35	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
As: Clairemont-----	100	Very limited Flooding Salinity	1.00 1.00	Very limited Salinity Flooding	1.00 0.40	Very limited Flooding Salinity	1.00 1.00
At: Attica-----	100	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy Slope	0.96 0.13
Ba: Blanket-----	100	Not limited		Not limited		Not limited	
Bb: Blanket-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Bc: Blanket-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Bf: Quinlan-----	60	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00
Clairemont-----	40	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
Ca: Canadian-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Cc: Case-----	50	Not limited		Not limited		Somewhat limited Slope	0.50
Clark-----	50	Not limited		Not limited		Somewhat limited Slope	0.50
Cd: Clairemont-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Cf: Clairemont-----	100	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
Ck: Clark-----	100	Not limited		Not limited		Not limited	
Fa: Farnum-----	100	Not limited		Not limited		Not limited	
Fm: Farnum-----	100	Not limited		Not limited		Not limited	
Fr: Farnum-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Fu: Farnum-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Ga: Grant-----	100	Not limited		Not limited		Not limited	
Gb: Grant-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Gc: Grant-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
GRP: Gravel Pits-----	100	Not rated		Not rated		Not rated	
INT: Aquolls-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Restricted permeability	1.00
		Restricted permeability	1.00	Restricted permeability	1.00	Depth to saturated zone	1.00
		Ponding	1.00	Ponding	1.00	Ponding	1.00
Ka: Kanza-----	100	Very limited Flooding Depth to saturated zone	1.00 0.98	Somewhat limited Too sandy Depth to saturated zone	0.92 0.75	Very limited Flooding Depth to saturated zone	1.00 0.98
		Too sandy	0.92	Flooding	0.40	Too sandy	0.92
Kf: Kingfisher-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Kv: Kingfisher-----	70	Not limited		Not limited		Somewhat limited Slope	0.00
Vernon-----	30	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39
						Content of large stones	0.03
						Slope	0.00
Kz: Kingfisher-----	60	Not limited		Not limited		Somewhat limited Slope	0.87
						Depth to bedrock	0.65
Vernon-----	40	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39	Somewhat limited Depth to bedrock	0.97
						Slope	0.87
						Restricted permeability	0.39
						Content of large stones	0.03
Ln: Lincoln-----	100	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
LNK: Lincoln-----	100	Very limited Flooding Too sandy	1.00 0.91	Somewhat limited Too sandy	0.91	Somewhat limited Too sandy Flooding	0.91 0.60
Ma: Mangum-----	100	Very limited Flooding Too clayey	1.00 0.50	Somewhat limited Too clayey Restricted permeability	0.50 0.45	Somewhat limited Flooding Too clayey	0.60 0.50
		Restricted permeability	0.45			Restricted permeability	0.45
Mg: Mangum-----	65	Very limited Flooding	1.00	Somewhat limited Too clayey	0.50	Somewhat limited Too clayey	0.50

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Drummond-----	35	Too clayey	0.50	Restricted permeability	0.45	Restricted permeability	0.45
		Restricted permeability	0.45				
		Very limited Flooding	1.00	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45
		Restricted permeability	0.45				
Mn: Minco-----	100	Not limited		Not limited		Not limited	
Na: Naron-----	100	Not limited		Not limited		Not limited	
Nb: Naron-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Os: Ost-----	100	Not limited		Not limited		Not limited	
Ot: Ost-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Pa: Pond Creek-----	100	Not limited		Not limited		Not limited	
Pd: Pond Creek-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Ph: Dale-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Pk: Buttermilk-----	100	Very limited Flooding Salinity	1.00 0.13	Somewhat limited Salinity	0.13	Somewhat limited Salinity	0.13
Ps: Pratt-----	100	Somewhat limited Too sandy Slope	0.37 0.00	Somewhat limited Too sandy Slope	0.37 0.00	Very limited Slope Too sandy	1.00 0.37
Pt: Pratt-----	70	Somewhat limited Too sandy Slope	0.37 0.16	Somewhat limited Too sandy Slope	0.37 0.16	Very limited Slope Too sandy	1.00 0.37
Tivoli-----	30	Somewhat limited Too sandy Slope	0.92 0.16	Somewhat limited Too sandy Slope	0.92 0.16	Very limited Slope Too sandy	1.00 0.92
Qn: Quinlan-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Slope	1.00 0.00
Qw: Quinlan-----	60	Very limited Depth to bedrock Slope	1.00 0.16	Very limited Depth to bedrock Slope	1.00 0.16	Very limited Depth to bedrock Slope	1.00 1.00
Woodward-----	40	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Rb: Knoco-----	70	Very limited Depth to bedrock Slope Sodium content Too clayey Restricted permeability	1.00 1.00 1.00 0.50 0.45	Very limited Depth to bedrock Slope Sodium content Too clayey Restricted permeability	1.00 1.00 1.00 0.50 0.45	Very limited Slope Depth to bedrock Sodium content Too clayey Restricted permeability	1.00 1.00 1.00 0.50 0.45
Rock Outcrop-----	30	Not rated		Not rated		Not rated	
Sb: Shellabarger-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
SBB: Shellabarger-----	100	Not limited		Not limited		Not limited	
Sc: Shellabarger-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
Tv: Tivoli-----	100	Very limited Too sandy Slope	1.00 0.84	Very limited Too sandy Slope	1.00 0.84	Very limited Too sandy Slope	1.00 1.00
Vn: Vernon-----	100	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39	Somewhat limited Depth to bedrock	0.97

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Vr: Vernon-----	100	Somewhat limited Restricted permeability Slope	0.39 0.16	Somewhat limited Restricted permeability Slope	0.39 0.16	Slope	0.50
						Restricted permeability	0.39
						Content of large stones	0.03
						Very limited Slope	1.00
Vs: Knoco-----	50	Very limited Depth to bedrock Sodium content Too clayey Restricted permeability	1.00 1.00 0.50 0.45	Very limited Depth to bedrock Sodium content Too clayey Restricted permeability	1.00 1.00 0.50 0.45	Depth to bedrock	0.97
						Restricted permeability	0.39
						Content of large stones	0.03
						Very limited	1.00
Vernon-----	50	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39	Depth to bedrock	1.00
						Sodium content	1.00
						Too clayey	0.50
						Restricted permeability	0.45
W: Water-----	100	Not rated		Not rated		Slope	0.13
						Somewhat limited Depth to bedrock	0.42
						Restricted permeability	0.39
						Slope	0.13
Wa: Waldeck-----	100	Very limited Flooding	1.00	Not limited		Content of large stones	0.03
						Not rated	
						Somewhat limited Flooding	0.60
Wo: Woodward-----	70	Not limited		Not limited			
						Somewhat limited Slope	0.00
						Very limited	
						Depth to bedrock	1.00
Quinlan-----	30	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Slope	0.00
						Somewhat limited Slope	0.87
						Very limited	
Ws: Woodward-----	65	Not limited		Not limited		Depth to bedrock	1.00
						Slope	0.87
						Very limited Depth to bedrock	1.00
Quinlan-----	35	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Slope	0.87
						Somewhat limited Flooding	0.60
Ya: Yahola-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
						Somewhat limited Flooding	0.60
Ze: Zenda-----	100	Very limited Flooding	1.00	Not limited			

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
033CP: Clairemont-----	100	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
033OA: Obaro-----	100	Not limited		Somewhat limited Depth to bedrock Slope	0.10 0.04
033OB: Obaro-----	75	Not limited		Somewhat limited Slope Depth to bedrock	0.96 0.10
Rock Outcrop-----	25	Not rated		Not rated	
033QR: Quinlan-----	55	Not limited		Very limited Depth to bedrock Droughty Slope	1.00 0.74 0.37
Woodward-----	45	Not limited		Somewhat limited Depth to bedrock Slope	0.46 0.37
033QT: Quinlan-----	55	Somewhat limited Slope	0.92	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.74
Woodward-----	45	Somewhat limited Slope	0.18	Very limited Slope Depth to bedrock	1.00 0.46
033YE: Yahola-----	100	Not limited		Somewhat limited Flooding	0.60
077BP: Woodward-----	65	Not limited		Somewhat limited Depth to bedrock Slope	0.42 0.37
Port-----	35	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
077FU: Farnum-----	100	Not limited		Not limited	
077GE: Gerlane-----	100	Not limited		Somewhat limited Flooding	0.60
077MC: Minco-----	100	Not limited		Not limited	
077MN: Minco-----	100	Not limited		Not limited	
077MO: Minco-----	100	Not limited		Not limited	
077QU: Quinlan-----	100	Not limited		Very limited Depth to bedrock Droughty	1.00 1.00
077SE: Shellabarger-----	100	Not limited		Not limited	
077WE: Quinlan-----	50	Not limited		Very limited Depth to bedrock Droughty	1.00 1.00
Woodward-----	50	Not limited		Somewhat limited Depth to bedrock	0.90
095AC: Albion-----	100	Not limited		Not limited	
095DA: Dillwyn-----	60	Somewhat limited Too sandy	0.37	Somewhat limited Droughty Depth to saturated zone	0.22 0.19
Plevna-----	40	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 1.00
095FA: Farnum-----	100	Not limited		Not limited	
095PB: Pratt-----	100	Somewhat limited Too sandy	0.37	Not limited	
095SB: Shellabarger-----	100	Not limited		Not limited	

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
095WA: Waldeck-----	100	Not limited		Somewhat limited Flooding	0.60
097LN: Lincoln-----	100	Not limited		Somewhat limited Droughty Flooding	0.80 0.60
097QW: Quinlan-----	55	Somewhat limited Slope	0.02	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.62
Woodward-----	45	Not limited		Somewhat limited Slope Depth to bedrock	0.84 0.71
097WA: Waldeck-----	100	Not limited		Somewhat limited Flooding	0.60
151AB: Albion-----	100	Not limited		Not limited	
151AO: Albion-----	100	Not limited		Not limited	
151AS: Albion-----	50	Not limited		Somewhat limited Slope	0.37
Shellabarger-----	50	Not limited		Somewhat limited Slope	0.37
151BC: Blanket-----	100	Not limited		Not limited	
151CA: Carwile-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
151CK: Case-----	60	Not limited		Somewhat limited Slope	0.37
Clark-----	40	Not limited		Somewhat limited Slope	0.16
151CM: Clark-----	100	Not limited		Not limited	
151CN: Clark-----	100	Not limited		Not limited	
151CO: Clark-----	70	Not limited		Not limited	
Ost-----	30	Not limited		Not limited	
151FA: Farnum-----	100	Not limited		Not limited	
151KP: Kanza-----	50	Somewhat limited Too sandy Depth to saturated zone Flooding	0.92 0.44 0.40	Very limited Flooding Depth to saturated zone Droughty	1.00 0.75 0.02
Plevna-----	50	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 1.00
151OS: Ost-----	100	Not limited		Not limited	
151PM: Pratt-----	100	Somewhat limited Too sandy	0.37	Not limited	
151PN: Pratt-----	100	Somewhat limited Too sandy	0.37	Somewhat limited Slope	0.16
151PO: Pratt-----	60	Somewhat limited Too sandy	0.37	Not limited	
Carwile-----	40	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
151SA: Albion-----	70	Not limited		Somewhat limited Slope	0.37
Kaski-----	30	Not limited		Somewhat limited Flooding	0.60
151SE: Shellabarger-----	100	Not limited		Not limited	
1439: Crisfield-----	100	Somewhat limited		Somewhat limited	

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
		Too sandy	0.00	Droughty	0.31
Ad:					
Albion-----	60	Not limited		Not limited	
Shellabarger-----	40	Not limited		Not limited	
Ae:					
Albion-----	55	Not limited		Somewhat limited Slope	0.16
Shellabarger-----	45	Not limited		Somewhat limited Slope	0.16
AED:					
Arents, Earthen Dam-	100	Not rated		Not rated	
AN:					
Albion-----	65	Not limited		Somewhat limited Slope	0.16
Shellabarger-----	35	Not limited		Somewhat limited Slope	0.16
As:					
Clairemont-----	100	Somewhat limited Flooding	0.40	Very limited Flooding Salinity	1.00 1.00
At:					
Attica-----	100	Somewhat limited Too sandy	0.96	Not limited	
Ba:					
Blanket-----	100	Not limited		Not limited	
Bb:					
Blanket-----	100	Not limited		Not limited	
Bc:					
Blanket-----	100	Not limited		Not limited	
Bf:					
Quinlan-----	60	Very limited Slope	1.00	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.74
Clairemont-----	40	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
Ca:					
Canadian-----	100	Not limited		Not limited	
Cc:					
Case-----	50	Not limited		Not limited	
Clark-----	50	Not limited		Not limited	
Cd:					
Clairemont-----	100	Not limited		Somewhat limited Flooding	0.60
Cf:					
Clairemont-----	100	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
Ck:					
Clark-----	100	Not limited		Not limited	
Fa:					
Farnum-----	100	Not limited		Not limited	
Fm:					
Farnum-----	100	Not limited		Not limited	
Fr:					
Farnum-----	100	Not limited		Not limited	
Fu:					
Farnum-----	100	Not limited		Not limited	
Ga:					
Grant-----	100	Not limited		Not limited	
Gb:					
Grant-----	100	Not limited		Not limited	
Gc:					
Grant-----	100	Not limited		Not limited	
GRP:					
Gravel Pits-----	100	Not rated		Not rated	
INT:					
Aquolls-----	100	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
Ka:					
Kanza-----	100	Somewhat limited Too sandy Depth to saturated zone Flooding	0.92 0.44 0.40	Very limited Flooding Depth to saturated zone Droughty	1.00 0.75 0.04

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Kf: Kingfisher-----	100	Not limited		Somewhat limited Depth to bedrock	0.65
Kv: Kingfisher-----	70	Not limited		Somewhat limited Depth to bedrock	0.65
Vernon-----	30	Not limited		Somewhat limited Depth to bedrock	0.97
				Droughty	0.11
				Content of large stones	0.03
Kz: Kingfisher-----	60	Not limited		Somewhat limited Depth to bedrock	0.65
Vernon-----	40	Not limited		Somewhat limited Depth to bedrock	0.97
				Droughty	0.11
				Content of large stones	0.03
Ln: Lincoln-----	100	Somewhat limited Flooding	0.40	Very limited Flooding Droughty	1.00 0.89
LNN: Lincoln-----	100	Somewhat limited Too sandy	0.91	Somewhat limited Droughty Flooding	0.92 0.60
Ma: Mangum-----	100	Somewhat limited Too clayey	0.50	Very limited Too clayey Flooding	1.00 0.60
Mg: Mangum-----	65	Somewhat limited Too clayey	0.50	Very limited Too clayey	1.00
Drummond-----	35	Not limited		Not limited	
Mn: Minco-----	100	Not limited		Not limited	
Na: Naron-----	100	Not limited		Not limited	
Nb: Naron-----	100	Not limited		Not limited	
Os: Ost-----	100	Not limited		Not limited	
Ot: Ost-----	100	Not limited		Not limited	
Pa: Pond Creek-----	100	Not limited		Not limited	
Pd: Pond Creek-----	100	Not limited		Not limited	
Ph: Dale-----	100	Not limited		Not limited	
Pk: Buttermilk-----	100	Not limited		Somewhat limited Salinity	0.13
Ps: Pratt-----	100	Somewhat limited Too sandy	0.37	Somewhat limited Slope	0.00
Pt: Pratt-----	70	Somewhat limited Too sandy	0.37	Somewhat limited Slope	0.16
Tivoli-----	30	Somewhat limited Too sandy	0.92	Somewhat limited Droughty Slope	0.98 0.16
Qn: Quinlan-----	100	Not limited		Very limited Depth to bedrock Droughty	1.00 0.74
Qw: Quinlan-----	60	Not limited		Very limited Depth to bedrock Droughty Slope	1.00 0.74 0.16
Woodward-----	40	Not limited		Somewhat limited Slope	0.16
Rb: Knoco-----	70	Very limited Slope Too clayey	1.00 0.50	Very limited Depth to bedrock Droughty Slope Too clayey Sodium content	1.00 1.00 1.00 1.00 1.00

RECREATIONAL INTERPRETATIONS--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Rock Outcrop-----	30	Not rated		Not rated	
Sb: Shellabarger-----	100	Not limited		Not limited	
SBB: Shellabarger-----	100	Not limited		Not limited	
Sc: Shellabarger-----	100	Not limited		Not limited	
Tv: Tivoli-----	100	Very limited Too sandy	1.00	Very limited Droughty Slope	1.00 0.84
Vn: Vernon-----	100	Not limited		Somewhat limited Depth to bedrock Droughty Content of large stones	0.97 0.11 0.03
Vr: Vernon-----	100	Not limited		Somewhat limited Depth to bedrock Slope Droughty Content of large stones	0.97 0.16 0.11 0.03
Vs: Knoco-----	50	Somewhat limited Too clayey	0.50	Very limited Depth to bedrock Droughty Too clayey Sodium content	1.00 1.00 1.00 1.00
Vernon-----	50	Not limited		Somewhat limited Depth to bedrock Droughty Content of large stones	0.42 0.11 0.03
W: Water-----	100	Not rated		Not rated	
Wa: Waldeck-----	100	Not limited		Somewhat limited Flooding	0.60
Wo: Woodward-----	70	Not limited		Not limited	
Quinlan-----	30	Not limited		Very limited Depth to bedrock Droughty	1.00 0.74
Ws: Woodward-----	65	Not limited		Not limited	
Quinlan-----	35	Not limited		Very limited Depth to bedrock Droughty	1.00 0.74
Ya: Yahola-----	100	Not limited		Somewhat limited Flooding	0.60
Ze: Zenda-----	100	Not limited		Somewhat limited Flooding	0.60

WILDLIFE INTERPRETATIONS
Barber County, Kansas

Use and Explanation of Wildlife Interpretations

Soils directly affect the kind and amount of vegetation that is available to wildlife as food and cover. They also affect the development of water impoundments. The kind and abundance of wildlife that populate an area depend largely on the amount and distribution of food, cover, water, and living space. If any one of these elements is missing, inadequate, or inaccessible, wildlife will be scarce or will not inhabit the area. If the soils have the potential, wildlife habitat can be created or improved by planting appropriate vegetation, properly managing the existing plant cover, and fostering the natural establishment of desirable plants.

In the Wildlife Interpretations table, the soils in the survey area are rated according to their potential for providing habitat for various kinds of wildlife. This information can be used in planning parks, wildlife refuges, nature study areas, and other developments for wildlife; in selecting soils that are suitable for establishing, improving, or maintaining specific elements of wildlife habitat; and in determining the intensity of management needed for each element of the habitat.

Suitability Ratings

The potential of the soil is rated good, fair, poor, or very poor.

Good - means that the element of wildlife habitat or the kind of habitat is easily created, improved, or maintained. Few or no limitations affect management, and satisfactory results can be expected if the soil is used for the designated purpose.

Fair - means that the element of wildlife habitat or kind of habitat can be created, improved, or maintained in most places. Moderately intensive management is required for satisfactory results.

Poor - means that limitations are severe for the designated element or kind of wildlife habitat. Habitat can be created, improved, or maintained in most places, but management is difficult and requires intensive effort.

Very Poor - means that limitations are very severe for the designated element or kind of wildlife habitat. Habitat is difficult to create, improve, or maintain in most places, and management is difficult and requires intensive effort.

Description of Wildlife Habitat Elements

Openland habitat consists of croplands, pastures, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. The kind of wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, killdeer, cottontail rabbit, red fox, and coyote.

Woodland habitat consists of hardwood or conifers, or a mixture of these and associated grasses, legumes and wild herbaceous plants. Examples of wildlife attracted to this habitat are wild turkey, thrushes, woodpeckers, owl, tree squirrels, raccoon, and deer.

Wetland habitat consists of water-tolerant plants in open, marshy or swampy, shallow water areas. Examples of wildlife attracted to this habitat are ducks, geese, herons, bitterns, rails, kingfishers, shorebirds, muskrat, mink, and beaver.

The elements of wildlife habitat are described in the following paragraphs.

Grain and seed crops are domestic grains and seed-producing herbaceous plants. Soil properties and features that affect the growth of grain and seed crops are depth of the root zone, texture of the surface layer, available water capacity, wetness, slope, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of grain and seed crops are corn, wheat, oats, and barley.

Grasses and legumes are domestic perennial grasses and herbaceous legumes. Soil properties and features that affect the growth of grasses and legumes are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, flooding, and slope. Soil temperature and soil moisture also are considerations. Examples of grasses and legumes are fescue, lovegrass, bromegrass, clover, and alfalfa.

Wild herbaceous plants are native or naturally established grasses and forbs, including weeds. Soil properties and features that affect the growth of these plants are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of wild herbaceous plants are bluestem, goldenrod, beggarweed, wheatgrass, and grama.

Hardwood trees and woody understory produce nuts or other fruit, buds, catkins, twigs, bark, and foliage. Soil properties and features that affect the growth of hardwood trees and shrubs are depth of the root zone, available water capacity, and wetness. Examples of these plants are oak, poplar, cherry, sweetgum, apple, hawthorn, dogwood, hickory, blackberry, and blueberry. Examples of fruit-producing shrubs that are suitable for planting on soils rated good are Russian-olive, autumn-olive, and crabapple.

Coniferous plants furnish browse and seeds. Soil properties and features that affect the growth of coniferous trees, shrubs, and ground cover are depth of the root zone, available water capacity, and wetness. Examples of coniferous plants are pine, spruce, fir, cedar, and juniper.

Shrubs are bushy woody plants that produce fruit, buds, twigs, bark, and foliage. Soil properties and features that affect the growth of shrubs are depth of the root zone, available water capacity, salinity, and soil moisture. Examples of shrubs are fragrant sumac, chokecherry, American plum, sand plum, and gorden currant.

Wetland plants are annual and perennial wild herbaceous plants that grow on moist or wet sites. Submerged or floating aquatic plants are excluded. Soil properties and features affecting wetland plants are texture of the surface layer, wetness, reaction, salinity, slope, and surface stoniness. Examples of wetland plants are smartweed, wild millet, saltgrass, cordgrass, rushes, sedges, and cattails.

Shallow water areas have an average depth of less than 5 feet. Some are naturally wet areas. Others are created by dams, levees, or other water-control structures. Soil properties and features affecting shallow water areas are depth to bedrock, wetness, surface stoniness, slope, and permeability. Examples of shallow water areas are marshes, waterfowl feeding areas, and ponds.

The habitat for various kinds of wildlife is described in the following paragraphs.

Habitat for openland wildlife consists of cropland, pasture, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. Wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, cottontail, red fox and coyote.

Habitat for woodland wildlife consists of areas of deciduous and/or coniferous plants and associated grasses, legumes, and wild herbaceous plants. Wildlife attracted to these areas include wild turkey, thrushes, woodpeckers, squirrels, gray fox, raccoon, and deer.

Habitat for wetland wildlife consists of open, marshy or swampy shallow water areas. Some of the wildlife attracted to such areas are ducks, geese, herons, shore birds, muskrat, mink, and beaver.

Habitat for rangeland wildlife consists of areas of shrubs and wild herbaceous plants. Wildlife attracted to rangeland include antelope, deer, cottontail rabbit, prairie chicken, meadowlark, quail, and pheasant.

WILDLIFE INTERPRETATIONS
Barber County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
033CP: CLAIREMONT-----	Very poor	Poor	Fair	---	Very poor	Good	Very poor	Very poor	Poor	---	---	Fair
033OA: OBARO-----	Poor	Fair	Fair	---	Very poor	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
033OB: OBARO-----	Poor	Fair	Fair	---	Very poor	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
ROCK OUTCROP----	---	---	---	---	---	---	---	---	---	---	---	---
033QR: QUINLAN-----	Poor	Poor	Fair	---	---	Poor	Very poor	Very poor	Fair	---	Very poor	Poor
WOODWARD-----	Fair	Good	Good	---	---	Fair	Very poor	Very poor	Good	---	Very poor	Fair
033QT: QUINLAN-----	Poor	Poor	Fair	---	---	Poor	Very poor	Very poor	Fair	---	Very poor	Poor
WOODWARD-----	Poor	Fair	Good	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
033YE: YAHOLA-----	Good	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor	Good
077BP: WOODWARD-----	Fair	Good	Good	---	---	Fair	Very poor	Very poor	Good	---	Very poor	Fair
PORT-----	Poor	Fair	Fair	---	---	Good	Poor	Very poor	Fair	---	Very poor	Fair
077FU: FARNUM-----	Fair	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
077GE: GERLANE-----	Good	Good	Good	---	---	Fair	Poor	Poor	Good	---	Poor	Fair
077MC: MINCO-----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
077MN: MINCO-----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
077MO: MINCO-----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
077QU: QUINLAN-----	Poor	Poor	Fair	---	---	Poor	Poor	Very poor	Fair	---	Very poor	Poor
077SE: SHELLABARGER----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
077WE: QUINLAN-----	Poor	Poor	Fair	---	---	Poor	Poor	Very poor	Fair	---	Very poor	Poor
WOODWARD-----	Fair	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
095AC: ALBION-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
095DA: DILLWYN-----	Poor	Fair	Good	---	---	Fair	Fair	Fair	Fair	---	Fair	Fair
PLEVNA-----	Poor	Fair	Fair	---	---	Fair	Good	Good	Fair	---	Good	Fair
095FA: FARNUM-----	Good	Good	Good	---	---	Good	Poor	Poor	Good	---	Poor	Good

WILDLIFE INTERPRETATIONS--Continued
Barber County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
095PB: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
095SB: SHELLABARGER----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
095WA: WALDECK-----	Fair	Good	Good	---	---	Good	Fair	Fair	Good	---	Fair	Good
097LN: LINCOLN-----	Fair	Fair	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
097QW: QUINLAN-----	Poor	Poor	Fair	---	---	Poor	Very poor	Very poor	Fair	---	Very poor	Poor
WOODWARD-----	Fair	Good	Good	---	---	Fair	Very poor	Very poor	Good	---	Very poor	Fair
097WA: WALDECK-----	Fair	Good	Good	---	---	Good	Fair	Fair	Good	---	Fair	Good
151AB: ALBION-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
151AO: ALBION-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
151AS: ALBION-----	Poor	Fair	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
SHELLABARGER----	Poor	Fair	Good	---	---	Good	Very poor	Very poor	Fair	---	Very poor	Good
151BC: BLANKET-----	Good	Good	Fair	---	Good	Good	Poor	Very poor	Good	---	Very poor	Fair
151CA: CARWILE-----	Fair	Good	Good	---	---	Good	Good	Fair	Good	---	Fair	Good
151CK: CASE-----	Poor	Fair	Fair	---	---	Fair	Poor	Very poor	Fair	---	Very poor	Fair
CLARK-----	Poor	Fair	Fair	Fair	Fair	Fair	Poor	Very poor	Fair	---	Very poor	Fair
151CM: CLARK-----	Fair	Good	Fair	Fair	Fair	Fair	Poor	Very poor	Fair	---	Very poor	Fair
151CN: CLARK-----	Fair	Good	Fair	Fair	Fair	Fair	Poor	Very poor	Fair	---	Very poor	Fair
151CO: CLARK-----	Good	Good	Fair	Fair	Fair	Fair	Poor	Very poor	Good	---	Very poor	Fair
OST-----	Good	Good	Fair	---	---	Fair	Poor	Poor	Good	---	Poor	Fair
151FA: FARNUM-----	Fair	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
151KP: KANZA-----	Very poor	Poor	Fair	---	---	Fair	Fair	Fair	Poor	---	Fair	Fair
PLEVNA-----	Poor	Fair	Fair	---	---	Fair	Good	Good	Fair	---	Good	Fair
151OS: OST-----	Good	Good	Fair	---	---	Fair	Poor	Poor	Good	---	Poor	Fair
151PM: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair

WILDLIFE INTERPRETATIONS--Continued
Barber County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
151PN: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
151PO: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
CARWILE-----	Fair	Good	Good	---	---	Good	Good	Fair	Good	---	Fair	Good
151SA: ALBION-----	Poor	Fair	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
KASKI-----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
151SE: SHELLABARGER----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
1439: CRISFIELD-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
Ad: ALBION-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
SHELLABARGER----	Fair	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
Ae: ALBION-----	Poor	Fair	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
SHELLABARGER----	Poor	Fair	Good	---	---	Good	Very poor	Very poor	Fair	---	Very poor	Good
AED: ARENTS, EARTHEN DAM-----	---	---	---	---	---	---	---	---	---	---	---	---
AN: ALBION-----	Poor	Fair	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
SHELLABARGER----	Poor	Fair	Good	---	---	Good	Very poor	Very poor	Fair	---	Very poor	Good
As: CLAIREMONT-----	Poor	Poor	Poor	---	---	Poor	Poor	Poor	Poor	---	Poor	Poor
At: ATTICA-----	Fair	Fair	Good	---	---	Fair	Poor	Very poor	Fair	---	Very poor	Fair
Ba: BLANKET-----	Good	Good	Fair	---	Good	Good	Poor	Very poor	Good	---	Very poor	Fair
Bb: BLANKET-----	Good	Good	Fair	---	Good	Good	Poor	Very poor	Good	---	Very poor	Fair
Bc: BLANKET-----	Fair	Good	Fair	---	Good	Good	Poor	Very poor	Fair	---	Very poor	Fair
Bf: QUINLAN-----	Very poor	Very poor	Fair	---	---	Poor	Very poor	Very poor	Poor	---	Very poor	Poor
CLAIREMONT-----	Very poor	Poor	Fair	---	Very poor	Good	Very poor	Very poor	Poor	---	---	Fair
Ca: CANADIAN-----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
Cc: CASE-----	Fair	Good	Fair	---	---	Fair	Poor	Very poor	Fair	---	Very poor	Fair

WILDLIFE INTERPRETATIONS--Continued
Barber County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
CLARK-----	Fair	Good	Fair	Fair	Fair	Fair	Poor	Very poor	Fair	---	Very poor	Fair
Cd: CLAIREMONT-----	Good	Good	Fair	---	Very poor	Good	Very poor	Very poor	Good	---	---	Fair
Cf: CLAIREMONT-----	Very poor	Poor	Fair	---	Very poor	Good	Very poor	Very poor	Poor	---	---	Fair
Ck: CLARK-----	Fair	Good	Fair	Fair	Fair	Fair	Poor	Very poor	Fair	---	Very poor	Fair
Fa: FARNUM-----	Good	Good	Good	---	---	Good	Poor	Poor	Good	---	Poor	Good
Fm: FARNUM-----	Good	Good	Good	---	---	Good	Poor	Poor	Good	---	Poor	Good
Fr: FARNUM-----	Good	Good	Good	---	---	Good	Poor	Poor	Good	---	Poor	Good
Fu: FARNUM-----	Good	Good	Good	---	---	Good	Poor	Poor	Good	---	Poor	Good
Ga: GRANT-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
Gb: GRANT-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
Gc: GRANT-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
GRP: GRAVEL PITS-----	---	---	---	---	---	---	---	---	---	---	---	---
INT: AQUOLLS-----	---	---	---	---	---	---	---	---	---	---	---	---
Ka: KANZA-----	Very poor	Poor	Fair	---	---	Fair	Fair	Fair	Poor	---	Fair	Fair
Kf: KINGFISHER-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
Kv: KINGFISHER-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
VERNON-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Kz: KINGFISHER-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
VERNON-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Ln: LINCOLN-----	Poor	Fair	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
LNN: LINCOLN-----	Fair	Fair	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Ma: MANGUM-----	Good	Good	Fair	---	Very poor	Fair	Poor	Poor	Good	---	Poor	Fair
Mg: MANGUM-----	Good	Good	Fair	---	Very poor	Fair	Poor	Poor	Good	---	Poor	Fair
DRUMMOND-----	Poor	Fair	Fair	---	Poor	Poor	Fair	Fair	Fair	---	Fair	Poor

WILDLIFE INTERPRETATIONS--Continued
Barber County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
Mn: MINCO-----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
Na: NARON-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
Nb: NARON-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
Os: OST-----	Good	Good	Fair	---	---	Fair	Poor	Poor	Good	---	Poor	Fair
Ot: OST-----	Good	Good	Fair	---	---	Fair	Poor	Poor	Good	---	Poor	Fair
Pa: POND CREEK-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
Pd: POND CREEK-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
Ph: DALE-----	Good	Good	Fair	---	---	Good	Poor	Very poor	Good	---	Very poor	Fair
Pk: BUTTERMILK-----	Fair	Fair	Fair	---	---	Fair	Poor	Very poor	Fair	---	Very poor	Fair
Ps: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Pt: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
TIVOLI-----	Poor	Poor	Fair	---	---	Poor	Very poor	Very poor	Poor	---	Very poor	Poor
Qn: QUINLAN-----	Poor	Poor	Fair	---	---	Poor	Poor	Very poor	Fair	---	Very poor	Poor
Qw: QUINLAN-----	Poor	Poor	Fair	---	---	Poor	Very poor	Very poor	Fair	---	Very poor	Poor
WOODWARD-----	Fair	Good	Good	---	---	Fair	Very poor	Very poor	Good	---	Very poor	Fair
Rb: KNOCO-----	Very poor	Very poor	Poor	---	---	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor
ROCK OUTCROP----	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor
Sb: SHELLABARGER----	Fair	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
SBB: SHELLABARGER----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
Sc: SHELLABARGER----	Fair	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
Tv: TIVOLI-----	Poor	Poor	Fair	---	---	Poor	Very poor	Very poor	Poor	---	Very poor	Poor
Vn: VERNON-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair

WILDLIFE INTERPRETATIONS--Continued
Barber County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
Vr: VERNON-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Vs: KNOCO-----	Very poor	Very poor	Poor	---	---	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor
VERNON-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
W: WATER-----	---	---	---	---	---	---	---	---	---	---	---	---
Wa: WALDECK-----	Fair	Good	Good	---	---	Good	Fair	Fair	Good	---	Fair	Good
Wo: WOODWARD-----	Fair	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
QUINLAN-----	Poor	Poor	Fair	---	---	Poor	Poor	Very poor	Fair	---	Very poor	Poor
Ws: WOODWARD-----	Fair	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
QUINLAN-----	Poor	Poor	Fair	---	---	Poor	Poor	Very poor	Fair	---	Very poor	Poor
Ya: YAHOLA-----	Good	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor	Good
Ze: ZENDA-----	Fair	Good	Good	---	---	Good	Fair	Fair	Good	---	Fair	Good

YIELDS PER ACRE OF PASTURE AND HAYLAND
Barber County, Kansas

Use and Explanation of Pastureland and Hayland Interpretations

This subsection provides information concerning the suitability of soils for the production of pasture and hayland. This subsection may contain pasture and hayland suitability groupings, land capability and yield estimates, yield estimates for individual grasses or legumes, or other information pertaining to the production of forage.

Pasture and Hayland Suitability Groupings

Soils are placed in pasture and hayland groups according to their suitability for the production of forage. The soils in each group are enough alike to be suited to the same grasses or legumes, to have similar limitations and hazards, to require similar management, and to have similar productivity and other responses to management. Thus, the pasture and hayland suitability group is a convenient way of grouping the soils for their management. If used, these groupings are identified and described in other reports in the subsection.

Yield Estimates

The average yields per acre that can be expected of the principal pasture or hayland crops, under a high level of management, are presented in this subsection. In any given year, yields may be higher or lower than those indicated in the tables because of variations in rainfall or other climatic factors. The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations are also considered.

Under good management, proper grazing is essential for the production of high quality forage, stand survival, and erosion control. Proper grazing helps plants maintain sufficient and generally vigorous top growth during the growing season. Brush control is essential in many areas, and weed control generally is needed. Rotation grazing and renovation are also important management practices.

The Pasture and Hayland table show yield estimates in tons per acre and animal unit months for pasture and hayland groups. An animal unit month is the amount of forage required by one animal unit (AU) for 30 days. On animal unit (AU) is one (1000 pound) mature cow and a calf up to weaning age (usually six months of age) or their equivalent. The Natural Resources Conservation Service uses 900 pounds of air dry forage as the amount needed to meet this requirement. To maintain a healthy and vigorous plant community, the degree of use should never be greater than 50 percent. Therefore only 25 percent of the total biomass grown is considered consumed by the grazing animal. Animal Unit Months can be converted to air dry pounds per acre production by multiplying the AUM by 30 days, then by 30 pounds per day, and then by four. This figure is the amount of total forage production.

Planners of management systems for individual fields or farms should consider the detailed information given in the description of each soil in the Nontechnical Description section. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
033CP: Clairemont-----	6w	---	---	---
033OA: Obaro-----	6e	---	---	---
033OB: Obaro-----	6e	---	---	---
Rock Outcrop-----	8	---	---	---
033QR: Quinlan-----	6e	---	---	---
Woodward-----	6e	---	---	---
033QT: Quinlan-----	7e	---	---	---
Woodward-----	6e	---	---	---
033YE: Yahola-----	2w	---	2.00	---
077BP: Woodward-----	6e	---	---	---
Port-----	5w	---	---	---
077FU: Farnum-----	3e	---	3.00	---
077GE: Gerlane-----	2e	---	3.40	---
077MC: Minco-----	1	---	3.50	---
077MN: Minco-----	2e	---	3.00	---
077MO: Minco-----	3e	---	2.50	---
077QU: Quinlan-----	4e	---	---	---
077SE: Shellabarger-----	2e	---	2.20	6.50
077WE: Quinlan-----	3e	---	---	---
Woodward-----	2e	---	---	---
095AC: Albion-----	4e	---	1.50	---
095DA: Dillwyn-----	4w	---	---	---
Plevna-----	5w	---	---	---
095FA: Farnum-----	3e	2e	2.50	6.50
095PB: Pratt-----	4e	3e	---	5.50
095SB: Shellabarger-----	2e	---	2.20	6.50
095WA: Waldeck-----	3w	---	3.50	5.00
097LN: Lincoln-----	6w	---	---	---
097QW: Quinlan-----	6e	---	---	---

YIELDS PER ACRE OF PASTURE AND HAYLAND--Continued
Barber County, Kansas

PAGE 3 OF 6

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Woodward-----	6e	---	---	---
097WA: Waldeck-----	3w	---	3.50	5.00
151AB: Albion-----	3e	---	2.00	---
151AO: Albion-----	4e	---	1.50	---
151AS: Albion-----	6e	---	---	---
Shellabarger-----	6e	---	---	---
151BC: Blanket-----	3e	---	---	---
151CA: Carwile-----	2w	---	---	---
151CK: Case-----	6e	---	---	---
Clark-----	6e	---	---	---
151CM: Clark-----	3e	---	---	---
151CN: Clark-----	3e	---	---	---
151CO: Clark-----	2c	---	---	---
Ost-----	2c	---	---	---
151FA: Farnum-----	4e	---	---	---
151KP: Kanza-----	5w	---	---	---
Plevna-----	5w	---	---	---
151OS: Ost-----	2e	---	---	---
151PM: Pratt-----	4e	3e	---	5.50
151PN: Pratt-----	6e	---	---	---
151PO: Pratt-----	4e	3e	---	5.50
Carwile-----	2w	---	---	---
151SA: Albion-----	6e	---	---	---
Kaski-----	2w	---	3.00	6.50
151SE: Shellabarger-----	2e	---	2.20	6.50
1439: Crisfield-----	3s	---	2.50	---
Ad: Albion-----	4e	---	1.50	---
Shellabarger-----	3e	---	2.00	6.00
Ae: Albion-----	6e	---	---	---
Shellabarger-----	6e	---	---	---

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
AED: Arents, Earthen Dam-----	8	---	---	---
AN: Albion-----	6e	---	---	---
Shellabarger-----	6e	---	---	---
As: Clairemont-----	6s	---	---	---
At: Attica-----	3e	---	2.50	6.00
Ba: Blanket-----	2c	---	---	---
Bb: Blanket-----	2e	---	---	---
Bc: Blanket-----	3e	---	---	---
Bf: Quinlan-----	7e	---	---	---
Clairemont-----	5w	---	---	---
Ca: Canadian-----	2e	---	2.50	---
Cc: Case-----	4e	---	---	---
Clark-----	4e	---	---	---
Cd: Clairemont-----	2w	2w	---	---
Cf: Clairemont-----	5w	---	---	---
Ck: Clark-----	3e	---	---	---
Fa: Farnum-----	2e	1	3.00	7.00
Fm: Farnum-----	2c	1	3.00	7.00
Fr: Farnum-----	2e	2e	3.00	6.50
Fu: Farnum-----	3e	---	2.50	---
Ga: Grant-----	1	---	2.80	---
Gb: Grant-----	2e	---	2.30	---
Gc: Grant-----	3e	---	2.00	---
GRP: Gravel Pits-----	---	---	---	---
INT: Aquolls-----	5w	---	---	---
Ka: Kanza-----	5w	---	---	---
Kf: Kingfisher-----	2e	---	---	---
Kv: Kingfisher-----	2e	---	---	---

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Vernon-----	3e	---	---	---
Kz: Kingfisher-----	3e	---	---	---
Vernon-----	4e	---	---	---
Ln: Lincoln-----	6w	---	---	---
LNN: Lincoln-----	6w	---	---	---
Ma: Mangum-----	3w	3w	---	---
Mg: Mangum-----	3s	2s	---	---
Drummond-----	5s	---	---	---
Mn: Minco-----	2e	---	2.50	---
Na: Naron-----	2e	1	2.50	7.00
Nb: Naron-----	3e	2e	2.50	6.50
Os: Ost-----	2c	---	---	---
Ot: Ost-----	2e	---	---	---
Pa: Pond Creek-----	1	---	3.00	---
Pd: Pond Creek-----	2e	---	3.00	---
Ph: Dale-----	1	---	4.00	---
Pk: Buttermilk-----	3s	---	---	---
Ps: Pratt-----	4e	3e	---	5.50
Pt: Pratt-----	6e	---	---	---
Tivoli-----	7e	---	---	---
Qn: Quinlan-----	3e	---	---	---
Qw: Quinlan-----	6e	---	---	---
Woodward-----	6e	---	---	---
Rb: Knoco-----	7s	---	---	---
Rock Outcrop-----	8s	---	---	---
Sb: Shellabarger-----	3e	---	2.00	6.00
SBB: Shellabarger-----	2e	---	2.50	7.00
Sc: Shellabarger-----	3e	---	1.50	5.50
Tv: Tivoli-----	7e	---	---	---

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Vn: Vernon-----	4e	---	---	---
Vr: Vernon-----	6e	---	---	---
Vs: Knoco-----	7s	---	---	---
Vernon-----	4e	---	---	---
W: Water-----	---	---	---	---
Wa: Waldeck-----	3w	---	3.50	5.00
Wo: Woodward-----	2e	---	---	---
Quinlan-----	3e	---	---	---
Ws: Woodward-----	3e	---	---	---
Quinlan-----	4e	---	---	---
Ya: Yahola-----	2w	---	3.50	---
Ze: Zenda-----	2w	---	4.00	5.50

CONSERVATION TREE AND SHRUB MANAGEMENT
Barber County, Kansas

A Conservation Tree/Shrub Suitability Group (CTSG), formerly Windbreak Suitability Group, is a physiographic unit or area having similar climatic and edaphic characteristics that control the selection and height growth of trees and shrubs.

In this table, the Conservation Tree and Shrub Grouping is expressed as a group index number. The group index for Conservation Tree and Shrub groups (CTSG) are a guide for species best suited for different kinds of soil and for prediction height, growth, and effectiveness. The groupings can be used when selection woody plants for windbreaks, wildlife plantings riparian buffers, reforestation, other environmental plantings, recreation, landscaping, wetland restoration or enhancement and critical area plantings. CTSG's are developed to assure satisfactory species selection and adaptation to specific conditions of soil, climate and physiography. CTSG's are a guide for selection species best suited for different kinds of soil and prediction height growth and effectiveness.

All soil series mapped in the state have been placed in 10 groups of similar soil characteristics. Groups 1, 2, 3, 4, 6, and 9 are further divided into subgroups. In addition, all groups provide information by Major Land Resource Areas.

Each tree or shrub species has certain climatic and physiographic limits. Within these parameters a tree or shrub may be well or poorly suited because of soil characteristics. Each tree or shrub also has definable potentials of height growth depending on the factors just mentioned. Accurate definitions of potential heights are necessary for proper windbreak planning and design.

Windbreaks protect livestock, buildings, roads and yards from wind and snow. They also protect fruit trees and gardens, and they furnish habitat for wildlife. Several rows of low-growing and high-growing broadleaf and coniferous trees and shrubs provide the most protection.

Field windbreaks are narrow plantings made at right angles to the prevailing wind and at specific intervals across the field. The interval depends on the erodibility of the soil. Field windbreaks protect cropland and crops from wind, help to keep snow on the fields, and provide food and cover for wildlife.

Environmental plantings help to beautify and screen houses and other buildings and to abate noise. The plants, mostly evergreen shrubs and trees, are closely spaced. To ensure plant survival, a healthy planting stock of suitable species should be planted properly on a well prepared site and maintained in good condition.

Windbreaks are often planted on land that did not grow trees originally. Knowledge of how trees perform on such land can be gained only by observing and recording their performance where trees have been planted and survived. The problem is compounded by the fact that many favorite windbreak species are not indigenous to the areas in which they are planted.

The Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups shows the adapted species listing for each group index number. Showing the height that locally grown trees and shrubs are expected to reach in 20 years on various soils. The estimates are based on measurements and observation of established plantings that have been given adequate care. This information should be used to determine the placement of a windbreak, the area protected and the arrangement of species.

A number of attributes are included in the CTSG species tables for each group number found in this section of the Field Office Technical Guide. These attributes were rated subjectively and assigned a relative value to further assist those unfamiliar with individual species characteristics or desirability for the intended use. Definitions and explanations can be found. Additional information on planning windbreaks and screens and planting and caring for trees and shrubs can be obtained from the local office of the Natural Resources Conservation Service or of the Cooperative Extension Service or from a commercial nursery. See part 537 of the National Forestry Manual for additional information.

In the Tree and Shrub Management table interpretive ratings are given for various aspects of forest and conservation tree and shrub management. Some rating class terms indicate the degree to which the soils are suited to a specified forest management practice. Well suited indicates that the soil has features that are favorable for the specified practice and has no limitations. Good performance can be expected, and little or no maintenance is needed. Moderately well suited indicates that the soil has features that are moderately favorable for the specified practice. One or more soil properties are less than desirable and fair performance can be expected. Some maintenance is needed. Poorly suited indicates that the soil has one or more properties that are unfavorable for the specified practice. Overcoming the unfavorable properties requires special design, extra maintenance, and costly alteration. Unsited indicates that the expected performance of the soil is unacceptable for the specified practice or that extreme measures are needed to overcome the undesirable soil properties.

The paragraphs that follow indicate the soil properties considered in rating the soils for forest and conservation tree and shrub management practices. More detailed information about the criteria used in the ratings is available in the "National Forestry Manual," which is available in local offices of the Natural Resources Conservation Service or on the Internet. Also, in the Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups.

Ratings in the columns suitability for hand planting and suitability for mechanical planting are based on slope, depth to a restrictive layer, content of sand, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, moderately well suited, poorly suited, or unsited to these methods of planting. It is assumed that necessary site preparation is completed before seedlings are planted.

Ratings in the column suitability for mechanical site preparation (surface) are based on slope, depth to a restrictive layer, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsited to this management activity. The part of the soil from the surface to a depth of about 1-foot is considered in the ratings.

Ratings in the column suitability for mechanical site preparation (deep) are based on slope, depth to a restrictive layer, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsited to this management activity. The part of the soil from the surface to a depth of about 3 feet is considered in the ratings.

Ratings in the column potential for seedling mortality are based on flooding, ponding, depth to a water table, content of lime, reaction, salinity, available water capacity, soil moisture regime, soil temperature regime, aspect, and slope. The soils are described as having a low, moderate, or high potential for seedling mortality. See the National Forestry Manual, Subpart B for criteria used in rating management concerns. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

CONSERVATION TREE AND SHRUB MANAGEMENT
Barber County,
Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
033CP: Clairemont-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
033OA: Obaro-----		Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Soil reaction
033OB: Obaro-----		Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Soil reaction
Rock Outcrop-----		Not rated	Not rated	Not rated	Not rated	Not rated
033QR: Quinlan-----	10	Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Soil reaction
Woodward-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Low
033QT: Quinlan-----	10	Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Soil reaction
Woodward-----	8	Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Low
033YE: Yahola-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
077BP: Woodward-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Port-----	1	Well suited	Well suited	Well suited	Well suited	Low
077FU: Farnum-----	3	Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
077GE: Gerlane-----	1	Well suited	Well suited	Well suited	Well suited	Low
077MC: Minco-----	3	Well suited	Well suited	Well suited	Well suited	Low
077MN: Minco-----	3	Well suited	Well suited	Well suited	Well suited	Low
077MO: Minco-----	3	Well suited	Moderately suited Slope	Well suited	Well suited	Low
077QU: Quinlan-----	10	Moderately suited Rock fragments Restrictive layer	Poorly suited Rock fragments Slope	Poorly suited Rock fragments	Well suited	Moderate Soil reaction
077SE: Shellabarger-----	5	Well suited	Well suited	Well suited	Well suited	Low
077WE: Quinlan-----	10	Moderately suited Rock fragments Restrictive layer	Poorly suited Rock fragments	Poorly suited Rock fragments	Well suited	Moderate Soil reaction
Woodward-----	8	Well suited	Well suited	Well suited	Well suited	Low
095AC: Albion-----	6G	Well suited	Moderately suited Slope	Well suited	Well suited	Low
095DA: Dillwyn-----	1	Well suited	Well suited	Well suited	Well suited	Low
Plevna-----	2	Well suited	Well suited	Well suited	Unsuited Wetness	High Wetness
095FA: Farnum-----	3	Well suited	Well suited	Well suited	Well suited	Low
095PB: Pratt-----	7	Well suited	Moderately suited Slope	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT
Barber County,
Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
095SB: Shellabarger-----	5	Well suited	Well suited	Well suited	Well suited	Low
095WA: Waldeck-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
097LN: Lincoln-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
097QW: Quinlan-----	10	Well suited	Poorly suited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Soil reaction
Woodward-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Low
097WA: Waldeck-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
151AB: Albion-----	6G	Well suited	Well suited	Well suited	Well suited	Low
151AO: Albion-----	6G	Well suited	Moderately suited Slope	Well suited	Well suited	Low
151AS: Albion-----	6G	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Shellabarger-----	5	Well suited	Moderately suited Slope	Well suited	Well suited	Low
151BC: Blanket-----	4C	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
151CA: Carwile-----	1	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	High Wetness
151CK: Case-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Soil reaction
Clark-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Soil reaction
151CM: Clark-----	8	Well suited	Well suited	Well suited	Well suited	Lime Soil reaction
151CN: Clark-----	8	Well suited	Well suited	Well suited	Well suited	Moderate Lime Soil reaction
151CO: Clark-----	8	Well suited	Well suited	Well suited	Well suited	Moderate Lime Soil reaction
Ost-----	3	Well suited	Well suited	Well suited	Well suited	Moderate Lime Soil reaction
151FA: Farnum-----	3	Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
151KP: Kanza-----	2	Well suited	Well suited	Well suited	Well suited	Low
Plevna-----	2	Well suited	Well suited	Well suited	Unsuited Wetness	High Wetness
151OS: Ost-----	3	Well suited	Well suited	Well suited	Well suited	Low
151PM: Pratt-----	7	Well suited	Moderately suited Slope	Well suited	Well suited	Low
151PN: Pratt-----	7	Well suited	Moderately suited Slope	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT
Barber County,
Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
151PO: Pratt-----	7	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Carwile-----	1	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	High Wetness
151SA: Albion-----	6G	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Kaski-----	1K	Well suited	Well suited	Well suited	Well suited	Low
151SE: Shellabarger-----	5	Well suited	Well suited	Well suited	Well suited	Low
1439: Crisfield-----	5	Well suited	Well suited	Well suited	Well suited	Low
Ad: Albion-----	6G	Well suited	Well suited	Well suited	Well suited	Low
Shellabarger-----	5	Well suited	Well suited	Well suited	Well suited	Low
Ae: Albion-----	6G	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Shellabarger-----	5	Well suited	Moderately suited Slope	Well suited	Well suited	Low
AED: Arents, Earthen Dam-		Not rated	Not rated	Not rated	Not rated	Not rated
AN: Albion-----	6G	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Shellabarger-----	5	Well suited	Moderately suited Slope	Well suited	Well suited	Low
As: Clairemont-----	1K	Well suited	Well suited	Well suited	Well suited	High Salinity Soil reaction
At: Attica-----	5	Well suited	Well suited	Well suited	Well suited	Low
Ba: Blanket-----	4C	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Bb: Blanket-----	4C	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Bc: Blanket-----	4C	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Bf: Quinlan-----	10	Well suited	Unsuited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Soil reaction
Clairemont-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Ca: Canadian-----	1	Well suited	Well suited	Well suited	Well suited	Low
Cc: Case-----	8	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Clark-----	8	Well suited	Well suited	Well suited	Well suited	Moderate Lime Soil reaction
Cd: Clairemont-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Cf: Clairemont-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Ck: Clark-----	8	Well suited	Well suited	Well suited	Well suited	Moderate Lime Soil reaction
Fa: Farnum-----	3	Moderately suited	Moderately suited	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT
Barber County,
Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Fm: Farnum-----	3	Stickiness	Stickiness			
Fr: Farnum-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Fu: Farnum-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Ga: Grant-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Gb: Grant-----	3	Well suited	Well suited	Well suited	Well suited	Low
Gc: Grant-----	3	Well suited	Well suited	Well suited	Well suited	Low
GRP: Gravel Pits-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
INT: Aquolls-----		Not rated	Not rated	Not rated	Not rated	Not rated
Ka: Kanza-----	2	Moderately suited Rock fragments	Poorly suited	Poorly suited	Well suited	High
Kf: Kingfisher-----	6D	Well suited	Rock fragments	Rock fragments		Wetness
Kv: Kingfisher-----	6D	Well suited	Well suited	Well suited		Soil reaction
Vernon-----	6D	Well suited	Well suited	Well suited	Well suited	Low
Kz: Kingfisher-----	6D	Moderately suited Stickiness	Well suited	Poorly suited	Well suited	Low
Vernon-----		Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Ln: Lincoln-----	1K	Well suited	Well suited	Well suited	Well suited	Low
LNN: Lincoln-----	1K	Well suited	Well suited	Well suited	Well suited	Low
Ma: Mangum-----	1K	Well suited	Well suited	Well suited	Well suited	Low
Mg: Mangum-----	1K	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Moderate Soil reaction
Drummond-----	9W	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Moderate Soil reaction
Mn: Minco-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Moderate
Na: Naron-----	5	Well suited	Well suited	Well suited	Well suited	Salinity
Nb: Naron-----	5	Well suited	Well suited	Well suited	Well suited	Low
Os: Ost-----	3	Well suited	Well suited	Well suited	Well suited	Low
Ot: Ost-----	3	Well suited	Well suited	Well suited	Well suited	Low
Pa: Pond Creek-----	3	Well suited	Well suited	Well suited	Well suited	Low
Pd: Pond Creek-----	3	Well suited	Well suited	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT
Barber County,
Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Ph: Dale-----	1	Well suited	Well suited	Well suited	Well suited	Low
Pk: Buttermilk-----	9L	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction Salinity
Ps: Pratt-----	7	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Pt: Pratt-----	7	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Tivoli-----	7	Moderately suited Sandiness	Moderately suited Slope Sandiness	Well suited	Well suited	Low
Qn: Quinlan-----	10	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Qw: Quinlan-----	10	Well suited	Moderately suited Slope	Well suited	Well suited	Moderate
Woodward-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Soil reaction Low
Rb: Knoco-----		Moderately suited Restrictive layer	Unsuited Slope	Poorly suited Slope	Poorly suited Slope	Moderate Lime
Rock Outcrop-----		Stickiness Not rated	Stickiness Not rated	Not rated	Not rated	Soil reaction Not rated
Sb: Shellabarger-----	5	Well suited	Moderately suited Slope	Well suited	Well suited	Low
SBB: Shellabarger-----	5	Well suited	Well suited	Well suited	Well suited	Low
Sc: Shellabarger-----	5	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Tv: Tivoli-----	7	Moderately suited Sandiness	Moderately suited Slope Sandiness	Well suited	Well suited	Low
Vn: Vernon-----	4C	Moderately suited Stickiness	Moderately suited Stickiness Rock fragments	Poorly suited Stickiness	Well suited	Low
Vr: Vernon-----	4C	Moderately suited Stickiness	Moderately suited Stickiness Slope Rock fragments	Poorly suited Stickiness	Well suited	Low
Vs: Knoco-----		Moderately suited Restrictive layer	Moderately suited Stickiness	Well suited	Well suited	Moderate Lime
Vernon-----	4C	Stickiness Moderately suited Stickiness	Moderately suited Stickiness Rock fragments	Poorly suited Stickiness	Well suited	Soil reaction Low
W: Water-----		Not rated	Not rated	Not rated	Not rated	Not rated

CONSERVATION TREE AND SHRUB MANAGEMENT
Barber County,
Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Wa: Waldeck-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Wo: Woodward-----	8	Well suited	Well suited	Well suited	Well suited	Low
Quinlan-----	10	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Ws: Woodward-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Quinlan-----	10	Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Soil reaction
Ya: Yahola-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Ze: Zenda-----	1	Well suited	Well suited	Well suited	Well suited	Low

ENGINEERING INDEX PROPERTIES
Barber County, Kansas

Engineering Index Properties table gives the engineering classifications and the range of index properties for the layers of each soil in the survey area. Depth to the upper and lower boundaries of each layer is indicated. Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter. Loam, for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, gravelly. Textural terms are defined in the Glossary.

Classification of the soils is determined according to the Unified soil classification system (ASTM, 1998) and the system adopted by the American Association of State Highway and Transportation Officials (AASHTO, 1998). The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection. If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional refinement, the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest. The AASHTO classification for soils tested, with group index numbers in parentheses, is given in Engineering Index Properties table.

Rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in the field to weight percentage. Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field.

Liquid limit and plasticity index (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination. The estimates of particle-size distribution, liquid limit, and plasticity index are generally rounded to the nearest 5 percent. Thus, if the ranges of gradation and Atterberg limits extend a marginal amount (1 or 2 percentage points) across classification boundaries, the classification in the marginal zone is generally omitted in the table.

ENGINEERING INDEX PROPERTIES--Continued
Barber County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
033CP: Clairemont-----	0-15	Loam	CL, CL-ML	A-4, A-6	0	0	100	98-100	85-100	60-90	20-35	4-17
	15-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	98-100	95-100	65-95	20-40	4-20
033OA: Obaro-----	0-35	Silty clay loam	CL, CL-ML	A-4, A-6	0	0	98-100	95-100	95-100	80-98	25-40	7-20
	>35	Weathered bedrock			---	---					---	---
033OB: Obaro-----	0-35	Silty clay loam	CL, CL-ML	A-4, A-6	0	0	98-100	95-100	95-100	80-98	25-40	7-20
	>35	Weathered bedrock			---	---					---	---
Rock Outcrop---	---	---	---	---	---	---	---	---	---	---	---	---
033QR: Quinlan-----	0-14	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	90-100	51-97	15-37	NP-14
	>14	Weathered bedrock			---	---					---	---
Woodward-----	0-30	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	51-95	15-31	NP-12
	>30	Weathered bedrock			---	---					---	---
033QT: Quinlan-----	0-14	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	90-100	51-97	15-37	NP-14
	>14	Weathered bedrock			---	---					---	---
Woodward-----	0-30	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	51-95	15-31	NP-12
	>30	Weathered bedrock			---	---					---	---
033YE: Yahola-----	0-12	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	95-100	90-100	36-60	15-26	NP-7
	12-40	Fine sandy loam	CL, ML, SC, SM	A-4	0	0	100	95-100	90-100	36-85	15-30	NP-10
	40-60	Stratified loam to loamy fine sand	CL, ML, SC, SM	A-2, A-4	0	0	100	95-100	90-100	15-85	15-30	NP-10
077BP: Woodward-----	0-24	Silt loam	CL, CL-ML, ML	A-6, A-4	0	0	100	100	90-100	51-95	15-31	NP-12
	>24	Weathered bedrock			---	---					---	---
Port-----	0-27	Silt loam	CL	A-4, A-6	0	0	100	100	96-100	65-97	27-37	8-14
	27-60	Silty clay loam	CL	A-4, A-6, A-7	0	0	100	100	96-100	65-98	27-43	8-20
077FU: Farnum-----	0-11	Loam	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	60-85	20-35	5-15
	11-41	Clay loam	CL, SC	A-6, A-7-6	0	0	100	100	70-100	45-80	35-50	15-30
	41-60	Clay loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6	0	0	100	95-100	65-100	30-80	20-35	5-15
077GE: Gerlane-----	0-17	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	95-100	70-95	40-55	15-26	NP-6
	17-40	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	95-100	70-95	35-55	15-26	NP-6
	40-48	Loamy sand	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	95-100	70-95	15-55	15-26	NP-6
	48-60	Clay loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	65-95	20-40	5-25
077MC: Minco-----	0-42	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	94-100	51-97	15-31	NP-10
	42-60	Silt loam	CL, ML, SC, SM	A-4	0	0	100	98-100	94-100	36-97	15-31	NP-10
077MN: Minco-----	0-42	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	94-100	51-97	15-31	NP-10
	42-60	Silt loam	CL, ML, SC, SM	A-4	0	0	100	98-100	94-100	36-97	15-31	NP-10
077MO: Minco-----	0-42	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	94-100	51-97	15-31	NP-10
	42-60	Silt loam	CL, ML, SC, SM	A-4	0	0	100	98-100	94-100	36-97	15-31	NP-10
077QU: Quinlan-----	0-9	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	90-100	51-97	15-37	NP-14
	>9	Weathered bedrock			---	---					---	---
077SE: Shellabarger---	0-13	Fine sandy loam	ML, SM	A-2, A-4	0	0	95-100	95-100	75-100	30-55	15-30	NP-5
	13-38	Fine sandy loam	SC	A-4, A-6	0	0	95-100	85-100	70-90	35-50	25-40	8-20
	38-60	Coarse sandy loam	SC, SC-SM, SM, SP-SM	A-2, A-4	0	0	80-100	70-100	50-80	10-40	15-30	NP-10
077WE: Woodward-----	0-24	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	51-95	15-31	NP-12
	>24	Weathered bedrock			---	---					---	---
Quinlan-----	0-9	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	90-100	51-97	15-37	NP-14
	>9	Weathered bedrock			---	---					---	---
095AC: Albion-----	0-8	Sandy loam	ML, SM	A-2, A-4	0	0	100	75-100	60-90	25-55	15-30	NP-5
	8-16	Sandy loam	ML, SM	A-2, A-4	0	0	85-100	75-100	45-90	30-55	20-35	NP-10
	16-26	Loamy sand	SM	A-1, A-2	0	0	85-100	75-90	40-70	15-30	15-30	NP-5
	26-60	Sand	GM, GP-GM, SM, SP-SM	A-1, A-2, A-3	0	0-5	40-100	40-90	30-70	5-30	15-30	NP-5

ENGINEERING INDEX PROPERTIES--Continued
Barber County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
095DA: Dillwyn-----	0-8	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	70-90	5-35	---	NP
	8-60	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	90-100	70-90	5-35	---	NP
Plevna-----	0-11	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	20-50	15-26	NP-6
	11-36	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	30-50	15-26	NP-6
	36-60	Sand	SM, SP	A-2, A-3	0	0	100	90-100	50-90	4-35	---	NP
095FA: Farnum-----	0-16	Sandy loam	ML, SM	A-2, A-4	0	0	100	100	70-100	30-55	15-30	NP-5
	16-50	Clay loam	CL, SC	A-6, A-7-6	0	0	100	100	70-100	45-80	35-50	15-30
	50-60	Clay loam	SC-SM, CL, CL-ML, SC	A-2, A-4, A-6	0	0	100	95-100	65-100	30-80	20-35	5-15
095PB: Pratt-----	0-12	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	---	NP
	12-36	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	36-60	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-35	---	NP
095SB: Shellabarger---	0-10	Sandy loam	ML, SM, CL-ML	A-2, A-4	0	0	95-100	95-100	75-100	30-55	15-30	NP-5
	10-45	Sandy clay loam	SC	A-4, A-6	0	0	95-100	85-100	70-90	35-50	25-40	8-20
	45-60	Coarse sandy loam	SC, SC-SM, SM, SP-SM	A-2, A-4	0	0	80-100	70-100	50-80	10-40	15-30	NP-10
095WA: Waldeck-----	0-12	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	95-100	75-100	25-55	15-25	NP-5
	12-36	Sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	30-50	15-25	NP-5
	36-60	Sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	40-60	1-35	---	NP
097LN: Lincoln-----	0-8	Sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	98-100	94-100	36-60	15-24	NP-7
	8-60	Stratified fine sand to loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	98-100	82-100	5-35	---	NP
097QW: Quinlan-----	0-15	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	90-100	51-97	15-37	NP-14
	>15	Weathered bedrock			---	---	---	---	---	---	---	---
Woodward-----	0-27	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	51-95	15-31	NP-12
	>27	Weathered bedrock			---	---	---	---	---	---	---	---
097WA: Waldeck-----	0-14	Loam	ML, CL-ML	A-4	0	0	100	95-100	75-100	50-60	15-30	NP-7
	14-41	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	30-50	15-25	NP-5
	41-60	Fine sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	40-60	1-35	---	NP
151AB: Albion-----	0-8	Sandy loam	ML, SM, CL-ML	A-2, A-4	0	0	100	75-100	60-90	25-55	15-30	NP-5
	8-18	Sandy loam	ML, SM	A-2, A-4	0	0	85-100	75-100	45-90	30-55	20-35	NP-10
	18-29	Coarse sandy loam	SM	A-1, A-2	0	0	85-100	75-90	40-70	15-30	15-30	NP-5
	29-60	Gravelly sand	GM, GP-GM, SM, SP-SM	A-1, A-2, A-3	0	0-5	40-100	40-90	30-70	5-30	15-30	NP-5
151AO: Albion-----	0-8	Sandy loam	ML, SM, CL-ML	A-2, A-4	0	0	100	75-100	60-90	25-55	15-30	NP-5
	8-18	Sandy loam	ML, SM	A-2, A-4	0	0	85-100	75-100	45-90	30-55	20-35	NP-10
	18-29	Coarse sandy loam	SM	A-1, A-2	0	0	85-100	75-90	40-70	15-30	15-30	NP-5
	29-60	Gravelly sand	SM, SP-SM, GM, GP-GM	A-1, A-2, A-3	0	0-5	40-100	40-90	30-70	5-30	15-30	NP-5
151AS: Albion-----	0-8	Sandy loam	ML, SM	A-2, A-4	0	0	100	75-100	60-90	25-55	15-30	NP-5
	8-18	Sandy loam	ML, SM	A-2, A-4	0	0	85-100	75-100	45-90	30-55	20-35	NP-10
	18-29	Coarse sandy loam	SM	A-1, A-2	0	0	85-100	75-90	40-70	15-30	15-30	NP-5
	29-60	Gravelly sand	GM, GP-GM, SM, SP-SM	A-1, A-2, A-3	0	0-5	40-100	40-90	30-70	5-30	15-30	NP-5
Shellabarger---	0-11	Fine sandy loam	ML, SM	A-2, A-4	0	0	95-100	95-100	75-100	30-55	15-30	NP-5
	11-34	Sandy clay loam	SC	A-4, A-6	0	0	95-100	85-100	70-90	35-50	25-40	8-20
	34-60	Coarse sandy loam	SC, SC-SM, SM, SP-SM	A-2, A-4	0	0	80-100	70-100	50-80	10-40	15-30	NP-10
151BC: Blanket-----	0-13	Silty clay loam	CH, CL	A-6, A-7	0	0	98-100	96-100	90-100	70-98	36-62	16-38
	13-46	Silty clay	CH, CL	A-7	0	0	98-100	96-100	85-100	70-90	41-64	20-38
	46-60	Silty clay loam	CH, CL	A-6, A-7	0	0	85-100	80-100	80-100	51-85	30-60	15-38
151CA: Carwile-----	0-12	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	98-100	90-100	36-60	15-26	NP-7
	12-20	Sandy clay loam	CL, SC	A-6, A-7	0	0	100	100	90-100	36-90	35-50	14-26
	20-33	Clay	CH, CL, SC	A-6, A-7	0	0	100	100	90-100	40-95	35-70	14-38
	33-60	Sandy clay loam	CH, CL, SC	A-4, A-6, A-7	0	0	100	100	90-100	36-95	25-70	7-38
151CK: Case-----	0-6	Clay loam	CL	A-6	0	0	90-100	90-100	85-100	55-85	30-40	10-20
	6-60	Clay loam	CL	A-6, A-7-6	0	0	90-100	90-100	85-100	55-85	25-45	10-25
Clark-----	0-8	Clay loam	CL	A-6	0	0	100	95-100	90-100	50-90	30-40	10-20
	8-60	Clay loam	CL	A-6	0	0	100	95-100	90-100	55-90	25-40	10-25
151CM: Clark-----	0-8	Clay loam	CL	A-6	0	0	100	95-100	90-100	50-90	30-40	10-20
	8-60	Clay loam	CL	A-6	0	0	100	95-100	90-100	55-90	25-40	10-25
151CN: Clark-----	0-8	Fine sandy loam	SC, SC-SM	A-4	0	0	100	95-100	90-100	35-50	15-25	5-10
	8-60	Clay loam	CL	A-6	0	0	100	95-100	90-100	55-90	25-40	10-25

ENGINEERING INDEX PROPERTIES--Continued
Barber County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
151CO: Clark-----	0-8	Clay loam	CL	A-6	0	0	100	95-100	90-100	50-90	30-40	10-20
	8-60	Clay loam	CL	A-6	0	0	100	95-100	90-100	55-90	25-40	10-25
Ost-----	0-9	Clay loam	CL	A-6	0	0	95-100	95-100	85-100	75-90	30-40	10-20
	9-14	Clay loam	CL	A-4, A-6	0	0	95-100	90-100	85-100	55-90	30-40	9-18
	14-23	Clay loam	CL, SC	A-2, A-4, A-6	0	0	95-100	90-100	80-100	30-90	25-40	8-18
	23-60	Clay loam	CL, ML, SC, SM	A-2, A-4, A-6	0	0	85-100	85-100	80-100	30-90	15-40	NP-18
151FA: Farnum-----	0-7	Clay loam	CL	A-6	0	0	100	100	90-100	60-85	30-40	10-20
	7-37	Clay loam	CL, SC	A-6, A-7-6	0	0	100	100	70-100	45-80	35-50	15-30
	37-60	Fine sandy loam	SC-SM, CL, CL-ML, SC	A-2, A-4, A-6	0	0	100	95-100	65-100	30-80	20-35	5-15
151KP: Kanza-----	0-11	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	90-100	5-35	---	NP
	11-40	Fine sand	SM, SP-SM	A-2, A-3	0	0	90-100	90-100	80-100	5-35	---	NP
Plevna-----	0-10	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	20-50	15-26	NP-6
	10-40	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	30-50	15-26	NP-6
	40-60	Fine sand	SM, SP	A-2, A-3	0	0	100	90-100	50-90	4-35	---	NP
151OS: Ost-----	0-9	Clay loam	CL	A-6	0	0	95-100	95-100	85-100	75-90	30-40	10-20
	9-14	Clay loam	CL	A-4, A-6	0	0	95-100	90-100	85-100	55-90	30-40	9-18
	14-23	Clay loam	CL, SC	A-2, A-4, A-6	0	0	95-100	90-100	80-100	30-90	25-40	8-18
	23-60	Clay loam	CL, ML, SC, SM	A-2, A-4, A-6	0	0	85-100	85-100	80-100	30-90	15-40	NP-18
151PM: Pratt-----	0-10	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	---	NP
	10-40	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	40-60	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-35	---	NP
151PN: Pratt-----	0-10	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	---	NP
	10-40	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	40-60	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-35	---	NP
151PO: Pratt-----	0-10	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	---	NP
	10-40	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	40-60	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-35	---	NP
Carwile-----	0-12	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	98-100	90-100	36-60	15-26	NP-7
	12-20	Sandy clay loam	CL, SC	A-6, A-7	0	0	100	100	90-100	36-90	35-50	14-26
	20-33	Clay	CH, CL, SC	A-6, A-7	0	0	100	100	90-100	40-95	35-70	14-38
	33-60	Sandy clay loam	CH, CL, SC	A-4, A-6, A-7	0	0	100	100	90-100	36-95	25-70	7-38
151SA: Albion-----	0-8	Fine sandy loam	ML, SM, CL-ML	A-2, A-4	0	0	100	75-100	60-90	25-55	15-30	NP-5
	8-18	Sandy loam	ML, SM, CL-ML	A-2, A-4	0	0	85-100	75-100	45-90	30-55	20-35	NP-10
	18-29	Coarse sandy loam	SM	A-1, A-2	0	0	85-100	75-90	40-70	15-30	15-30	NP-5
	29-60	Gravelly sand	GM, GP-GM, SM, SP-SM	A-1, A-2, A-3	0	0-5	40-100	40-90	30-70	5-30	15-30	NP-5
Kaski-----	0-26	Loam	CL, CL-ML	A-4, A-6, A-7	0	0	100	100	85-100	50-85	20-45	5-25
	26-40	Loam	CL, SC	A-4, A-6, A-7	0	0	100	95-100	85-100	45-85	25-45	7-25
	40-60	Sandy loam	CL, ML, SC, SM	A-2, A-4, A-6	0	0	100	95-100	60-100	30-80	15-35	NP-20
151SE: Shellabarger---	0-11	Fine sandy loam	ML, SM, CL-ML	A-2, A-4	0	0	95-100	95-100	75-100	30-55	15-30	NP-5
	11-34	Sandy clay loam	SC	A-4, A-6	0	0	95-100	85-100	70-90	35-50	25-40	8-20
	34-60	Coarse sandy loam	SC, SC-SM, SM, SP-SM	A-2, A-4	0	0	80-100	70-100	50-80	10-40	15-30	NP-10
1439: Crisfield-----	0-12	Sandy loam	SC, SM	A-6, A-2-6, A-4	0	0	95-100	95-100	65-75	30-45	10-20	NP-12
	12-24	Sandy loam	SC, SM	A-6, A-2-6, A-4	0	0	100	95-100	70-75	30-60	10-20	NP-12
	24-80	Coarse sand	SP-SM	A-3, A-1-b, A-2-4	0	0	95-100	90-100	45-55	5-40	0-5	NP-2
Ad: Albion-----	0-8	Sandy loam	ML, SM	A-2, A-4	0	0	100	75-100	60-90	25-55	15-30	NP-5
	8-16	Sandy loam	ML, SM	A-2, A-4	0	0	85-100	75-100	45-90	30-55	20-35	NP-10
	16-27	Loamy sand	SM	A-1, A-2	0	0	85-100	75-90	40-70	15-30	15-30	NP-5
	27-60	Sand	GM, GP-GM, SM, SP-SM	A-1, A-2, A-3	0	0-5	40-100	40-90	30-70	5-30	15-30	NP-5
Shellabarger---	0-14	Sandy loam	ML, SM	A-2, A-4	0	0	95-100	95-100	75-100	30-55	15-30	NP-5
	14-48	Sandy clay loam	SC	A-4, A-6	0	0	95-100	85-100	70-90	35-50	25-40	8-20
	48-60	Coarse sandy loam	SC, SC-SM, SM, SP-SM	A-2, A-4	0	0	80-100	70-100	50-80	10-40	15-30	NP-10
Ae: Albion-----	0-8	Sandy loam	ML, SM, CL-ML	A-2, A-4	0	0	100	75-100	60-90	25-55	15-30	NP-5
	8-16	Sandy loam	ML, SM, CL-ML	A-2, A-4	0	0	85-100	75-100	45-90	30-55	20-35	NP-10
	16-27	Loamy sand	SM	A-1, A-2	0	0	85-100	75-90	40-70	15-30	15-30	NP-5
	27-60	Sand	GM, GP-GM, SM, SP-SM	A-1, A-2, A-3	0	0-5	40-100	40-90	30-70	5-30	15-30	NP-5
Shellabarger---	0-14	Sandy loam	ML, SM	A-2, A-4	0	0	95-100	95-100	75-100	30-55	15-30	NP-5
	14-48	Sandy clay loam	SC	A-4, A-6	0	0	95-100	85-100	70-90	35-50	25-40	8-20
	48-60	Coarse sandy loam	SC, SC-SM, SM, SP-SM	A-2, A-4	0	0	80-100	70-100	50-80	10-40	15-30	NP-10
AED: Arents, Earthen Dam-----	---	---	---	---	---	---	---	---	---	---	---	---

ENGINEERING INDEX PROPERTIES--Continued
Barber County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
AN:												
Albion-----	0-8	Sandy loam	CL-ML, ML, SM	A-2, A-4	0	0	100	75-100	60-90	25-55	15-30	NP-5
	8-16	Sandy loam	CL-ML, ML, SM	A-2, A-4	0	0	85-100	75-100	45-90	30-55	20-35	NP-10
	16-28	Loamy sand	SM	A-1, A-2	0	0	85-100	75-90	40-70	15-30	15-30	NP-5
	28-60	Sand	GM, GP-GM, SM, SP-SM	A-1, A-2, A-3	0	0-5	40-100	40-90	30-70	5-30	15-30	NP-5
Shellabarger---	0-11	Sandy loam	CL-ML, ML, SM	A-2, A-4	0	0	95-100	95-100	75-100	30-55	15-30	NP-5
	11-38	Sandy clay loam	SC	A-4, A-6	0	0	95-100	85-100	70-90	35-50	25-40	8-20
	38-60	Coarse sandy loam	SC, SC-SM, SM, SP-SM	A-2, A-4	0	0	80-100	70-100	50-80	10-40	15-30	NP-10
As:												
Clairemont----	0-8	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	98-100	85-100	65-95	22-40	5-20
	8-60	Loam	CL, CL-ML	A-4, A-6, A-7-6	0	0	100	98-100	90-100	65-95	22-45	5-25
At:												
Attica-----	0-12	Loamy fine sand	SM, SP-SM	A-2	0	0	100	95-100	70-100	10-35	---	NP
	12-30	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	95-100	75-100	30-55	15-26	NP-7
	30-60	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	85-100	80-100	70-100	20-50	15-26	NP-7
Ba:												
Blanket-----	0-9	Silt loam	CL	A-6	0	0	98-100	96-100	85-100	65-95	28-40	11-20
	9-30	Silty clay	CH, CL	A-7	0	0	98-100	96-100	85-100	70-90	41-64	20-38
	30-60	Silty clay loam	CH, CL	A-6, A-7	0	0	85-100	80-100	80-100	51-85	30-60	15-38
Bb:												
Blanket-----	0-9	Silt loam	CL	A-6	0	0	98-100	96-100	85-100	65-95	28-40	11-20
	9-30	Silty clay	CH, CL	A-7	0	0	98-100	96-100	85-100	70-90	41-64	20-38
	30-60	Silty clay loam	CL, CH	A-6, A-7	0	0	85-100	80-100	80-100	51-85	30-60	15-38
Bc:												
Blanket-----	0-9	Silty clay loam	CH, CL	A-7, A-6	0	0	98-100	96-100	90-100	70-98	36-62	16-38
	9-30	Silty clay	CH, CL	A-7	0	0	98-100	96-100	85-100	70-90	41-64	20-38
	30-60	Silty clay loam	CH, CL	A-6, A-7	0	0	85-100	80-100	80-100	51-85	30-60	15-38
Bf:												
Quinlan-----	0-14	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	90-100	51-97	15-37	NP-14
	>14				---	---	---	---	---	---	---	---
Clairemont----	0-14	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	98-100	85-100	60-90	20-35	4-17
	14-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	98-100	95-100	65-95	20-40	4-20
Ca:												
Canadian-----	0-18	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	98-100	94-100	36-65	15-26	NP-7
	18-48	Fine sandy loam	CL, ML, SC, SM	A-4	0	0	100	98-100	94-100	36-85	15-31	NP-10
	48-60	Fine sandy loam	CL, ML, SC, SM	A-2, A-4	0	0	100	98-100	90-100	15-85	15-31	NP-10
Cc:												
Case-----	0-6	Clay loam	CL	A-6	0	0	90-100	90-100	85-100	55-85	30-40	10-20
	6-60	Clay loam	CL	A-6, A-7-6	0	0	90-100	90-100	85-100	55-85	25-45	10-25
Clark-----	0-10	Clay loam	CL	A-6	0	0	100	95-100	90-100	50-90	30-40	10-20
	10-60	Clay loam	CL	A-6	0	0	100	95-100	90-100	55-90	25-40	10-25
Cd:												
Clairemont----	0-14	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	98-100	85-100	60-90	20-35	4-17
	14-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	98-100	95-100	65-95	20-40	4-20
Cf:												
Clairemont----	0-14	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	98-100	85-100	60-90	20-35	4-17
	14-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	98-100	95-100	65-95	20-40	4-20
Ck:												
Clark-----	0-10	Clay loam	CL	A-6	0	0	100	95-100	90-100	50-90	30-40	10-20
	10-60	Clay loam	CL	A-6	0	0	100	95-100	90-100	55-90	25-40	10-25
Fa:												
Farnum-----	0-9	Fine sandy loam	ML, SM	A-2, A-4	0	0	100	100	70-100	30-55	15-30	NP-3
	9-60	Clay loam	CL, SC	A-6, A-7-6	0	0	100	100	70-100	45-80	35-50	15-30
Fm:												
Farnum-----	0-9	Loam	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	60-85	20-35	5-15
	9-60	Clay loam	CL, SC	A-6, A-7-6	0	0	100	100	70-100	45-80	35-50	15-30
Fr:												
Farnum-----	0-9	Loam	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	60-85	20-35	5-15
	9-60	Clay loam	CL, SC	A-6, A-7-6	0	0	100	100	70-100	45-80	35-50	15-30
Fu:												
Farnum-----	0-9	Clay loam	CL	A-6	0	0	100	100	90-100	60-85	30-40	10-20
	9-60	Clay loam	CL, SC	A-6, A-7-6	0	0	100	100	70-100	45-80	35-50	15-30
Ga:												
Grant-----	0-13	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	90-100	70-90	20-32	1-10
	13-50	Silt loam	CL, ML	A-4, A-6, A-7	0	0	100	100	90-100	70-90	30-42	8-19
	50-60	Silt loam	CL, CL-ML, ML	A-4	0	0	70-100	70-100	65-100	55-90	20-32	1-10
Gb:												
Grant-----	0-13	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	90-100	70-90	20-32	1-10
	13-50	Silt loam	CL, ML	A-4, A-6, A-7	0	0	100	100	90-100	70-90	30-42	8-19
	50-60	Silt loam	CL, CL-ML, ML	A-4	0	0	70-100	70-100	65-100	55-90	20-32	1-10
Gc:												
Grant-----	0-13	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	90-100	70-90	20-32	1-10
	13-50	Silt loam	CL, ML	A-4, A-6, A-7	0	0	100	100	90-100	70-90	30-42	8-19
	50-60	Silt loam	CL, CL-ML, ML	A-4	0	0	70-100	70-100	65-100	55-90	20-32	1-10
GRP:												
Gravel Pits----	---	---	---	---	---	---	---	---	---	---	---	---
INT:												
Aquolls-----	0-60	Variable			---	---	---	---	---	---	---	---
Ka:												
Kanza-----	0-7	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	90-100	5-35	---	NP
	7-48	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	90-100	90-100	80-100	5-35	---	NP

ENGINEERING INDEX PROPERTIES--Continued
Barber County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
Kf: Kingfisher-----	0-9 9-17 17-28 >28	Silt loam Silty clay loam Silty clay loam Weathered bedrock	CL, CL-ML, ML CL CL CL	A-4, A-6 A-4, A-6 A-6, A-7	0 0 0 ---	0 0 0 ---	100 100 100 ---	100 100 100 ---	96-100 96-100 96-100 ---	65-97 80-98 80-98 ---	22-37 30-40 33-43 ---	3-14 8-17 12-20 ---
Kv: Kingfisher-----	0-9 9-17 17-28 >28	Silt loam Silty clay loam Silty clay loam Weathered bedrock	CL, CL-ML, ML CL CL CL	A-4, A-6 A-4, A-6 A-6, A-7	0 0 0 ---	0 0 0 ---	100 100 100 ---	100 100 100 ---	96-100 96-100 96-100 ---	65-97 80-98 80-98 ---	22-37 30-40 33-43 ---	3-14 8-17 12-20 ---
Vernon-----	0-5 5-22 22-60	Clay loam Silty clay Weathered bedrock	CL CH, CL	A-6, A-7 A-7	--- --- ---	0-15 0-15 ---	90-100 90-100 ---	85-95 70-100 ---	80-95 70-95 ---	60-95 60-90 ---	37-50 41-70 ---	14-25 20-40 ---
Kz: Kingfisher-----	0-9 9-17 17-28 >28	Silt loam Silty clay loam Silty clay loam Weathered bedrock	CL, CL-ML, ML CL CL CL	A-4, A-6 A-4, A-6 A-6, A-7	0 0 0 ---	0 0 0 ---	100 100 100 ---	100 100 100 ---	96-100 96-100 96-100 ---	65-97 80-98 80-98 ---	22-37 30-40 33-43 ---	3-14 8-17 12-20 ---
Vernon-----	0-5 5-22 22-60	Clay loam Silty clay Weathered bedrock	CL CH, CL	A-6, A-7 A-7	--- --- ---	0-15 0-15 ---	90-100 90-100 ---	85-95 70-100 ---	80-95 70-95 ---	60-95 60-90 ---	37-50 41-70 ---	14-25 20-40 ---
Ln: Lincoln-----	0-6 6-60	Fine sandy loam Stratified fine sand to clay loam	CL-ML, ML, SC-SM, SM SM, SP-SM	A-4 A-2, A-3	0 0	0 0	100 100	98-100 98-100	94-100 82-100	36-60 5-35	15-24 ---	NP-7 NP
LNN: Lincoln-----	0-10 10-60	Loamy sand Sand	SM SM, SP-SM	A-2 A-2, A-3	0 0	0 0	100 100	98-100 98-100	90-100 82-100	15-35 5-35	--- ---	NP NP
Ma: Mangum-----	0-10 10-50 50-60	Clay Clay Clay loam	CH, CL CH, CL CL	A-7-6 A-7-6 A-6, A-7-6	0 0 0	0 0 0	100 100 100	100 100 100	98-100 95-100 95-100	90-100 90-100 90-100	40-70 41-70 28-50	20-45 20-45 11-25
Mg: Mangum-----	0-10 10-50 50-60	Clay Clay Clay loam	CH, CL CH, CL CL	A-7-6 A-7-6 A-6, A-7-6	0 0 0	0 0 0	100 100 100	100 100 100	98-100 95-100 95-100	90-100 90-100 90-100	40-70 41-70 28-50	20-45 20-45 11-25
Drummond-----	0-8 8-30 30-60	Silty clay loam Clay loam Variable	CL CH, CL ---	A-6, A-7 A-6, A-7	0 0 ---	0 0 ---	100 100 ---	100 100 ---	96-100 96-100 ---	80-98 80-98 ---	37-50 35-60 ---	15-26 15-35 ---
Mn: Minco-----	0-27 27-60	Silt loam Silt loam	CL, CL-ML, ML CL, ML, SC, SM	A-4 A-4	0 0	0 0	100 100	100 98-100	94-100 94-100	51-97 36-97	15-31 15-31	NP-10 NP-10
Na: Naron-----	0-12 12-37 37-60	Fine sandy loam Fine sandy loam Fine sandy loam	CL-ML, ML, SC-SM, SM CL, SC SC-SM, SM	A-2, A-4 A-4, A-6 A-2, A-4	0 0 0	0 0 0	100 100 100	95-100 95-100 95-100	75-100 80-100 75-100	25-60 36-60 20-50	15-26 26-40 15-26	1-7 8-18 NP-7
Nb: Naron-----	0-12 12-37 37-60	Fine sandy loam Fine sandy loam Fine sandy loam	CL-ML, ML, SC-SM, SM CL, SC SC-SM, SM	A-2, A-4 A-4, A-6 A-2, A-4	0 0 0	0 0 0	100 100 100	95-100 95-100 95-100	75-100 80-100 75-100	25-60 36-60 20-50	15-26 26-40 15-26	1-7 8-18 NP-7
Os: Ost-----	0-8 8-22 22-60	Clay loam Clay loam Clay loam	CL CL CL, ML, SC, SM	A-6 A-4, A-6 A-2, A-4, A-6	0 0 0	0 0 0	95-100 95-100 85-100	95-100 90-100 85-100	85-100 85-100 80-100	75-90 55-90 30-90	30-40 30-40 15-40	10-20 9-18 NP-18
Ot: Ost-----	0-8 8-22 22-60	Clay loam Clay loam Clay loam	CL CL CL, ML, SC, SM	A-6 A-4, A-6 A-2, A-4, A-6	0 0 0	0 0 0	95-100 95-100 85-100	95-100 90-100 85-100	85-100 85-100 80-100	75-90 55-90 30-90	30-40 30-40 15-40	10-20 9-18 NP-18
Pa: Pond Creek-----	0-11 11-60	Silt loam Silty clay loam	CL, CL-ML, ML CL	A-4, A-6 A-4, A-6, A-7	0 0	0 0	100 100	100 100	96-100 96-100	65-97 65-98	22-37 30-43	3-14 8-20
Pd: Pond Creek-----	0-11 11-60	Silt loam Silty clay loam	CL, CL-ML, ML CL	A-4, A-6 A-4, A-6, A-7	0 0	0 0	100 100	100 100	96-100 96-100	65-97 65-98	22-37 30-43	3-14 8-20
Ph: Dale-----	0-20 20-60	Silt loam Silt loam	CL, CL-ML CL	A-4, A-6 A-4, A-6, A-7	0 0	0 0	95-100 95-100	95-100 95-100	90-100 90-100	65-98 65-98	25-35 30-43	5-15 8-20
Pk: Buttermilk-----	0-20 20-32 32-60	Silt loam Silt loam Silt loam	CL, CL-ML CL CL	A-4, A-6 A-4, A-6, A-7 A-4, A-6, A-7	0 0 0	0 0 0	100 100 100	100 100 100	90-100 90-100 90-100	80-100 85-100 85-100	25-35 30-50 30-50	5-15 8-25 8-25
Ps: Pratt-----	0-7 7-30 30-60	Loamy fine sand Loamy fine sand Loamy fine sand	SM SC-SM, SM SM, SP-SM	A-2 A-2, A-4 A-2, A-3	0 0 0	0 0 0	100 100 100	95-100 95-100 95-100	70-100 90-100 80-100	15-35 15-40 5-35	--- 15-20 ---	NP NP-6 NP

ENGINEERING INDEX PROPERTIES--Continued
Barber County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
Pt:												
Pratt-----	0-7	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	---	NP
	7-30	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	30-60	Loamy fine sand	SP-SM, SM	A-2, A-3	0	0	100	95-100	80-100	5-35	---	NP
Tivoli-----	0-5	Loamy fine sand	SM	A-2	0	0	100	95-100	90-100	15-35	---	NP
	5-60	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-25	---	NP
Qn:												
Quinlan-----	0-14	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	90-100	51-97	15-37	NP-14
	>14	Weathered bedrock			---	---	---	---	---	---	---	---
Qw:												
Quinlan-----	0-14	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	90-100	51-97	15-37	NP-14
	>14	Weathered bedrock			---	---	---	---	---	---	---	---
Woodward-----	0-40	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	51-95	15-31	NP-12
	>40	Weathered bedrock			---	---	---	---	---	---	---	---
Rb:												
Knoco-----	0-9	Clay	CH, CL	A-6, A-7-6	0	0-2	90-100	80-100	75-100	60-90	32-60	14-38
	9-19		CH, CL	A-6, A-7-6	0	0-2	90-100	85-100	70-100	70-100	30-60	13-38
	19-60		CH, CL	A-6, A-7-6	0	0-2	90-100	85-100	70-100	70-100	30-60	13-38
Rock Outcrop---	0-10	Unweathered bedrock			---	---	---	---	---	---	---	---
Sb:												
Shellabarger---	0-14	Sandy loam	ML, SM	A-2, A-4	0	0	95-100	95-100	75-100	30-55	15-30	NP-5
	14-48	Sandy clay loam	SC	A-4, A-6	0	0	95-100	85-100	70-90	35-50	25-40	8-20
	48-60	Coarse sandy loam	SC, SC-SM, SM, SP-SM	A-2, A-4	0	0	80-100	70-100	50-80	10-40	15-30	NP-10
SBB:												
Shellabarger---	0-11	Fine sandy loam	ML, SM	A-2, A-4	0	0	95-100	95-100	75-100	30-55	15-30	NP-5
	11-34	Sandy clay loam	SC	A-4, A-6	0	0	95-100	85-100	70-90	35-50	25-40	8-20
	34-60	Coarse sandy loam	SC, SC-SM, SM, SP-SM	A-2, A-4	0	0	80-100	70-100	50-80	10-40	15-30	NP-10
Sc:												
Shellabarger---	0-14	Sandy loam	ML, SM	A-2, A-4	0	0	95-100	95-100	75-100	30-55	15-30	NP-5
	14-48	Sandy clay loam	SC	A-4, A-6	0	0	95-100	85-100	70-90	35-50	25-40	8-20
	48-60	Coarse sandy loam	SC, SC-SM, SM, SP-SM	A-2, A-4	0	0	80-100	70-100	50-80	10-40	15-30	NP-10
Tv:												
Tivoli-----	0-5	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-25	---	NP
	5-60	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-25	---	NP
Vn:												
Vernon-----	0-5	Clay loam	CL	A-6, A-7	---	0-15	90-100	85-95	80-95	60-95	37-50	14-25
	5-22	Clay	CH, CL	A-7	---	0-15	90-100	70-100	70-95	60-90	41-70	20-40
	22-60	Weathered bedrock			---	---	---	---	---	---	---	---
Vr:												
Vernon-----	0-5	Clay loam	CL	A-6, A-7	---	0-15	90-100	85-95	80-95	60-95	37-50	14-25
	5-22	Clay	CH, CL	A-7	0	0-15	90-100	70-100	70-95	60-90	41-70	20-40
	22-60	Weathered bedrock			---	---	---	---	---	---	---	---
Vs:												
Vernon-----	0-5	Clay loam	CL	A-6, A-7	---	0-15	90-100	85-95	80-95	60-95	37-50	14-25
	5-22	Silty clay	CH, CL	A-7	0	0-15	90-100	70-100	70-95	60-90	41-70	20-40
	22-60	Weathered bedrock			---	---	---	---	---	---	---	---
Knoco-----	0-9	Clay	CH, CL	A-6, A-7-6	0	0-2	90-100	80-100	75-100	60-90	32-60	14-38
	9-19	Clay	CH, CL	A-6, A-7-6	0	0-2	90-100	85-100	70-100	70-100	30-60	13-38
	19-60		CH, CL	A-6, A-7-6	0	0-2	90-100	85-100	70-100	70-100	30-60	13-38
W:												
Water-----	---	---	---	---	---	---	---	---	---	---	---	---
Wa:												
Waldeck-----	0-12	Sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	95-100	75-100	25-55	15-25	NP-5
	12-36	Sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	30-50	15-25	NP-5
	36-60	Sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	40-60	1-35	---	NP
Wo:												
Woodward-----	0-40	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	51-95	15-31	NP-12
	>40	Weathered bedrock			---	---	---	---	---	---	---	---
Quinlan-----	0-14	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	90-100	51-97	15-37	NP-14
	>14	Weathered bedrock			---	---	---	---	---	---	---	---
Ws:												
Woodward-----	0-40	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	51-95	15-31	NP-12
	>40	Weathered bedrock			---	---	---	---	---	---	---	---
Quinlan-----	0-14	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	90-100	51-97	15-37	NP-14
	>14	Weathered bedrock			---	---	---	---	---	---	---	---
Ya:												
Yahola-----	0-14	Sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	95-100	90-100	36-60	15-26	NP-7
	14-60	Fine sandy loam	CL, ML, SC, SM, CL-ML	A-4	0	0	100	95-100	90-100	36-85	15-30	NP-10
	60-66	Stratified loam to loamy fine sand	CL, ML, SC, SM, CL-ML	A-2, A-4	0	0	100	95-100	90-100	15-85	15-30	NP-10

ENGINEERING INDEX PROPERTIES--Continued
Barber County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
Ze: Zenda-----	In											
	0-20	Clay loam	CL	A-6	0	0	100	95-100	85-100	55-80	30-40	10-20
	20-60	Clay loam	CL	A-6	0	0	100	95-100	85-100	55-80	25-40	10-25

Physical Properties table shows estimates of some physical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Particle size is the effective diameter of a soil particle as measured by sedimentation, sieving, or micrometric methods. Particle sizes are expressed as classes with specific effective diameter class limits. The broad classes are sand, silt, and clay, ranging from the larger to the smaller.

Sand as a soil separate consists of mineral soil particles that are 0.05 millimeter to 2 millimeters in diameter. In this table, the estimated sand content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Silt as a soil separate consists of mineral soil particles that are 0.002 to 0.05 millimeter in diameter. In this table, the estimated silt content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Clay as a soil separate consists of mineral soil particles that are less than 0.002 millimeter in diameter. In this table, the estimated clay content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of sand, silt, and clay affects the physical behavior of a soil. Particle size is important for engineering and agronomic interpretations, for determination of soil hydrologic qualities, and for soil classification.

The amount and kind of clay affect the fertility and physical condition of the soil and the ability of the soil to adsorb cations and to retain moisture. They influence shrink-swell potential, permeability, plasticity, the ease of soil dispersion, and other soil properties. The amount and kind of clay in a soil also affect tillage and earth moving operations.

Moist bulk density is the weight of soil (oven-dry) per unit volume. Volume is measured when the soil is at field moisture capacity, that is, the moisture content at 1/3- or 1/10-bar (33kPa or 10kPa) moisture tension. Weight is determined after the soil is dried at 105 degrees C. In the table, the estimated moist bulk density of each soil horizon is expressed in grams per cubic centimeter of soil material that is less than 2 millimeters in diameter. Bulk density data are used to compute shrink-swell potential, available water capacity, total pore space, and other soil properties. The moist bulk density of a soil indicates the pore space available for water and roots. Depending on soil texture, a bulk density of more than 1.4 can restrict water storage and root penetration. Moist bulk density is influenced by texture, kind of clay, content of organic matter, and soil structure.

Permeability ($K \rightarrow \text{sat}$) refers to the ability of a soil to transmit water or air. The term "permeability," as used in soil surveys, indicates saturated hydraulic conductivity ($K \rightarrow \text{sat}$). The estimates in the table indicate the rate of water movement, in inches per hour, when the soil is saturated. They are based on soil characteristics observed in the field, particularly structure, porosity, and texture. Permeability is considered in the design of soil drainage systems and septic tank absorption fields.

Available water capacity refers to the quantity of water that the soil is capable of storing for use by plants. The capacity for water storage is given in inches of water per inch of soil for each soil layer. The capacity varies, depending on soil properties that affect retention of water. The most important properties are the content of organic matter, soil texture, bulk density, and soil structure. Available water capacity is an important factor in the choice of plants or crops to be grown and in the design and management of irrigation systems. Available water capacity is not an estimate of the quantity of water actually available to plants at any given time.

Linear extensibility refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. It is an expression of the volume change between the water content of the clod at 1/3- or 1/10-bar tension (33kPa or 10kPa tension) and oven dryness. The volume change is reported in the table as percent change for the whole soil. Volume change is influenced by the amount and type of clay minerals in the soil.

Linear extensibility is used to determine the shrink-swell potential of soils. The shrink-swell potential is low if the soil has a linear extensibility of less than 3 percent; moderate if 3 to 6 percent; high if 6 to 9 percent; and very high if more than 9 percent. If the linear extensibility is more than 3, shrinking and swelling can cause damage to buildings, roads, and other structures and to plant roots. Special design commonly is needed.

Organic matter is the plant and animal residue in the soil at various stages of decomposition. In Physical Properties table, the estimated content of organic matter is expressed as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter. The content of organic matter in a soil can be maintained by returning crop residue to the soil. Organic matter has a positive effect on available water capacity, water infiltration, soil organism activity, and tilth. It is a source of nitrogen and other nutrients for crops and soil organisms.

Erosion factors are shown in the Physical Properties table as the K factor (K_w and K_f) and the T factor. Erosion factor K indicates the susceptibility of a soil to sheet and rill erosion by water. Factor K is one of six factors used in the Universal Soil Loss Equation (USLE) and the Revised Universal Soil Loss Equation (RUSLE) to predict the average annual rate of soil loss by sheet and rill erosion in tons per acre per year. The estimates are based primarily on percentage of silt, sand, and organic matter and on soil structure and permeability. Values of K range from 0.02 to 0.69. Other factors being equal, the higher the value, the more susceptible the soil is to sheet and rill erosion by water.

Erosion factor K_w indicates the erodibility of the whole soil. The estimates are modified by the presence of rock fragments.

Erosion factor K_f indicates the erodibility of the fine-earth fraction, or the material less than 2 millimeters in size.

Erosion factor T is an estimate of the maximum average annual rate of soil erosion by wind or water that can occur without affecting crop productivity over a sustained period. The rate is in tons per acre per year.

Wind erodibility groups are made up of soils that have similar properties affecting their susceptibility to

wind erosion in cultivated areas. The soils assigned to group 1 are the most susceptible to wind erosion, and those assigned to group 8 are the least susceptible. The groups are as follows:

1. Coarse sands, sands, fine sands, and very fine sands.
2. Loamy coarse sands, loamy sands, loamy fine sands, loamy very fine sands, ash material, and sapric soil material.
3. Coarse sandy loams, sandy loams, fine sandy loams, and very fine sandy loams.
- 4L. Calcareous loams, silt loams, clay loams, and silty clay loams.
4. Clays, silty clays, noncalcareous clay loams, and silty clay loams that are more than 35 percent clay.
5. Noncalcareous loams and silt loams that are less than 20 percent clay and sandy clay loams, sandy clays, and hemic soil material.
6. Noncalcareous loams and silt loams that are more than 20 percent clay and noncalcareous clay loams that are less than 35 percent clay.
7. Silts, noncalcareous silty clay loams that are less than 35 percent clay, and fibric soil material.
8. Soils that are not subject to wind erosion because of coarse fragments on the surface or because of surface wetness.

Wind erodibility index is a numerical value indicating the susceptibility of soil to wind erosion, or the tons per acre per year that can be expected to be lost to wind erosion. There is a close correlation between wind erosion and the texture of the surface layer, the size and durability of surface clods, rock fragments, organic matter, and a calcareous reaction. Soil moisture and frozen soil layers also influence wind erosion.

Explanation of Wind Erodibility Groups

Soil erodibility by wind is directly related to the percentage of dry non-erodible surface soil aggregates larger than 0.84 mm in diameter. From this percentage, the wind erodibility index (I-factor) is determined. The I-factor is an expression of the stability of these soil aggregates against breakdown by tillage and abrasion from wind erosion. Soils are placed in Wind Erodibility Groups (WEG) having similar percentages of dry soil aggregates larger than 0.84 mm as shown in the following table.

WEG	Properties of Soil Surface Layer	Dry Soil Aggregates >0.84mm Percent	Wind Erodibility Index T/Ac/Yr (I)
1	Very fine sand, fine sand, sand, or coarse sand	1 2 3 5 7	310 1/ 250 220 180 160
2	Loamy very fine sand, loamy fine sand, loamy sand, loamy coarse sand, organic soil materials.	10	134
3	Very fine sandy loam, fine sandy loam, sandy loam, or coarse sandy loam.	25	86
4	Clay, silty clay, non-calcareous clay loam, or silty clay loam with >35 percent clay content.	25	86
4L	Calcareous 2/ loam, silt loam, clay loam, or silty clay loam.	25	86
5	Non-calcareous loam and silt loam with <20 percent clay content, or sandy clay loam, sandy clay, and hemic 3/ organic soil materials.	40	56
6	Non-calcareous loam and silt loam with >20 percent clay content, or non-calcareous clay loam with <35 percent clay content.	45	48
7	Silt, non-calcareous silty clay loam with >35 percent clay content and fibric 3/ organic soil material.	50	38
8	Soils not suitable for cultivation due to coarse fragments or wetness; wind erosion is not a problem.	--	0

1/ The "I" values for WEG 1 vary from 160 for coarse sands to 310 for very fine sands. Use an "I" of 220 as an average figure. For coarser sand that has gravel, use a lower figure. For a soil that has no gravel and very fine sand, use a higher figure. (Modification for coarse fragments is preparation.)

2/ Calcareous is a strongly or violently effervescent reaction to cold dilute (1N) HCL.

3/ See Soil Taxonomy for definition.

PHYSICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
033CP: Clairemont---	0-15	37	42	15-27	1.40-1.60	0.60-2.00	0.16-0.22	0.0-2.9	0.0-2.0	.43	.43	5	4L	86
	15-60	9	64	18-35	1.40-1.65	0.60-2.00	0.16-0.22	0.0-2.9	---	.43	.43			
033OA: Obaro-----	0-35	7	62	27-35	1.30-1.45	0.60-2.00	0.14-0.20	0.0-2.9	0.0-1.0	.43	.43	3	4L	86
	>35			---	---	---	---	---	---	---	---			
033OB: Obaro-----	0-35	7	62	27-35	1.30-1.45	0.60-2.00	0.14-0.20	0.0-2.9	0.0-1.0	.43	.43	3	4L	86
	>35			---	---	---	---	---	---	---	---			
Rock Outcrop-	---			---	---	---	---	---	---	---	---	-	---	---
033QR: Quinlan-----	0-14	42	37	15-27	1.30-1.55	0.60-2.00	0.13-0.24	0.0-2.9	0.0-1.0	.37	.37	2	4L	86
	>14			---	---	---	---	---	---	---	---			
Woodward----	0-30	43	43	10-18	1.30-1.60	0.60-2.00	0.13-0.20	0.0-2.9	0.5-2.0	.37	.37	3	4L	86
	>30			---	---	---	---	---	---	---	---			
033QT: Quinlan-----	0-14	42	37	15-27	1.30-1.55	0.60-2.00	0.13-0.24	0.0-2.9	0.0-1.0	.37	.37	2	4L	86
	>14			---	---	---	---	---	---	---	---			
Woodward----	0-30	43	43	10-18	1.30-1.60	0.60-2.00	0.13-0.20	0.0-2.9	0.5-2.0	.37	.37	3	4L	86
	>30			---	---	---	---	---	---	---	---			
033YE: Yahola-----	0-12	66	20	10-18	1.30-1.60	2.00-6.00	0.11-0.15	0.0-2.9	0.5-1.0	.20	.20	5	3	86
	12-40	62	26	5-18	1.40-1.70	2.00-6.00	0.11-0.20	0.0-2.9	---	.32	.32			
	40-60	45	43	5-18	1.50-1.70	2.00-6.00	0.07-0.20	0.0-2.9	---	.32	.32			
077BP: Woodward----	0-24	14	72	10-18	1.30-1.60	0.60-2.00	0.13-0.20	0.0-2.9	0.5-2.0	.37	.37	3	4L	86
	>24			---	---	---	---	---	---	---	---			
Port-----	0-27	12	69	12-26	1.30-1.55	0.60-2.00	0.15-0.24	0.0-2.9	1.0-3.0	.37	.37	5	6	48
	27-60	7	65	20-35	1.30-1.60	0.60-2.00	0.15-0.24	3.0-5.9	---	.37	.37			
077FU: Farnum-----	0-11	42	38	14-27	1.35-1.45	0.60-2.00	0.19-0.22	0.0-2.9	1.0-3.0	.28	.28	5	6	48
	11-41	34	36	25-35	1.40-1.50	0.60-2.00	0.15-0.19	3.0-5.9	---	.28	.28			
	41-60	38	41	12-29	1.40-1.55	0.60-2.00	0.13-0.16	0.0-2.9	---	.28	.28			
077GE: Gerlane-----	0-17	66	20	10-18	1.35-1.50	2.00-6.00	0.13-0.18	0.0-2.9	1.0-3.0	.20	.20	5	3	86
	17-40	66	20	10-18	1.35-1.50	2.00-6.00	0.12-0.17	0.0-2.9	---	.20	.20			
	40-48	84	2	10-18	1.50-1.60	2.00-6.00	0.08-0.13	0.0-2.9	---	.20	.20			
	48-60	36	39	20-30	1.50-1.60	0.20-2.00	0.15-0.18	0.0-2.9	---	.20	.20			
077MC: Minco-----	0-42	14	73	8-18	1.35-1.60	0.60-2.00	0.13-0.24	0.0-2.9	1.0-3.0	.37	.37	5	5	56
	42-60	14	73	8-18	1.35-1.60	0.60-2.00	0.11-0.24	0.0-2.9	---	.37	.37			
077MN: Minco-----	0-42	14	73	8-18	1.35-1.60	0.60-2.00	0.13-0.24	0.0-2.9	1.0-3.0	.37	.37	5	5	56
	42-60	14	73	8-18	1.35-1.60	0.60-2.00	0.11-0.24	0.0-2.9	---	.37	.37			
077MO: Minco-----	0-42	14	73	8-18	1.35-1.60	0.60-2.00	0.13-0.24	0.0-2.9	1.0-3.0	.37	.37	5	5	56
	42-60	14	73	8-18	1.35-1.60	0.60-2.00	0.11-0.24	0.0-2.9	---	.37	.37			
077QU: Quinlan-----	0-9	42	37	15-27	1.30-1.55	0.60-2.00	0.13-0.24	0.0-2.9	0.0-1.0	.37	.37	2	4L	86
	>9			---	---	---	---	---	---	---	---			
077SE: Shellabarger-	0-13	68	20	8-16	1.35-1.50	0.60-2.00	0.13-0.21	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	13-38	59	18	18-27	1.45-1.60	0.60-2.00	0.16-0.18	0.0-2.9	---	.28	.32			
	38-60	66	24	3-18	1.50-1.65	0.60-2.00	0.05-0.16	0.0-2.9	---	.28	.32			
077WE: Quinlan-----	0-9	42	37	15-27	1.30-1.55	0.60-2.00	0.13-0.24	0.0-2.9	0.0-1.0	.37	.37	2	4L	86
	>9			---	---	---	---	---	---	---	---			
Woodward----	0-24	43	43	10-18	1.30-1.60	0.60-2.00	0.13-0.20	0.0-2.9	0.5-2.0	.37	.37	3	4L	86
	>24			---	---	---	---	---	---	---	---			
095AC: Albion-----	0-8	66	23	7-15	1.35-1.50	2.00-6.00	0.13-0.17	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	8-16	67	19	10-18	1.45-1.60	2.00-6.00	0.12-0.18	0.0-2.9	---	.20	.24			
	16-26	82	9	4-15	1.45-1.60	2.00-6.00	0.09-0.12	0.0-2.9	---	.15	.20			
	26-60	92	2	2-10	1.50-1.65	5.95-19.98	0.03-0.10	0.0-2.9	---	.15	.32			
095DA: Dillwyn-----	0-8	79	16	2-8	1.50-1.60	5.95-19.98	0.08-0.12	0.0-2.9	0.0-2.0	.17	.17	5	2	134
	8-60	79	16	2-8	1.50-1.60	5.95-19.98	0.06-0.10	0.0-2.9	---	.17	.17			
Plevna-----	0-11	67	20	8-18	1.40-1.50	2.00-6.00	0.14-0.16	0.0-2.9	1.0-4.0	.20	.20	5	3	86
	11-36	67	20	8-18	1.40-1.50	2.00-6.00	0.12-0.16	0.0-2.9	---	.20	.20			
	36-60	95	1	1-7	1.50-1.60	2.00-6.00	0.05-0.07	0.0-2.9	---	.20	.20			
095FA: Farnum-----	0-16	66	23	8-14	1.45-1.55	2.00-6.00	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	16-50	34	36	25-35	1.40-1.50	0.60-2.00	0.15-0.19	3.0-5.9	---	.28	.28			
	50-60	42	38	12-29	1.40-1.55	0.60-2.00	0.13-0.16	0.0-2.9	---	.28	.28			
095PB: Pratt-----	0-12	79	16	2-8	1.40-1.55	5.95-19.98	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	12-36	86	7	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	---	.17	.17			
	36-60	79	16	1-8	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	---	.17	.17			
095SB: Shellabarger-	0-10	68	20	8-16	1.35-1.50	0.60-2.00	0.13-0.21	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	10-45	60	18	18-27	1.45-1.60	0.60-2.00	0.16-0.18	0.0-2.9	---	.28	.32			
	45-60	66	24	3-18	1.50-1.65	0.60-2.00	0.05-0.16	0.0-2.9	---	.28	.32			

PHYSICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
095WA:														
Waldeck-----	0-12	68	20	8-16	1.50-1.60	2.00-6.00	0.14-0.18	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	12-36	68	20	8-16	1.50-1.60	2.00-6.00	0.12-0.17	0.0-2.9	---	.20	.20			
	36-60	97	1	1-4	1.55-1.65	5.95-19.98	0.05-0.07	0.0-2.9	---	.20	.24			
097LN:														
Lincoln-----	0-8	67	19	10-18	1.30-1.60	5.95-19.98	0.10-0.15	0.0-2.9	0.0-1.0	.20	.20	5	3	86
	8-60			5-15	1.30-1.60	5.95-19.98	0.02-0.08	0.0-2.9	---	.17	.17			
097QW:														
Quinlan-----	0-15	42	37	15-27	1.30-1.55	0.60-2.00	0.13-0.24	0.0-2.9	0.0-1.0	.37	.37	2	4L	86
	>15			---	---	---	---	---	---	---	---			
Woodward-----	0-27	43	43	10-18	1.30-1.60	0.60-2.00	0.13-0.20	0.0-2.9	0.5-2.0	.37	.37	3	4L	86
	>27			---	---	---	---	---	---	---	---			
097WA:														
Waldeck-----	0-14	45	41	10-18	1.40-1.55	2.00-6.00	0.18-0.20	0.0-2.9	1.0-3.0	.28	.28	4	5	56
	14-41	68	20	8-16	1.50-1.60	2.00-6.00	0.12-0.17	0.0-2.9	---	.20	.20			
	41-60	97	1	1-4	1.55-1.65	5.95-19.98	0.05-0.07	0.0-2.9	---	.20	.24			
151AB:														
Albion-----	0-8	66	23	7-15	1.35-1.50	2.00-6.00	0.13-0.17	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	8-18	67	19	10-18	1.45-1.60	2.00-6.00	0.12-0.18	0.0-2.9	---	.20	.24			
	18-29	66	24	4-15	1.45-1.60	2.00-6.00	0.09-0.12	0.0-2.9	---	.17	.20			
	29-60	92	2	2-10	1.50-1.65	5.95-19.98	0.03-0.10	0.0-2.9	---	.15	.32			
151AO:														
Albion-----	0-8	66	23	7-15	1.35-1.50	2.00-6.00	0.13-0.17	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	8-18	67	19	10-18	1.45-1.60	2.00-6.00	0.12-0.18	0.0-2.9	---	.20	.24			
	18-29	66	24	4-15	1.45-1.60	2.00-6.00	0.09-0.12	0.0-2.9	---	.17	.20			
	29-60	92	2	2-10	1.50-1.65	5.95-19.98	0.03-0.10	0.0-2.9	---	.15	.32			
151AS:														
Albion-----	0-8	66	23	7-15	1.35-1.50	2.00-6.00	0.13-0.17	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	8-18	67	19	10-18	1.45-1.60	2.00-6.00	0.12-0.18	0.0-2.9	---	.20	.24			
	18-29	66	24	4-15	1.45-1.60	2.00-6.00	0.09-0.12	0.0-2.9	---	.17	.20			
	29-60	92	2	2-10	1.50-1.65	5.95-19.98	0.03-0.10	0.0-2.9	---	.15	.32			
Shellabarger-														
	0-11	68	20	8-16	1.35-1.50	0.60-2.00	0.13-0.21	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	11-34	60	18	18-27	1.45-1.60	0.60-2.00	0.16-0.18	0.0-2.9	---	.28	.32			
	34-60	66	24	3-18	1.50-1.65	0.60-2.00	0.05-0.16	0.0-2.9	---	.28	.32			
151BC:														
Blanket-----	0-13	20	49	27-35	1.30-1.45	0.60-2.00	0.15-0.20	3.0-5.9	1.0-3.0	.37	.37	5	6	48
	13-46	8	50	35-50	1.35-1.55	0.20-0.60	0.12-0.18	3.0-5.9	---	.43	.43			
	46-60	7	51	35-50	1.35-1.55	0.60-2.00	0.12-0.18	3.0-5.9	---	.43	.37			
151CA:														
Carwile-----	0-12	62	26	5-18	1.30-1.65	0.60-2.00	0.11-0.20	0.0-2.9	1.0-3.0	.24	.24	5	3	86
	12-20	35	33	25-39	1.45-1.75	0.20-2.00	0.12-0.20	3.0-5.9	---	.37	.37			
	20-33	25	27	35-60	1.35-1.75	0.06-0.20	0.12-0.20	6.0-8.9	---	.37	.37			
	33-60	35	33	20-45	1.35-1.75	0.20-2.00	0.12-0.20	6.0-8.9	---	.32	.32			
151CK:														
Case-----	0-6	34	37	27-32	1.35-1.45	0.60-2.00	0.17-0.22	0.0-2.9	0.5-2.0	.32	.32	5	4L	86
	6-60	35	38	18-35	1.35-1.70	0.60-2.00	0.15-0.19	3.0-5.9	---	.32	.32			
Clark-----	0-8	34	37	27-32	1.35-1.45	0.60-2.00	0.17-0.22	3.0-5.9	1.0-2.0	.28	.28	5	4L	86
	8-60	35	38	18-35	1.35-1.70	0.60-2.00	0.14-0.19	3.0-5.9	---	.28	.28			
151CM:														
Clark-----	0-8	34	37	27-32	1.35-1.45	0.60-2.00	0.17-0.22	3.0-5.9	1.0-2.0	.28	.28	5	4L	86
	8-60	35	38	18-35	1.35-1.70	0.60-2.00	0.14-0.19	3.0-5.9	---	.28	.28			
151CN:														
Clark-----	0-8	65	20	10-20	1.30-1.45	0.60-2.00	0.15-0.19	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	8-60	35	38	18-35	1.35-1.70	0.60-2.00	0.14-0.19	3.0-5.9	---	.28	.28			
151CO:														
Clark-----	0-8	34	37	27-32	1.35-1.45	0.60-2.00	0.17-0.22	3.0-5.9	1.0-2.0	.28	.28	5	4L	86
	8-60	35	38	18-35	1.35-1.70	0.60-2.00	0.14-0.19	3.0-5.9	---	.28	.28			
Ost-----	0-9	34	37	27-30	1.35-1.40	0.20-0.60	0.15-0.20	0.0-2.9	1.0-3.0	.32	.32	5	6	48
	9-14	35	38	20-34	1.45-1.65	0.20-0.60	0.15-0.20	3.0-5.9	---	.32	.32			
	14-23	36	38	18-34	1.40-1.60	0.20-0.60	0.15-0.20	3.0-5.9	---	.32	.32			
	23-60	39	43	5-30	1.40-1.60	0.20-0.60	0.13-0.20	0.0-2.9	---	.32	.37			
151FA:														
Farnum-----	0-7	34	38	27-29	1.35-1.45	0.60-2.00	0.19-0.22	0.0-2.9	1.0-3.0	.28	.28	5	6	48
	7-37	34	36	25-35	1.40-1.50	0.60-2.00	0.15-0.19	3.0-5.9	---	.28	.28			
	37-60	61	19	12-29	1.40-1.55	0.60-2.00	0.13-0.16	0.0-2.9	---	.28	.28			
151KP:														
Kanza-----	0-11	86	7	3-12	1.50-1.70	5.95-19.98	0.08-0.13	0.0-2.9	1.0-3.0	.17	.17	5	2	134
	11-40	92	1	1-12	1.50-1.70	5.95-19.98	0.06-0.11	0.0-2.9	---	.17	.20			
Plevna-----	0-10	67	20	8-18	1.40-1.50	2.00-6.00	0.14-0.16	0.0-2.9	1.0-4.0	.20	.20	5	3	86
	10-40	67	20	8-18	1.40-1.50	2.00-6.00	0.12-0.16	0.0-2.9	---	.20	.20			
	40-60	95	1	1-7	1.50-1.60	2.00-6.00	0.05-0.07	0.0-2.9	---	.20	.20			
151OS:														
Ost-----	0-9	34	37	27-30	1.35-1.40	0.20-0.60	0.15-0.20	0.0-2.9	1.0-3.0	.32	.32	5	6	48
	9-14	35	38	20-34	1.45-1.65	0.20-0.60	0.15-0.20	3.0-5.9	---	.32	.32			
	14-23	36	38	18-34	1.40-1.60	0.20-0.60	0.15-0.20	3.0-5.9	---	.32	.32			
	23-60	39	43	5-30	1.40-1.60	0.20-0.60	0.13-0.20	0.0-2.9	---	.32	.37			
151PM:														
Pratt-----	0-10	79	16	2-8	1.40-1.55	5.95-19.98	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	10-40	86	7	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	---	.17	.17			
	40-60	79	16	1-8	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	---	.17	.17			
151PN:														
Pratt-----	0-10	79	16	2-8	1.40-1.55	5.95-19.98	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	10-40	86	7	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	---	.17	.17			
	40-60	79	16	1-8	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	---	.17	.17			

PHYSICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
151PO:														
Pratt-----	0-10	79	16	2-8	1.40-1.55	5.95-19.98	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	10-40	86	7	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	---	.17	.17			
	40-60	79	16	1-8	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	---	.17	.17			
Carwile-----	0-12	62	26	5-18	1.30-1.65	0.60-2.00	0.11-0.20	0.0-2.9	1.0-3.0	.24	.24	5	3	86
	12-20	54	14	25-39	1.45-1.75	0.20-2.00	0.12-0.20	3.0-5.9	---	.37	.37			
	20-33	23	29	35-60	1.35-1.75	0.06-0.20	0.12-0.20	6.0-8.9	---	.37	.37			
	33-60	54	14	20-45	1.35-1.75	0.20-2.00	0.12-0.20	6.0-8.9	---	.32	.32			
151SA:														
Albion-----	0-8	63	26	7-15	1.35-1.50	2.00-6.00	0.13-0.17	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	8-18	67	19	10-18	1.45-1.60	2.00-6.00	0.12-0.18	0.0-2.9	---	.20	.24			
	18-29	66	24	4-15	1.45-1.60	2.00-6.00	0.09-0.12	0.0-2.9	---	.17	.20			
	29-60	92	2	2-10	1.50-1.65	5.95-19.98	0.03-0.10	0.0-2.9	---	.15	.32			
Kaski-----	0-26	40	38	18-27	1.35-1.45	0.60-2.00	0.18-0.22	3.0-5.9	1.0-3.0	.28	.28	5	6	48
	26-40	38	36	18-35	1.40-1.50	0.60-2.00	0.13-0.19	3.0-5.9	---	.28	.28			
	40-60	66	15	8-30	1.45-1.55	0.60-2.00	0.13-0.19	0.0-2.9	---	.28	.28			
151SE:														
Shellabarger--	0-11	68	20	8-16	1.35-1.50	0.60-2.00	0.13-0.21	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	11-34	60	18	18-27	1.45-1.60	0.60-2.00	0.16-0.18	0.0-2.9	---	.28	.32			
	34-60	66	24	3-18	1.50-1.65	0.60-2.00	0.05-0.16	0.0-2.9	---	.28	.32			
1439:														
Crisfield----	0-12	70	23	5-12	1.35-1.50	2.00-6.00	0.05-0.18	0.0-2.9	0.2-0.8	.15	.15	5	3	86
	12-24	70	22	5-12	1.45-1.65	2.00-6.00	0.05-0.18	0.0-2.9	0.2-0.8	.15	.15			
	24-80	88	9	2-8	1.50-1.65	6.00-19.98	0.00-0.12	0.0-2.9	0.0-0.1	.05	.05			
Ad:														
Albion-----	0-8	66	23	7-15	1.35-1.50	2.00-6.00	0.13-0.17	0.0-2.9	1.0-2.0	.20	.24	4	3	86
	8-16	67	19	10-18	1.45-1.60	2.00-6.00	0.12-0.18	0.0-2.9	---	.20	.24			
	16-27	82	9	4-15	1.45-1.60	2.00-6.00	0.09-0.12	0.0-2.9	---	.17	.20			
	27-60	92	2	2-10	1.50-1.65	5.95-19.98	0.03-0.10	0.0-2.9	---	.15	.32			
Shellabarger--	0-14	68	20	8-16	1.35-1.50	0.60-2.00	0.13-0.21	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	14-48	60	18	18-27	1.45-1.60	0.60-2.00	0.16-0.18	0.0-2.9	---	.28	.32			
	48-60	66	24	3-18	1.50-1.65	0.60-2.00	0.05-0.16	0.0-2.9	---	.28	.32			
Ae:														
Albion-----	0-8	66	23	7-15	1.35-1.50	2.00-6.00	0.13-0.17	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	8-16	67	19	10-18	1.45-1.60	2.00-6.00	0.12-0.18	0.0-2.9	---	.20	.24			
	16-27	82	9	4-15	1.45-1.60	2.00-6.00	0.09-0.12	0.0-2.9	---	.17	.20			
	27-60	92	2	2-10	1.50-1.65	5.95-19.98	0.03-0.10	0.0-2.9	---	.15	.32			
Shellabarger--	0-14	68	20	8-16	1.35-1.50	0.60-2.00	0.13-0.21	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	14-48	60	18	18-27	1.45-1.60	0.60-2.00	0.16-0.18	0.0-2.9	---	.28	.32			
	48-60	66	24	3-18	1.50-1.65	0.60-2.00	0.05-0.16	0.0-2.9	---	.28	.32			
AED:														
Arents,	---			---	---	---	---	---	---	---	---	---	---	---
Earthen Dam-														
AN:														
Albion-----	0-8	66	23	7-15	1.35-1.50	2.00-6.00	0.13-0.17	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	8-16	67	19	10-18	1.45-1.60	2.00-6.00	0.12-0.18	0.0-2.9	---	.20	.24			
	16-28	82	9	4-15	1.45-1.60	2.00-6.00	0.09-0.12	0.0-2.9	---	.17	.20			
	28-60	92	2	2-10	1.50-1.65	5.95-19.98	0.03-0.10	0.0-2.9	---	.15	.32			
Shellabarger--	0-11	68	20	8-16	1.35-1.50	0.60-2.00	0.13-0.21	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	11-38	60	18	18-27	1.45-1.60	0.60-2.00	0.16-0.18	0.0-2.9	---	.28	.32			
	38-60	66	24	3-18	1.50-1.65	0.60-2.00	0.05-0.16	0.0-2.9	---	.28	.32			
As:														
Clairemont---	0-8	10	68	18-27	1.30-1.45	0.60-2.00	0.10-0.15	0.0-2.9	0.0-2.0	.43	.43	5	4L	86
	8-60	32	42	18-35	1.35-1.55	0.60-2.00	0.10-0.15	0.0-2.9	---	.43	.43			
At:														
Attica-----	0-12	87	7	2-10	1.50-1.60	2.00-6.00	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	12-30	67	20	8-18	1.50-1.60	2.00-6.00	0.12-0.17	0.0-2.9	---	.24	.24			
	30-60	88	1	4-18	1.50-1.60	2.00-6.00	0.08-0.16	0.0-2.9	---	.24	.28			
Ba:														
Blanket-----	0-9	26	53	15-27	1.30-1.50	0.60-2.00	0.15-0.20	0.0-2.9	1.0-3.0	.37	.37	5	5	56
	9-30	8	50	35-50	1.35-1.55	0.20-0.60	0.12-0.18	3.0-5.9	---	.43	.43			
	30-60	7	51	35-50	1.35-1.55	0.60-2.00	0.12-0.18	3.0-5.9	---	.43	.43			
Bb:														
Blanket-----	0-9	26	53	15-27	1.30-1.50	0.60-2.00	0.15-0.20	0.0-2.9	1.0-3.0	.37	.37	5	5	56
	9-30	8	50	35-50	1.35-1.55	0.20-0.60	0.12-0.18	3.0-5.9	---	.43	.43			
	30-60	7	51	35-50	1.35-1.55	0.60-2.00	0.12-0.18	3.0-5.9	---	.43	.43			
Bc:														
Blanket-----	0-9	20	49	27-35	1.30-1.45	0.60-2.00	0.15-0.20	3.0-5.9	1.0-3.0	.37	.37	5	6	48
	9-30	8	50	35-50	1.35-1.55	0.20-0.60	0.12-0.18	3.0-5.9	---	.43	.43			
	30-60	7	51	35-50	1.35-1.55	0.60-2.00	0.12-0.18	3.0-5.9	---	.43	.43			
Bf:														
Quinlan-----	0-14	42	37	15-27	1.30-1.55	0.60-2.00	0.13-0.24	0.0-2.9	0.0-1.0	.37	.37	2	4L	86
	>14			---	---	---	---	---	---	---	---			
Clairemont---	0-14	11	68	15-27	1.40-1.60	0.60-2.00	0.16-0.22	0.0-2.9	0.0-2.0	.43	.43	5	4L	86
	14-60	9	64	18-35	1.40-1.65	0.60-2.00	0.16-0.22	0.0-2.9	---	.43	.43			
Ca:														
Canadian-----	0-18	62	26	5-18	1.30-1.60	2.00-6.00	0.10-0.15	0.0-2.9	1.0-3.0	.20	.20	5	3	86
	18-48	66	20	10-18	1.40-1.70	2.00-6.00	0.10-0.20	0.0-2.9	---	.20	.20			
	48-60	62	26	5-18	1.40-1.70	1.98-19.98	0.07-0.20	0.0-2.9	---	.20	.20			
Cc:														
Case-----	0-6	34	37	27-32	1.35-1.45	0.60-2.00	0.17-0.22	0.0-2.9	0.5-2.0	.32	.32	5	4L	86
	6-60	35	38	18-35	1.35-1.70	0.60-2.00	0.15-0.19	3.0-5.9	---	.32	.32			
Clark-----	0-10	34	37	27-32	1.35-1.45	0.60-2.00	0.17-0.22	3.0-5.9	1.0-2.0	.28	.28	5	4L	86
	10-60	38	36	18-35	1.35-1.70	0.60-2.00	0.14-0.19	3.0-5.9	---	.28	.28			

PHYSICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Cd: Clairemont---	0-14	11	68	15-27	1.40-1.60	0.60-2.00	0.16-0.22	0.0-2.9	0.0-2.0	.43	.43	5	4L	86
	14-60	9	64	18-35	1.40-1.65	0.60-2.00	0.16-0.22	0.0-2.9	---	.43	.43			
Cf: Clairemont---	0-14	11	68	15-27	1.40-1.60	0.60-2.00	0.16-0.22	0.0-2.9	0.0-2.0	.43	.43	5	4L	86
	14-60	9	64	18-35	1.40-1.65	0.60-2.00	0.16-0.22	0.0-2.9	---	.43	.43			
Ck: Clark-----	0-10	34	37	27-32	1.35-1.45	0.60-2.00	0.17-0.22	3.0-5.9	1.0-2.0	.28	.28	5	4L	86
	10-60	34	38	18-35	1.35-1.70	0.60-2.00	0.14-0.19	3.0-5.9	---	.28	.28			
Fa: Farnum-----	0-9	63	26	8-14	1.45-1.55	2.00-6.00	0.13-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	9-60	34	36	25-35	1.40-1.50	0.60-2.00	0.15-0.19	3.0-5.9	---	.28	.28			
Fm: Farnum-----	0-9	42	38	14-27	1.35-1.45	0.60-2.00	0.19-0.22	0.0-2.9	1.0-3.0	.28	.28	5	6	48
	9-60	34	36	25-35	1.40-1.50	0.60-2.00	0.15-0.19	3.0-5.9	---	.28	.28			
Fr: Farnum-----	0-9	42	38	14-27	1.35-1.45	0.60-2.00	0.19-0.22	0.0-2.9	1.0-3.0	.28	.28	5	6	48
	9-60	34	36	25-35	1.40-1.50	0.60-2.00	0.15-0.19	3.0-5.9	---	.28	.28			
Fu: Farnum-----	0-9	34	38	27-29	1.35-1.45	0.60-2.00	0.19-0.22	0.0-2.9	1.0-3.0	.28	.28	5	6	48
	9-60	34	36	25-35	1.40-1.50	0.60-2.00	0.15-0.19	3.0-5.9	---	.28	.28			
Ga: Grant-----	0-13	11	68	15-26	1.30-1.50	0.60-6.00	0.15-0.20	0.0-2.9	1.0-3.0	.37	.37	4	5	56
	13-50	9	64	18-35	1.40-1.70	0.60-2.00	0.15-0.20	0.0-2.9	---	.37	.37			
	50-60	11	68	15-26	1.40-1.65	0.60-2.00	0.16-0.22	0.0-2.9	---	.37	.37			
Gb: Grant-----	0-13	11	68	15-26	1.30-1.50	0.60-6.00	0.15-0.20	0.0-2.9	1.0-3.0	.37	.37	4	5	56
	13-50	9	64	18-35	1.40-1.70	0.60-2.00	0.15-0.20	0.0-2.9	---	.37	.37			
	50-60	11	68	15-26	1.40-1.65	0.60-2.00	0.16-0.22	0.0-2.9	---	.37	.37			
Gc: Grant-----	0-13	11	68	15-26	1.30-1.50	0.60-6.00	0.15-0.20	0.0-2.9	1.0-3.0	.37	.37	4	5	56
	13-50	9	64	18-35	1.40-1.70	0.60-2.00	0.15-0.20	0.0-2.9	---	.37	.37			
	50-60	11	68	15-26	1.40-1.65	0.60-2.00	0.16-0.22	0.0-2.9	---	.37	.37			
GRP: Gravel Pits--	---			---	---	---	---	---	---	---	---	-	---	---
INT: Aquolls-----	0-60			---	---	---	---	---	---	---	---	-	---	0
Ka: Kanza-----	0-7	86	7	3-12	1.50-1.70	5.95-19.98	0.08-0.13	0.0-2.9	1.0-3.0	.17	.17	5	2	134
	7-48	87	7	1-12	1.50-1.70	5.95-19.98	0.06-0.11	0.0-2.9	---	.17	.20			
Kf: Kingfisher---	0-9	11	68	15-27	1.30-1.55	0.60-2.00	0.15-0.24	0.0-2.9	1.0-3.0	.37	.37	3	5	56
	9-17	7	63	25-35	1.40-1.70	0.20-0.60	0.15-0.24	3.0-5.9	---	.37	.37			
	17-28	7	62	27-35	1.45-1.70	0.20-0.60	0.15-0.22	3.0-5.9	---	.32	.32			
	>28			---	---	---	---	---	---	---	---			
Kv: Kingfisher---	0-9	11	68	15-27	1.30-1.55	0.60-2.00	0.15-0.24	0.0-2.9	1.0-3.0	.37	.37	3	5	56
	9-17	7	63	25-35	1.40-1.70	0.20-0.60	0.15-0.24	3.0-5.9	---	.37	.37			
	17-28	7	62	27-35	1.45-1.70	0.20-0.60	0.15-0.22	3.0-5.9	---	.32	.32			
	>28			---	---	---	---	---	---	---	---			
Vernon-----	0-5	35	34	27-35	1.30-1.60	0.20-0.60	0.15-0.22	3.0-5.9	0.5-1.0	.37	.37	3	4	86
	5-22	6	47	35-60	1.35-1.65	0.06-0.20	0.10-0.20	6.0-8.9	---	.37	.37			
	22-60			---	---	---	---	---	---	---	---			
Kz: Kingfisher---	0-9	11	68	15-27	1.30-1.55	0.60-2.00	0.15-0.24	0.0-2.9	1.0-3.0	.37	.37	3	5	56
	9-17	7	63	25-35	1.40-1.70	0.20-0.60	0.15-0.24	3.0-5.9	---	.37	.37			
	17-28	7	62	27-35	1.45-1.70	0.20-0.60	0.15-0.22	3.0-5.9	---	.32	.32			
	>28			---	---	---	---	---	---	---	---			
Vernon-----	0-5	35	34	27-35	1.30-1.60	0.20-0.60	0.15-0.22	3.0-5.9	0.5-1.0	.37	.37	3	4	86
	5-22	6	47	35-60	1.35-1.65	0.06-0.20	0.10-0.20	6.0-8.9	---	.37	.37			
	22-60			---	---	---	---	---	---	---	---			
Ln: Lincoln-----	0-6	66	20	10-18	1.30-1.60	5.95-19.98	0.10-0.15	0.0-2.9	0.0-1.0	.20	.20	5	3	86
	6-60			5-15	1.30-1.60	5.95-19.98	0.02-0.08	0.0-2.9	---	.17	.17			
LNN: Lincoln-----	0-10	86	4	5-15	1.35-1.50	6.00-19.99	0.06-0.11	0.0-2.9	0.5-0.5	.17	.17	5	2	134
	10-60	90	0	5-15	1.30-1.60	6.00-19.99	0.02-0.08	0.0-2.9	---	.17	.17			
Ma: Mangum-----	0-10	22	28	40-60	1.30-1.55	0.00-0.06	0.14-0.18	6.0-8.9	0.0-3.0	.32	.32	5	4	86
	10-50	22	28	40-60	1.30-1.65	0.00-0.06	0.14-0.18	6.0-8.9	---	.32	.32			
	50-60	28	30	30-55	1.40-1.65	0.00-0.06	0.15-0.20	3.0-5.9	---	.32	.32			
Mg: Mangum-----	0-10	22	28	40-60	1.30-1.55	0.00-0.06	0.14-0.18	6.0-8.9	0.0-3.0	.32	.32	5	4	86
	10-50	22	28	40-60	1.30-1.65	0.00-0.06	0.14-0.18	6.0-8.9	---	.32	.32			
	50-60	28	30	30-55	1.40-1.65	0.00-0.06	0.15-0.20	3.0-5.9	---	.32	.32			
Drummond-----	0-8	18	52	27-32	1.30-1.60	0.20-0.60	0.13-0.20	3.0-5.9	0.5-1.0	.43	.43	2	4L	86
	8-30	25	27	35-60	1.40-1.65	0.00-0.06	0.09-0.17	6.0-8.9	---	.55	.55			
	30-60			---	---	---	---	---	---	---	---			
Mn: Minco-----	0-27	14	73	8-18	1.35-1.60	0.60-2.00	0.13-0.24	0.0-2.9	1.0-3.0	.37	.37	5	5	56
	27-60	14	73	8-18	1.35-1.60	0.60-2.00	0.11-0.24	0.0-2.9	---	.37	.37			

PHYSICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Na:														
Naron-----	0-12	63	26	8-14	1.40-1.50	2.00-6.00	0.14-0.18	0.0-2.9	1.0-3.0	.20	.20	5	3	86
	12-37	59	18	18-27	1.45-1.55	0.60-2.00	0.15-0.18	0.0-2.9	---	.32	.32			
	37-60	65	27	2-14	1.50-1.60	2.00-6.00	0.10-0.15	0.0-2.9	---	.32	.32			
Nb:														
Naron-----	0-12	63	26	8-14	1.40-1.50	2.00-6.00	0.14-0.18	0.0-2.9	1.0-3.0	.20	.20	5	3	86
	12-37	59	18	18-27	1.45-1.55	0.60-2.00	0.15-0.18	0.0-2.9	---	.32	.32			
	37-60	65	27	2-14	1.50-1.60	2.00-6.00	0.10-0.15	0.0-2.9	---	.32	.32			
Os:														
Ost-----	0-8	34	37	27-30	1.35-1.40	0.20-0.60	0.15-0.20	0.0-2.9	1.0-3.0	.32	.32	5	6	48
	8-22	35	38	20-34	1.45-1.65	0.20-0.60	0.15-0.20	3.0-5.9	---	.32	.32			
	22-60	39	43	5-30	1.40-1.60	0.20-0.60	0.13-0.20	0.0-2.9	---	.32	.37			
Ot:														
Ost-----	0-8	34	37	27-30	1.35-1.40	0.20-0.60	0.15-0.20	0.0-2.9	1.0-3.0	.32	.32	5	6	48
	8-22	35	38	20-34	1.45-1.65	0.20-0.60	0.15-0.20	3.0-5.9	---	.32	.32			
	22-60	39	43	5-30	1.40-1.60	0.20-0.60	0.13-0.20	0.0-2.9	---	.32	.32			
Pa:														
Pond Creek---	0-11	11	68	15-26	1.30-1.50	0.60-2.00	0.15-0.20	0.0-2.9	1.0-3.0	.37	.37	5	6	48
	11-60	7	65	20-35	1.40-1.70	0.20-0.60	0.15-0.22	3.0-5.9	---	.37	.37			
Pd:														
Pond Creek---	0-11	11	68	15-26	1.30-1.50	0.60-2.00	0.15-0.20	0.0-2.9	1.0-3.0	.37	.37	5	6	48
	11-60	7	65	20-35	1.40-1.70	0.20-0.60	0.15-0.22	3.0-5.9	---	.37	.37			
Ph:														
Dale-----	0-20	11	68	15-26	1.30-1.50	0.60-2.00	0.15-0.24	0.0-2.9	1.0-3.0	.37	.37	5	5	56
	20-60	9	64	18-35	1.40-1.70	0.60-2.00	0.15-0.24	3.0-5.9	---	.37	.37			
Pk:														
Buttermilk---	0-20	11	67	18-26	1.30-1.45	0.60-2.00	0.15-0.24	0.0-2.9	1.0-3.0	.32	.32	5	5	56
	20-32	9	64	18-35	1.35-1.50	0.60-2.00	0.10-0.15	3.0-5.9	---	.37	.37			
	32-60	9	64	18-35	1.35-1.50	0.60-2.00	0.10-0.15	3.0-5.9	---	.37	.37			
Ps:														
Pratt-----	0-7	79	16	2-8	1.40-1.55	5.95-19.98	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	7-30	86	7	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	---	.17	.17			
	30-60	79	16	1-8	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	---	.17	.17			
Pt:														
Pratt-----	0-7	79	16	2-8	1.40-1.55	5.95-19.98	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	7-30	86	7	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	---	.17	.17			
	30-60	79	16	1-8	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	---	.17	.17			
Tivoli-----	0-5	86	7	5-10	1.35-1.50	5.95-19.98	0.07-0.11	0.0-2.9	0.0-1.0	.17	.17	5	2	134
	5-60	93	1	1-10	1.50-1.70	5.95-19.98	0.02-0.08	0.0-2.9	---	.17	.17			
Qn:														
Quinlan-----	0-14	42	37	15-27	1.30-1.55	0.60-2.00	0.13-0.24	0.0-2.9	0.0-1.0	.37	.37	2	4L	86
	>14			---	---	---	---	---	---	---	---			
Qw:														
Quinlan-----	0-14	42	37	15-27	1.30-1.55	0.60-2.00	0.13-0.24	0.0-2.9	0.0-1.0	.37	.37	2	4L	86
	>14			---	---	---	---	---	---	---	---			
Woodward-----	0-40	43	43	10-18	1.30-1.60	0.60-2.00	0.13-0.20	0.0-2.9	0.5-2.0	.37	.37	3	4L	86
	>40			---	---	---	---	---	---	---	---			
Rb:														
Knoco-----	0-9	23	29	40-55	1.35-1.55	0.00-0.06	0.10-0.17	6.0-8.9	0.0-2.0	.37	.37	2	4	86
	9-19	23	29	40-55	1.45-1.70	0.00-0.06	0.07-0.10	6.0-8.9	0.0-1.0	.32	.32			
	19-60	23	29	40-55	1.70-2.25	0.00-0.06	0.00-0.08	6.0-8.9	0.5-0.5	.32	.32			
Rock Outcrop-	0-10			---	---	---	---	---	---	---	---	-	---	0
Sb:														
Shellabarger-	0-14	68	20	8-16	1.35-1.50	0.60-2.00	0.13-0.21	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	14-48	60	18	18-27	1.45-1.60	0.60-2.00	0.16-0.18	0.0-2.9	---	.28	.32			
	48-60	66	24	3-18	1.50-1.65	0.60-2.00	0.05-0.16	0.0-2.9	---	.28	.32			
SBB:														
Shellabarger-	0-11	68	20	8-16	1.35-1.50	0.60-2.00	0.13-0.21	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	11-34	60	18	18-27	1.45-1.60	0.60-2.00	0.16-0.18	0.0-2.9	---	.28	.32			
	34-60	66	24	3-18	1.50-1.65	0.60-2.00	0.05-0.16	0.0-2.9	---	.28	.32			
Sc:														
Shellabarger-	0-14	68	20	8-16	1.35-1.50	0.60-2.00	0.13-0.21	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	14-48	60	18	18-27	1.45-1.60	0.60-2.00	0.16-0.18	0.0-2.9	---	.28	.32			
	48-60	66	24	3-18	1.50-1.65	0.60-2.00	0.05-0.16	0.0-2.9	---	.28	.32			
Tv:														
Tivoli-----	0-5	93	1	1-10	1.35-1.50	5.95-19.98	0.02-0.08	0.0-2.9	0.0-1.0	.17	.17	5	1	250
	5-60	93	1	1-10	1.50-1.70	5.95-19.98	0.02-0.08	0.0-2.9	---	.17	.17			
Vn:														
Vernon-----	0-5	35	34	27-35	1.30-1.60	0.20-0.60	0.15-0.22	3.0-5.9	0.5-1.0	.37	.37	3	4	86
	5-22	23	29	35-60	1.35-1.65	0.06-0.20	0.10-0.20	6.0-8.9	---	.37	.37			
	22-60			---	---	---	---	---	---	---	---			
Vr:														
Vernon-----	0-5	35	34	27-35	1.30-1.60	0.20-0.60	0.15-0.22	3.0-5.9	0.5-1.0	.37	.37	4	4	86
	5-22	23	29	35-60	1.35-1.65	0.06-0.20	0.10-0.20	6.0-8.9	---	.37	.37			
	22-60			---	---	---	---	---	---	---	---			
Vs:														
Knoco-----	0-9	23	29	40-55	1.35-1.55	0.00-0.06	0.10-0.17	6.0-8.9	0.0-2.0	.37	.37	2	4	86
	9-19	23	29	40-55	1.45-1.70	0.00-0.06	0.07-0.10	6.0-8.9	0.0-1.0	.32	.43			
	19-60	23	29	40-55	1.70-2.25	0.00-0.06	0.00-0.08	6.0-8.9	0.5-0.5	.32	.55			
Vernon-----	0-5	35	34	27-35	1.30-1.60	0.20-0.60	0.15-0.22	3.0-5.9	0.5-1.0	.37	.37	4	4	86
	5-22	6	47	35-60	1.35-1.65	0.06-0.20	0.10-0.20	6.0-8.9	---	.37	.37			
	22-60			---	---	---	---	---	---	---	---			
W:														
Water-----	---			---	---	---	---	---	---	---	---	-	---	---

PHYSICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Wa:														
Waldeck-----	0-12	68	20	8-16	1.50-1.60	2.00-6.00	0.14-0.18	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	12-36	68	20	8-16	1.50-1.60	2.00-6.00	0.12-0.17	0.0-2.9	---	.20	.20			
	36-60	96	2	1-4	1.55-1.65	5.95-19.98	0.05-0.07	0.0-2.9	---	.20	.24			
Wo:														
Woodward-----	0-40	43	43	10-18	1.30-1.60	0.60-2.00	0.13-0.20	0.0-2.9	0.5-2.0	.37	.37	3	4L	86
	>40				---	---	---	---	---	---	---			
Quinlan-----	0-14	42	37	15-27	1.30-1.55	0.60-2.00	0.13-0.24	0.0-2.9	0.0-1.0	.37	.37	2	4L	86
	>14				---	---	---	---	---	---	---			
Ws:														
Woodward-----	0-40	43	43	10-18	1.30-1.60	0.60-2.00	0.13-0.20	0.0-2.9	0.5-2.0	.37	.37	3	4L	86
	>40				---	---	---	---	---	---	---			
Quinlan-----	0-14	42	37	15-27	1.30-1.55	0.60-2.00	0.13-0.24	0.0-2.9	0.0-1.0	.37	.37	2	4L	86
	>14				---	---	---	---	---	---	---			
Ya:														
Yahola-----	0-14	67	19	10-18	1.30-1.60	2.00-6.00	0.11-0.15	0.0-2.9	0.5-1.0	.20	.20	5	3	86
	14-60	62	26	5-18	1.40-1.70	2.00-6.00	0.11-0.20	0.0-2.9	---	.32	.32			
	60-66			5-18	1.50-1.70	2.00-6.00	0.07-0.20	0.0-2.9	---	.32	.32			
Ze:														
Zenda-----	0-20	34	37	27-32	1.45-1.55	0.60-2.00	0.17-0.22	3.0-5.9	1.0-3.0	.28	.28	5	6	48
	20-60	35	38	18-35	1.45-1.60	0.60-2.00	0.15-0.19	3.0-5.9	---	.28	.28			

CHEMICAL PROPERTIES OF THE SOILS
Barber County, Kansas

The Chemical Properties table shows estimates of some characteristics and features that affect soil behavior. These estimates are given for the major layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils. Depth to the upper and lower boundaries of each layer is indicated.

Cation-exchange capacity is the total amount of extractable bases that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. Soils having a high cation-exchange capacity can retain cations. The ability to retain cations helps to prevent the pollution of ground water.

Soil reaction is a measure of acidity or alkalinity and is expressed as a range in pH values. The range in pH of each major horizon is based on many field tests. For many soils, values have been verified by laboratory analyses. Soil reaction is important in selecting crops and other plants, in evaluating soil amendments for fertility and stabilization, and in determining the risk of corrosion.

Calcium carbonate equivalent is the percent of carbonates, by weight, in the fraction of the soil less than 2 millimeters in size. The availability of plant nutrients is influenced by the amount of carbonates in the soil. Incorporating nitrogen fertilizer into calcareous soils helps to prevent nitrite accumulation and ammonium-N volatilization.

Gypsum is expressed as a percent, by weight, of hydrated calcium sulfates in the fraction of the soil less than 20 millimeters in size. Gypsum is partially soluble in water and can be dissolved and removed by water. Soils that have a high content of gypsum may collapse if the gypsum is removed by percolating water.

Salinity is a measure of soluble salts in the soil at saturation. It is expressed as the electrical conductivity of the saturation extract, in millimhos per centimeter at 25 degrees C. Estimates are based on field and laboratory measurements at representative sites of nonirrigated soils. The salinity of irrigated soils is affected by the quality of the irrigation water and by the frequency of water application. Hence, the salinity of soils in individual fields can differ greatly from the value given in the table. Salinity affects the suitability of a soil for crop production, the stability of soil if used as construction material, and the potential of the soil to corrode metal and concrete.

Sodium adsorption ratio (SAR) is a measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste. It is the ratio of the Na concentration divided by the square root of one-half of the Ca + Mg concentration. Soils that have SAR values of 13 or more may be characterized by an increased dispersion of organic matter and clay particles, reduced permeability and aeration, and a general degradation of soil structure.

CHEMICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas

PAGE 2 of 8

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
033CP: Clairemont-----	0-15 15-60	6.0-18 7.0-21	7.9-8.4 7.9-8.4	--- ---	--- ---	--- ---	--- ---
033OA: Obaro-----	0-35 >35	10-22 ---	7.9-8.4 ---	0-10 ---	0 ---	--- ---	0 ---
033OB: Obaro-----	0-35 >35	10-22 ---	7.9-8.4 ---	0-10 ---	0 ---	--- ---	0 ---
Rock Outcrop----	---	---	---	---	---	---	---
033QR: Quinlan-----	0-14 >14	6.0-17 ---	7.4-8.4 ---	0-5 ---	0 ---	0 ---	0 ---
Woodward-----	0-30 >30	4.0-12 ---	6.6-8.4 ---	0-5 ---	0 ---	0 ---	0 ---
033QT: Quinlan-----	0-14 >14	6.0-17 ---	7.4-8.4 ---	0-5 ---	0 ---	0 ---	0 ---
Woodward-----	0-30 >30	4.0-12 ---	6.6-8.4 ---	0-5 ---	0 ---	0 ---	0 ---
033YE: Yahola-----	0-12 12-40 40-60	4.0-11 2.0-11 2.0-11	7.4-8.4 7.9-8.4 7.9-8.4	--- --- ---	0 0 0	0 0 0	0 0 0
077BP: Woodward-----	0-24 >24	4.0-12 ---	6.6-8.4 ---	--- ---	0 ---	0 ---	0 ---
Port-----	0-27 27-60	5.0-18 8.0-21	5.6-7.8 6.1-8.4	--- 0	0 0	0 0	0 0
077FU: Farnum-----	0-11 11-41 41-60	6.0-18 10-21 4.0-18	5.6-7.3 6.1-8.4 6.6-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
077GE: Gerlane-----	0-17 17-40 40-48 48-60	4.0-13 4.0-11 4.0-11 8.0-18	6.6-8.4 6.6-8.4 7.4-8.4 7.4-8.4	--- --- --- ---	--- --- --- ---	--- --- --- ---	--- --- --- ---
077MC: Minco-----	0-42 42-60	3.0-13 3.0-11	5.6-8.4 6.1-8.4	0 0	0 0	0 0	0 0
077MN: Minco-----	0-42 42-60	3.0-13 3.0-11	5.6-8.4 6.1-8.4	0 0	0 0	0 0	0 0
077MO: Minco-----	0-42 42-60	3.0-13 3.0-11	5.6-8.4 6.1-8.4	0 0	0 0	0 0	0 0
077QU: Quinlan-----	0-9 >9	6.0-17 ---	7.4-8.4 ---	0-5 ---	0 ---	0 ---	0 ---
077SE: Shellabarger----	0-13 13-38 38-60	3.0-11 7.0-16 1.0-11	5.1-6.5 6.1-7.8 6.1-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
077WE: Woodward-----	0-24 >24	4.0-12 ---	6.6-8.4 ---	0-5 ---	0 ---	0 ---	0 ---
Quinlan-----	0-9 >9	6.0-17 ---	7.4-8.4 ---	0-5 ---	0 ---	0 ---	0 ---
095AC: Albion-----	0-8 8-16 16-26 26-60	3.0-10 4.0-11 1.0-9.0 0.0-6.0	5.6-6.5 6.1-7.8 6.1-8.4 6.1-8.4	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0
095DA: Dillwyn-----	0-8 8-60	0.0-6.0 0.0-5.0	5.6-7.3 5.6-7.8	--- ---	--- ---	--- ---	--- ---
Plevna-----	0-11 11-36 36-60	3.0-13 3.0-11 0.0-4.0	6.6-8.4 6.6-8.4 6.6-8.4	0 0 0	0 0 0	0 0 0	0 0 0

CHEMICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas

PAGE 3 of 8

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
095FA: Farnum-----	0-16 16-50 50-60	3.0-10 10-21 4.0-18	5.6-7.3 6.1-8.4 6.6-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
095PB: Pratt-----	0-12 12-36 36-60	1.0-5.0 1.0-7.0 0.0-5.0	5.6-7.3 5.6-7.3 6.1-7.3	--- --- ---	--- --- ---	--- --- ---	--- --- ---
095SB: Shellabarger----	0-10 10-45 45-60	3.0-11 7.0-16 1.0-11	5.1-6.5 6.1-7.8 6.1-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
095WA: Waldeck-----	0-12 12-36 36-60	3.0-11 3.0-10 0.0-3.0	7.4-8.4 7.4-8.4 7.4-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
097LN: Lincoln-----	0-8 8-60	4.0-11 2.0-9.0	7.4-8.4 7.9-8.4	--- ---	--- ---	--- ---	--- ---
097QW: Quinlan-----	0-15 >15	6.0-17 ---	7.4-8.4 ---	--- ---	0 ---	0 ---	0 ---
Woodward-----	0-27 >27	4.0-12 ---	6.6-8.4 ---	--- ---	0 ---	0 ---	0 ---
097WA: Waldeck-----	0-14 14-41 41-60	4.0-13 3.0-10 0.0-3.0	7.4-8.4 7.4-8.4 7.4-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
151AB: Albion-----	0-8 8-18 18-29 29-60	3.0-10 4.0-11 1.0-9.0 0.0-6.0	5.6-6.5 6.1-7.8 6.1-8.4 6.1-8.4	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0
151AO: Albion-----	0-8 8-18 18-29 29-60	3.0-10 4.0-11 1.0-9.0 0.0-6.0	5.6-6.5 6.1-7.8 6.1-8.4 6.1-8.4	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0
151AS: Albion-----	0-8 8-18 18-29 29-60	3.0-10 4.0-11 1.0-9.0 0.0-6.0	5.6-6.5 6.1-7.8 6.1-8.4 6.1-8.4	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0
Shellabarger----	0-11 11-34 34-60	3.0-11 7.0-16 1.0-11	5.1-6.5 6.1-7.8 6.1-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
151BC: Blanket-----	0-13 13-46 46-60	11-23 14-30 14-30	6.1-7.8 6.1-8.4 7.9-8.4	0 --- ---	0 0 0	0 0 0	0 0 0
151CA: Carwile-----	0-12 12-20 20-33 33-60	2.0-13 10-24 14-36 8.0-27	5.1-7.3 5.1-7.3 6.1-8.4 6.6-8.4	--- --- --- ---	--- --- --- ---	--- --- --- ---	--- --- --- ---
151CK: Case-----	0-6 6-60 0-8 8-60	11-21 7.0-21 11-21 7.0-21	7.4-8.4 7.4-8.4 7.4-8.4 7.4-8.4	0-5 5-25 0-5 15-45	--- --- --- ---	--- --- --- ---	--- --- --- ---
151CM: Clark-----	0-8 8-60	11-21 7.0-21	7.4-8.4 7.4-8.4	0-5 15-45	--- ---	--- ---	--- ---
151CN: Clark-----	0-8 8-60	4.0-13 7.0-21	7.4-8.4 7.4-8.4	0-5 15-45	--- ---	--- ---	--- ---

CHEMICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas

PAGE 4 of 8

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
151CO:							
Clark-----	0-8	11-21	7.4-8.4	0-5	---	---	---
	8-60	7.0-21	7.4-8.4	15-45	---	---	---
Ost-----	0-9	11-20	6.1-8.4	---	---	---	---
	9-14	8.0-21	6.6-8.4	---	---	---	---
	14-23	7.0-21	7.4-8.4	15-34	---	---	---
	23-60	2.0-18	7.4-8.4	15-34	---	---	---
151FA:							
Farnum-----	0-7	11-19	5.6-7.3	---	---	---	---
	7-37	10-21	6.1-8.4	---	---	---	---
	37-60	4.0-18	6.6-8.4	---	---	---	---
151KP:							
Kanza-----	0-11	1.0-9.0	5.6-6.5	---	---	---	---
	11-40	0.0-7.0	5.6-8.4	---	---	---	---
Plevna-----	0-10	3.0-13	6.6-8.4	0	0	0	0
	10-40	3.0-11	6.6-8.4	0	0	0	0
	40-60	0.0-4.0	6.6-8.4	0	0	0	0
151OS:							
Ost-----	0-9	11-20	6.1-8.4	---	---	---	---
	9-14	8.0-21	6.6-8.4	---	---	---	---
	14-23	7.0-21	7.4-8.4	15-34	---	---	---
	23-60	2.0-18	7.4-8.4	15-34	---	---	---
151PM:							
Pratt-----	0-10	1.0-5.0	5.6-7.3	---	---	---	---
	10-40	1.0-7.0	5.6-7.3	---	---	---	---
	40-60	0.0-5.0	6.1-7.3	---	---	---	---
151PN:							
Pratt-----	0-10	1.0-5.0	5.6-7.3	---	---	---	---
	10-40	1.0-7.0	5.6-7.3	---	---	---	---
	40-60	0.0-5.0	6.1-7.3	---	---	---	---
151PO:							
Pratt-----	0-10	1.0-5.0	5.6-7.3	---	---	---	---
	10-40	1.0-7.0	5.6-7.3	---	---	---	---
	40-60	0.0-5.0	6.1-7.3	---	---	---	---
Carwile-----	0-12	2.0-13	5.1-7.3	---	---	---	---
	12-20	10-24	5.1-7.3	---	---	---	---
	20-33	14-36	6.1-8.4	---	---	---	---
	33-60	8.0-27	6.6-8.4	---	---	---	---
151SA:							
Albion-----	0-8	3.0-10	5.6-6.5	0	0	0	0
	8-18	4.0-11	6.1-7.8	0	0	0	0
	18-29	1.0-9.0	6.1-8.4	0	0	0	0
	29-60	0.0-6.0	6.1-8.4	0	0	0	0
Kaski-----	0-26	7.0-18	5.6-7.3	---	---	---	---
	26-40	7.0-21	5.6-7.8	---	---	---	---
	40-60	3.0-18	5.6-8.4	---	---	---	---
151SE:							
Shellabarger----	0-11	3.0-11	5.1-6.5	---	---	---	---
	11-34	7.0-16	6.1-7.8	---	---	---	---
	34-60	1.0-11	6.1-8.4	---	---	---	---
1439:							
Crisfield-----	0-12	4.0-12	5.1-7.3	0	---	0	0
	12-24	4.0-11	5.6-7.3	0	---	0	0
	24-80	1.0-5.0	6.1-7.3	0	---	0	0
Ad:							
Albion-----	0-8	3.0-10	5.6-6.5	0	0	0	0
	8-16	4.0-11	6.1-7.8	0	0	0	0
	16-27	1.0-9.0	6.1-8.4	0	0	0	0
	27-60	0.0-6.0	6.1-8.4	0	0	0	0
Shellabarger----	0-14	3.0-11	5.1-6.5	---	---	---	---
	14-48	7.0-16	6.1-7.8	---	---	---	---
	48-60	1.0-11	6.1-8.4	---	---	---	---
Ae:							
Albion-----	0-8	3.0-10	5.6-6.5	0	0	0	0
	8-16	4.0-11	6.1-7.8	0	0	0	0
	16-27	1.0-9.0	6.1-8.4	0	0	0	0
	27-60	0.0-6.0	6.1-8.4	0	0	0	0
Shellabarger----	0-14	3.0-11	5.1-6.5	---	---	---	---
	14-48	7.0-16	6.1-7.8	---	---	---	---
	48-60	1.0-11	6.1-8.4	---	---	---	---

CHEMICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas

PAGE 5 of 8

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
AED: Arents, Earthen Dam-----	---	---	---	---	---	---	---
AN: Albion-----	0-8 8-16 16-28 28-60	3.0-10 4.0-11 1.0-9.0 0.0-6.0	5.6-6.5 6.1-7.8 6.1-8.4 6.1-8.4	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0
Shellabarger----	0-11 11-38 38-60	3.0-11 7.0-16 1.0-11	5.1-6.5 6.1-7.8 6.1-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
As: Clairemont-----	0-8 8-60	7.0-18 7.0-21	7.9-8.4 7.9-8.4	--- ---	--- ---	4.0-16.0 4.0-16.0	--- ---
At: Attica-----	0-12 12-30 30-60	1.0-7.0 3.0-11 1.0-11	5.6-7.3 5.6-6.5 6.1-7.8	--- --- ---	--- --- ---	--- --- ---	--- --- ---
Ba: Blanket-----	0-9 9-30 30-60	6.0-18 14-30 14-30	6.1-7.8 6.1-8.4 7.9-8.4	0 --- ---	0 0 0	0 0 0	0 0 0
Bb: Blanket-----	0-9 9-30 30-60	6.0-18 14-30 14-30	6.1-7.8 6.1-8.4 7.9-8.4	0 --- ---	0 0 0	0 0 0	0 0 0
Bc: Blanket-----	0-9 9-30 30-60	11-23 14-30 14-30	6.1-7.8 6.1-8.4 7.9-8.4	0 --- ---	0 0 0	0 0 0	0 0 0
Bf: Quinlan-----	0-14 >14	6.0-17 ---	7.4-8.4 ---	--- ---	0 ---	0 ---	0 ---
Clairemont-----	0-14 14-60	6.0-18 7.0-21	7.9-8.4 7.9-8.4	--- ---	--- ---	--- ---	--- ---
Ca: Canadian-----	0-18 18-48 48-60	2.0-13 4.0-11 2.0-11	5.6-7.3 6.1-8.4 6.1-8.4	0 0 0	0 0 0	0 0 0	0 0 0
Cc: Case-----	0-6 6-60	11-21 7.0-21	7.4-8.4 7.4-8.4	0-5 5-25	--- ---	--- ---	--- ---
Clark-----	0-10 10-60	11-21 7.0-21	7.4-8.4 7.4-8.4	0-5 15-45	--- ---	--- ---	--- ---
Cd: Clairemont-----	0-14 14-60	6.0-18 7.0-21	7.9-8.4 7.9-8.4	--- ---	--- ---	--- ---	--- ---
Cf: Clairemont-----	0-14 14-60	6.0-18 7.0-21	7.9-8.4 7.9-8.4	--- ---	--- ---	--- ---	--- ---
Ck: Clark-----	0-10 10-60	11-21 7.0-21	7.4-8.4 7.4-8.4	0-5 15-45	--- ---	--- ---	--- ---
Fa: Farnum-----	0-9 9-60	3.0-10 10-21	5.6-7.3 6.1-8.4	--- ---	--- ---	--- ---	--- ---
Fm: Farnum-----	0-9 9-60	6.0-18 10-21	5.6-7.3 6.1-8.4	--- ---	--- ---	--- ---	--- ---
Fr: Farnum-----	0-9 9-60	6.0-18 10-21	5.6-7.3 6.1-8.4	--- ---	--- ---	--- ---	--- ---
Fu: Farnum-----	0-9 9-60	11-19 10-21	5.6-7.3 6.1-8.4	--- ---	--- ---	--- ---	--- ---
Ga: Grant-----	0-13 13-50 50-60	6.0-18 7.0-21 6.0-16	6.1-7.8 6.1-8.4 7.4-8.4	0 --- ---	0 0 0	0 0 0	0 0 0

CHEMICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas

PAGE 6 of 8

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Gb:							
Grant-----	0-13	6.0-18	6.1-7.8	0	0	0	0
	13-50	7.0-21	6.1-8.4	---	0	0	0
	50-60	6.0-16	7.4-8.4	---	0	0	0
Gc:							
Grant-----	0-13	6.0-18	6.1-7.8	0	0	0	0
	13-50	7.0-21	6.1-8.4	---	0	0	0
	50-60	6.0-16	7.4-8.4	---	0	0	0
GRP:							
Gravel Pits----	---	---	---	---	---	---	---
INT:							
Aquolls-----	0-60	---	---	---	---	---	---
Ka:							
Kanza-----	0-7	1.0-9.0	5.6-6.5	---	---	---	---
	7-48	0.0-7.0	5.6-8.4	---	---	---	---
Kf:							
Kingfisher-----	0-9	6.0-18	6.1-7.8	0	---	0	---
	9-17	10-21	6.1-7.8	0-5	---	0	---
	17-28	10-21	6.6-8.4	---	---	0	---
	>28	---	---	---	---	---	---
Kv:							
Kingfisher-----	0-9	6.0-18	6.1-7.8	0	---	0	---
	9-17	10-21	6.1-7.8	0-5	---	0	---
	17-28	10-21	6.6-8.4	---	---	0	---
	>28	---	---	---	---	---	---
Vernon-----	0-5	11-22	6.6-8.4	0-5	0	0	0
	5-22	14-36	7.9-8.4	0-5	0	0	0
	22-60	---	---	0	0	---	0
Kz:							
Kingfisher-----	0-9	6.0-18	6.1-7.8	0	---	0	---
	9-17	10-21	6.1-7.8	0-5	---	0	---
	17-28	10-21	6.6-8.4	---	---	0	---
	>28	---	---	---	---	---	---
Vernon-----	0-5	11-22	6.6-8.4	0-5	0	0	0
	5-22	14-36	7.9-8.4	0-5	0	0	0
	22-60	---	---	0	0	---	0
Ln:							
Lincoln-----	0-6	4.0-11	7.4-8.4	---	---	---	---
	6-60	2.0-9.0	7.9-8.4	---	---	---	---
LNN:							
Lincoln-----	0-10	2.0-9.0	7.4-8.4	---	---	---	---
	10-60	2.0-9.0	7.9-8.4	---	---	---	---
Ma:							
Mangum-----	0-10	16-38	7.9-8.4	0	0	0.0-2.0	0
	10-50	16-36	7.9-8.4	0-5	0	0.0-4.0	0
	50-60	12-33	7.9-8.4	---	0	0.0-8.0	0
Mg:							
Mangum-----	0-10	16-38	7.9-8.4	0	0	0.0-2.0	0
	10-50	16-36	7.9-8.4	0-5	0	0.0-4.0	0
	50-60	12-33	7.9-8.4	---	0	0.0-8.0	0
Drummond-----	0-8	11-20	6.1-8.4	---	---	0.0-4.0	---
	8-30	14-36	7.4-9.0	---	---	2.0-8.0	---
	30-60	---	---	---	---	---	---
Mn:							
Minco-----	0-27	3.0-13	5.6-8.4	0	0	0	0
	27-60	3.0-11	6.1-8.4	0	0	0	0
Na:							
Naron-----	0-12	3.0-10	5.6-7.3	---	---	---	---
	12-37	7.0-16	5.6-7.8	---	---	---	---
	37-60	0.0-9.0	6.1-8.4	---	---	---	---
Nb:							
Naron-----	0-12	3.0-10	5.6-7.3	---	---	---	---
	12-37	7.0-16	5.6-7.8	---	---	---	---
	37-60	0.0-9.0	6.1-8.4	---	---	---	---
Os:							
Ost-----	0-8	11-20	6.1-8.4	---	---	---	---
	8-22	8.0-21	6.6-8.4	---	---	---	---
	22-60	2.0-18	7.4-8.4	15-34	---	---	---

CHEMICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas

PAGE 7 of 8

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Ot:							
Ost-----	0-8	11-20	6.1-8.4	---	---	---	---
	8-22	8.0-21	6.6-8.4	---	---	---	---
	22-60	2.0-18	7.4-8.4	15-34	---	---	---
Pa:							
Pond Creek-----	0-11	6.0-18	5.1-7.3	---	---	---	---
	11-60	8.0-21	6.1-8.4	---	---	---	---
Pd:							
Pond Creek-----	0-11	6.0-18	5.1-7.3	---	---	---	---
	11-60	8.0-21	6.1-8.4	---	---	---	---
Ph:							
Dale-----	0-20	6.0-18	6.1-7.8	0	0	0	0
	20-60	7.0-21	7.4-8.4	0-5	0	0	0
Pk:							
Buttermilk-----	0-20	7.0-18	7.4-8.4	---	---	2.0-8.0	---
	20-32	7.0-21	7.4-8.4	---	---	4.0-16.0	---
	32-60	7.0-21	7.4-8.4	---	---	2.0-8.0	---
Ps:							
Pratt-----	0-7	1.0-5.0	5.6-7.3	---	---	---	---
	7-30	1.0-7.0	5.6-7.3	---	---	---	---
	30-60	0.0-5.0	6.1-7.3	---	---	---	---
Pt:							
Pratt-----	0-7	1.0-5.0	5.6-7.3	---	---	---	---
	7-30	1.0-7.0	5.6-7.3	---	---	---	---
	30-60	0.0-5.0	6.1-7.3	---	---	---	---
	0-5	2.0-7.0	6.1-7.8	---	---	---	---
	5-60	0.0-6.0	6.1-8.4	---	---	---	---
Qn:							
Quinlan-----	0-14	6.0-17	7.4-8.4	0-5	0	0	0
	>14	---	---	---	---	---	---
Qw:							
Quinlan-----	0-14	6.0-17	7.4-8.4	0-5	0	0	0
	>14	---	---	---	---	---	---
	0-40	4.0-12	6.6-8.4	0-5	0	0	0
	>40	---	---	---	---	---	---
Rb:							
Knoco-----	0-9	16-34	7.9-8.4	3-15	---	0	0-2
	9-19	16-34	7.9-8.4	3-30	0-25	0	2-20
	19-60	16-33	7.9-8.4	1-20	0-25	0	5-26
	0-10	---	---	---	---	---	---
Rock Outcrop----							
Sb:							
Shellabarger----	0-14	3.0-11	5.1-6.5	---	---	---	---
	14-48	7.0-16	6.1-7.8	---	---	---	---
	48-60	1.0-11	6.1-8.4	---	---	---	---
SBB:							
Shellabarger----	0-11	3.0-11	5.1-6.5	---	---	---	---
	11-34	7.0-16	6.1-7.8	---	---	---	---
	34-60	1.0-11	6.1-8.4	---	---	---	---
Sc:							
Shellabarger----	0-14	3.0-11	5.1-6.5	---	---	---	---
	14-48	7.0-16	6.1-7.8	---	---	---	---
	48-60	1.0-11	6.1-8.4	---	---	---	---
Tv:							
Tivoli-----	0-5	0.0-7.0	6.1-7.8	---	---	---	---
	5-60	0.0-6.0	6.1-8.4	---	---	---	---
Vn:							
Vernon-----	0-5	11-22	6.6-8.4	0-5	0	0	0
	5-22	14-36	7.9-8.4	0-5	0	0	0
	22-60	---	---	0	0	---	0
Vr:							
Vernon-----	0-5	11-22	6.6-8.4	0-5	---	---	---
	5-22	14-36	7.9-8.4	0-5	---	---	---
	22-60	---	---	---	---	---	---
Vs:							
Vernon-----	0-5	11-22	6.6-8.4	0-5	---	---	---
	5-22	14-36	7.9-8.4	0-5	---	---	---
	22-60	---	---	---	---	---	---
	0-9	16-34	7.9-8.4	3-15	---	0	0-2
	9-19	16-34	7.9-8.4	3-30	0-25	0	2-20
	19-60	16-33	7.9-8.4	1-20	0-25	0	5-26

CHEMICAL PROPERTIES OF THE SOILS--Continued
Barber County, Kansas

PAGE 8 of 8

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
W:							
Water-----	---	---	---	---	---	---	---
Wa:							
Waldeck-----	0-12	3.0-11	7.4-8.4	---	---	---	---
	12-36	3.0-10	7.4-8.4	---	---	---	---
	36-60	0.0-3.0	7.4-8.4	---	---	---	---
Wo:							
Woodward-----	0-40	4.0-12	6.6-8.4	0-5	0	0	0
	>40	---	---	---	---	---	---
Quinlan-----	0-14	6.0-17	7.4-8.4	0-5	0	0	0
	>14	---	---	---	---	---	---
Ws:							
Woodward-----	0-40	4.0-12	6.6-8.4	0-5	0	0	0
	>40	---	---	---	---	---	---
Quinlan-----	0-14	6.0-17	7.4-8.4	0-5	0	0	0
	>14	---	---	---	---	---	---
Ya:							
Yahola-----	0-14	4.0-11	7.4-8.4	---	---	---	---
	14-60	2.0-11	7.9-8.4	---	0	0	0
	60-66	2.0-11	7.9-8.4	---	0	0	0
Ze:							
Zenda-----	0-20	11-21	6.6-8.4	---	---	0.0-4.0	---
	20-60	7.0-21	7.4-8.4	---	---	0.0-4.0	---

WATER FEATURES Barber County, Kansas

The Water Features table gives estimates of various water features. The estimates are used in land use planning that involves engineering considerations. Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The four hydrologic soil groups are:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

The months in the table indicate the portion of the year in which the feature is most likely to be a concern.

Water table refers to a saturated zone in the soil. The Water Features table indicates, by month, depth to the top (upper limit) and base (lower limit) of the saturated zone in most years. Estimates of the upper and lower limits are based mainly on observations of the water table at selected sites and on evidence of a saturated zone, namely grayish colors or mottles (redoximorphic features) in the soil. A saturated zone that lasts for less than a month is not considered a water table. Ponding is standing water in a closed depression. Unless a drainage system is installed, the water is removed only by percolation, transpiration, or evaporation. The Water Features table indicates surface water depth and the duration and frequency of ponding. Duration is expressed as very brief if less than 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, rare, occasional, and frequent. None means that ponding is not probable; rare that it is unlikely but possible under unusual weather conditions (the chance of ponding is nearly 0 percent to 5 percent in any year); occasional that it occurs, on the average, once or less in 2 years (the chance of ponding is 5 to 50 percent in any year); and frequent that it occurs, on the average, more than once in 2 years (the chance of ponding is more than 50 percent in any year).

Flooding, the temporary inundation of an area, is caused by overflowing streams, by runoff from adjacent slopes, or by tides. Water standing for short periods after rainfall or snowmelt is not considered flooding, and water standing in swamps and marshes is considered ponding rather than flooding.

Duration and frequency are estimated. Duration is expressed as extremely brief if 0.1 hour to 4 hours, very brief if 4 hours to 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, very rare, rare, occasional, frequent, and very frequent. None means that flooding is not probable; very rare that it is very unlikely but possible under extremely unusual weather conditions (the chance of flooding is less than 1 percent in any year); rare that it is unlikely but possible under unusual weather conditions (the chance of flooding is 1 to 5 percent in any year); occasional that it occurs infrequently under normal weather conditions (the chance of flooding is 5 to 50 percent in any year); frequent that it is likely to occur often under normal weather conditions (the chance of flooding is more than 50 percent in any year but is less than 50 percent in all months in any year); and very frequent that it is likely to occur very often under normal weather conditions (the chance of flooding is more than 50 percent in all months of any year).

The information is based on evidence in the soil profile, namely thin strata of gravel, sand, silt, or clay deposited by floodwater; irregular decrease in organic matter content with increasing depth; and little or no horizon development.

Also considered are local information about the extent and levels of flooding and the relation of each soil on the landscape to historic floods. Information on the extent of flooding based on soil data is less specific than that provided by detailed engineering surveys that delineate flood-prone areas at specific flood frequency levels.

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
033CP: Clairemont-----	B	April	---	---	---	---	---	Very brief	Frequent
		May	---	---	---	---	---	Very brief	Frequent
		June	---	---	---	---	---	Very brief	Frequent
		July	---	---	---	---	---	Very brief	Frequent
		August	---	---	---	---	---	Very brief	Frequent
		September	---	---	---	---	---	Very brief	Frequent
		October	---	---	---	---	---	Very brief	Frequent
		November	---	---	---	---	---	Very brief	Frequent
033OA: Obaro-----	B		---	---	---	---	---	---	---
033OB: Obaro-----	B		---	---	---	---	---	---	---
Rock Outcrop-----	D		---	---	---	---	---	---	---
033QR: Quinlan-----	C		---	---	---	---	---	---	---
Woodward-----	B		---	---	---	---	---	---	---
033QT: Quinlan-----	C		---	---	---	---	---	---	---
Woodward-----	B		---	---	---	---	---	---	---
033YE: Yahola-----	B	April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		October	---	---	---	---	---	Very brief	Occasional
077BP: Woodward-----	B		---	---	---	---	---	---	---
Port-----	B	March	---	---	---	---	---	Brief	Frequent
		April	---	---	---	---	---	Brief	Frequent
		May	---	---	---	---	---	Brief	Frequent
		June	---	---	---	---	---	Brief	Frequent
		July	---	---	---	---	---	Brief	Frequent
		August	---	---	---	---	---	Brief	Frequent
077FU: Farnum-----	B		---	---	---	---	---	---	---
077GE: Gerlane-----	B	January	2.0-6.0	>6.0	---	---	---	---	None
		February	2.0-6.0	>6.0	---	---	---	---	None
		March	2.0-6.0	>6.0	---	---	---	Very brief	Occasional
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		December	2.0-6.0	>6.0	---	---	---	---	None
077MC: Minco-----	B		---	---	---	---	---	---	---
077MN: Minco-----	B		---	---	---	---	---	---	---
077MO: Minco-----	B		---	---	---	---	---	---	---
077QU: Quinlan-----	C		---	---	---	---	---	---	---
077SE: Shellabarger-----	B		---	---	---	---	---	---	---
077WE: Quinlan-----	C		---	---	---	---	---	---	---
Woodward-----	B		---	---	---	---	---	---	---
095AC: Albion-----	B		---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
095DA: Dillwyn-----	A		---	---	---	---	---	---	---
		January	1.0-3.0	>6.0	---	---	---	---	None
		February	1.0-3.0	>6.0	---	---	---	---	None
		March	1.0-3.0	>6.0	---	---	---	---	None
		April	1.0-3.0	>6.0	---	---	---	---	None
		May	1.0-3.0	>6.0	---	---	---	---	None
		June	1.0-3.0	>6.0	---	---	---	---	None
		July	1.0-3.0	>6.0	---	---	---	---	None
		August	1.0-3.0	>6.0	---	---	---	---	None
		September	1.0-3.0	>6.0	---	---	---	---	None
		October	1.0-3.0	>6.0	---	---	---	---	None
		November	1.0-3.0	>6.0	---	---	---	---	None
		December	1.0-3.0	>6.0	---	---	---	---	None
Plevna-----	D								
		January	0.0-2.0	>6.0	---	---	---	---	None
		February	0.0-2.0	>6.0	---	---	---	---	None
		March	0.0-2.0	>6.0	---	---	---	Long	Frequent
		April	0.0-2.0	>6.0	---	---	---	Long	Frequent
		May	0.0-2.0	>6.0	---	---	---	Long	Frequent
		June	0.0-2.0	>6.0	---	---	---	Long	Frequent
		July	0.0-2.0	>6.0	---	---	---	Long	Frequent
		August	0.0-2.0	>6.0	---	---	---	Long	Frequent
		September	0.0-2.0	>6.0	---	---	---	Long	Frequent
		October	0.0-2.0	>6.0	---	---	---	Long	Frequent
		November	0.0-2.0	>6.0	---	---	---	---	None
		December	0.0-2.0	>6.0	---	---	---	---	None
095FA: Farnum-----	B								
			---	---	---	---	---	---	---
095PB: Pratt-----	A								
			---	---	---	---	---	---	---
095SB: Shellabarger-----	B								
			---	---	---	---	---	---	---
095WA: Waldeck-----	C								
		January	2.0-4.0	>6.0	---	---	---	---	None
		February	2.0-4.0	>6.0	---	---	---	---	None
		March	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		April	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		May	---	---	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		November	2.0-4.0	>6.0	---	---	---	---	None
		December	2.0-4.0	>6.0	---	---	---	---	None
097LN: Lincoln-----	A								
		January	5.0-6.0	>6.0	---	---	---	---	None
		February	5.0-6.0	>6.0	---	---	---	---	None
		March	5.0-6.0	>6.0	---	---	---	---	None
		April	5.0-6.0	>6.0	---	---	---	Very brief	Occasional
		May	5.0-6.0	>6.0	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		October	---	---	---	---	---	Very brief	Occasional
		November	5.0-6.0	>6.0	---	---	---	---	None
		December	5.0-6.0	>6.0	---	---	---	---	None
097QW: Quinlan-----	C								
			---	---	---	---	---	---	---
Woodward-----	B								
			---	---	---	---	---	---	---
097WA: Waldeck-----	C								
		January	2.0-4.0	>6.0	---	---	---	---	None
		February	2.0-4.0	>6.0	---	---	---	---	None
		March	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		April	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		May	---	---	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		November	2.0-4.0	>6.0	---	---	---	---	None
		December	2.0-4.0	>6.0	---	---	---	---	None

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
151AB: Albion-----	B		Ft	Ft	Ft				
151AO: Albion-----	B		---	---	---	---	---	---	---
151AS: Albion-----	B		---	---	---	---	---	---	---
Shellabarger-----	B		---	---	---	---	---	---	---
151BC: Blanket-----	C		---	---	---	---	---	---	---
151CA: Carwile-----	D		---	---	---	---	---	---	---
		January	0.0	>6.0	0.0-1.0	Brief	---	---	None
		February	0.0	>6.0	0.0-1.0	Brief	---	---	None
		March	0.0	>6.0	0.0-1.0	Brief	---	---	None
		April	0.0	>6.0	0.0-1.0	Brief	---	---	None
		May	---	---	0.0-	---	---	---	None
		June	---	---	0.0-	---	---	---	None
		July	---	---	0.0-	---	---	---	None
		August	---	---	0.0-	---	---	---	None
		September	---	---	0.0-	---	---	---	None
		October	0.0	>6.0	0.0-1.0	Brief	---	---	None
		November	0.0	>6.0	0.0-1.0	Brief	---	---	None
		December	0.0	>6.0	0.0-1.0	Brief	---	---	None
151CK: Case-----	B		---	---	---	---	---	---	---
Clark-----	B		---	---	---	---	---	---	---
151CM: Clark-----	B		---	---	---	---	---	---	---
151CN: Clark-----	B		---	---	---	---	---	---	---
151CO: Clark-----	B		---	---	---	---	---	---	---
Ost-----	B		---	---	---	---	---	---	---
151FA: Farnum-----	B		---	---	---	---	---	---	---
151KP: Kanza-----	D		---	---	---	---	---	---	---
		January	0.0-3.0	>6.0	---	---	---	Very brief	Frequent
		February	0.0-3.0	>6.0	---	---	---	Very brief	Frequent
		March	0.0-3.0	>6.0	---	---	---	Very brief	Frequent
		April	---	---	---	---	---	Very brief	Frequent
		May	---	---	---	---	---	Very brief	Frequent
		June	---	---	---	---	---	Very brief	Frequent
		July	---	---	---	---	---	Very brief	Frequent
		August	---	---	---	---	---	Very brief	Frequent
		September	---	---	---	---	---	Very brief	Frequent
		October	---	---	---	---	---	Very brief	Frequent
		November	---	---	---	---	---	Very brief	Frequent
		December	0.0-3.0	>6.0	---	---	---	Very brief	Frequent
Plevna-----	D		---	---	---	---	---	---	---
		January	0.0-2.0	>6.0	---	---	---	---	None
		February	0.0-2.0	>6.0	---	---	---	---	None
		March	0.0-2.0	>6.0	---	---	---	Long	Frequent
		April	0.0-2.0	>6.0	---	---	---	Long	Frequent
		May	0.0-2.0	>6.0	---	---	---	Long	Frequent
		June	0.0-2.0	>6.0	---	---	---	Long	Frequent
		July	0.0-2.0	>6.0	---	---	---	Long	Frequent
		August	0.0-2.0	>6.0	---	---	---	Long	Frequent
		September	0.0-2.0	>6.0	---	---	---	Long	Frequent
		October	0.0-2.0	>6.0	---	---	---	Long	Frequent
		November	0.0-2.0	>6.0	---	---	---	---	None
		December	0.0-2.0	>6.0	---	---	---	---	None
151OS: Ost-----	B		---	---	---	---	---	---	---
151PM: Pratt-----	A		---	---	---	---	---	---	---
151PN: Pratt-----	A		---	---	---	---	---	---	---
151PO:			---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Pratt-----	A		Ft	Ft	Ft				
Carwile-----	D		---	---	---	---	---	---	---
		January	0.0	>6.0	0.0-1.0	Long	---	---	None
		February	0.0	>6.0	0.0-1.0	Long	---	---	None
		March	0.0	>6.0	0.0-1.0	Long	---	---	None
		April	0.0	>6.0	0.0-1.0	Long	---	---	None
		May	---	---	0.0-	---	---	---	None
		June	---	---	0.0-	---	---	---	None
		July	---	---	0.0-	---	---	---	None
		August	---	---	0.0-	---	---	---	None
		September	---	---	0.0-	---	---	---	None
		October	0.0	>6.0	0.0-1.0	Long	---	---	None
		November	0.0	>6.0	0.0-1.0	Long	---	---	None
		December	0.0	>6.0	0.0-1.0	Long	---	---	None
151SA: Albion-----	B		---	---	---	---	---	---	---
Kaski-----	B		---	---	---	---	---	---	---
		March	---	---	---	---	---	Very brief	Occasional
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
151SE: Shellabarger-----	B		---	---	---	---	---	---	---
1439: Crisfield-----	B		---	---	---	---	---	---	---
		March	3.3-6.1	>6.0	---	---	---	---	Rare
		April	3.3-6.1	>6.0	---	---	---	---	Rare
		May	3.3-6.1	>6.0	---	---	---	---	Rare
		June	3.3-6.1	>6.0	---	---	---	---	Rare
		July	3.3-6.1	>6.0	---	---	---	---	None
Ad: Albion-----	B		---	---	---	---	---	---	---
Shellabarger-----	B		---	---	---	---	---	---	---
Ae: Albion-----	B		---	---	---	---	---	---	---
Shellabarger-----	B		---	---	---	---	---	---	---
AN: Albion-----	B		---	---	---	---	---	---	---
Shellabarger-----	B		---	---	---	---	---	---	---
As: Clairemont-----	B		---	---	---	---	---	---	---
		April	---	---	---	---	---	Very brief	Frequent
		May	---	---	---	---	---	Very brief	Frequent
		June	---	---	---	---	---	Very brief	Frequent
		July	---	---	---	---	---	Very brief	Frequent
		August	---	---	---	---	---	Very brief	Frequent
		September	---	---	---	---	---	Very brief	Frequent
		October	---	---	---	---	---	Very brief	Frequent
		November	---	---	---	---	---	Very brief	Frequent
At: Attica-----	B		---	---	---	---	---	---	---
Ba: Blanket-----	C		---	---	---	---	---	---	---
Bb: Blanket-----	C		---	---	---	---	---	---	---
Bc: Blanket-----	C		---	---	---	---	---	---	---
Bf: Quinlan-----	C		---	---	---	---	---	---	---
Clairemont-----	B		---	---	---	---	---	---	---
		April	---	---	---	---	---	Very brief	Frequent
		May	---	---	---	---	---	Very brief	Frequent
		June	---	---	---	---	---	Very brief	Frequent
		July	---	---	---	---	---	Very brief	Frequent
		August	---	---	---	---	---	Very brief	Frequent
		September	---	---	---	---	---	Very brief	Frequent
		October	---	---	---	---	---	Very brief	Frequent
		November	---	---	---	---	---	Very brief	Frequent

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Ca: Canadian-----	B	March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
Cc: Case-----	B		---	---	---	---	---	---	---
Clark-----	B		---	---	---	---	---	---	---
Cd: Clairemont-----	B		---	---	---	---	---	---	---
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		October	---	---	---	---	---	Very brief	Occasional
		November	---	---	---	---	---	Very brief	Occasional
Cf: Clairemont-----	B	April	---	---	---	---	---	Very brief	Frequent
		May	---	---	---	---	---	Very brief	Frequent
		June	---	---	---	---	---	Very brief	Frequent
		July	---	---	---	---	---	Very brief	Frequent
		August	---	---	---	---	---	Very brief	Frequent
		September	---	---	---	---	---	Very brief	Frequent
		October	---	---	---	---	---	Very brief	Frequent
		November	---	---	---	---	---	Very brief	Frequent
Ck: Clark-----	B		---	---	---	---	---	---	---
Fa: Farnum-----	B		---	---	---	---	---	---	---
Fm: Farnum-----	B		---	---	---	---	---	---	---
Fr: Farnum-----	B		---	---	---	---	---	---	---
Fu: Farnum-----	B		---	---	---	---	---	---	---
Ga: Grant-----	B		---	---	---	---	---	---	---
Gb: Grant-----	B		---	---	---	---	---	---	---
Gc: Grant-----	B		---	---	---	---	---	---	---
GRP: Gravel Pits-----	---		---	---	---	---	---	---	---
INT: Aquolls-----	C	February		---	---	---	None	---	None
		March		---	---	---	None	---	None
		April		---	0.0-0.8	Brief	Occasional	---	None
		May	0.0 0.5-1.5 3.0-5.0	3.0-5.0 >6.0 0.5-1.5	0.0-0.8	Brief	Occasional	---	None
		June	0.0 0.5-1.5 3.0-5.0	3.0-5.0 >6.0 0.5-1.5	0.0-0.8	Brief	Occasional	---	None
		November	---	---	---	---	None	---	None
Ka:									

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Kanza-----	D		Ft	Ft	Ft				
		January	0.0-3.0	>6.0	---	---	---	Very brief	Frequent
		February	0.0-3.0	>6.0	---	---	---	Very brief	Frequent
		March	0.0-3.0	>6.0	---	---	---	Very brief	Frequent
		April	---	---	---	---	---	Very brief	Frequent
		May	---	---	---	---	---	Very brief	Frequent
		June	---	---	---	---	---	Very brief	Frequent
		July	---	---	---	---	---	Very brief	Frequent
		August	---	---	---	---	---	Very brief	Frequent
		September	---	---	---	---	---	Very brief	Frequent
		October	---	---	---	---	---	Very brief	Frequent
		November	---	---	---	---	---	Very brief	Frequent
		December	0.0-3.0	>6.0	---	---	---	Very brief	Frequent
Kf: Kingfisher-----	B		---	---	---	---	---	---	---
Kv: Kingfisher-----	B		---	---	---	---	---	---	---
Vernon-----	D		---	---	---	---	---	---	---
Kz: Kingfisher-----	B		---	---	---	---	---	---	---
Vernon-----	D		---	---	---	---	---	---	---
Ln: Lincoln-----	A								
		January	5.0-6.0	>6.0	---	---	---	---	None
		February	5.0-6.0	>6.0	---	---	---	---	None
		March	5.0-6.0	>6.0	---	---	---	---	None
		April	5.0-6.0	>6.0	---	---	---	Brief	Frequent
		May	5.0-6.0	>6.0	---	---	---	Brief	Frequent
		June	---	---	---	---	---	Brief	Frequent
		July	---	---	---	---	---	Brief	Frequent
		August	---	---	---	---	---	Brief	Frequent
		September	---	---	---	---	---	Brief	Frequent
		October	---	---	---	---	---	Brief	Frequent
		November	5.0-6.0	>6.0	---	---	---	---	None
		December	5.0-6.0	>6.0	---	---	---	---	None
LNN: Lincoln-----	A								
		January	5.0-6.0	>6.0	---	---	---	---	None
		February	5.0-6.0	>6.0	---	---	---	---	None
		March	5.0-6.0	>6.0	---	---	---	---	None
		April	5.0-6.0	>6.0	---	---	---	Very brief	Occasional
		May	5.0-6.0	>6.0	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		October	---	---	---	---	---	Very brief	Occasional
		November	5.0-6.0	>6.0	---	---	---	---	None
		December	5.0-6.0	>6.0	---	---	---	---	None
Ma: Mangum-----	D								
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		October	---	---	---	---	---	Very brief	Occasional
		November	---	---	---	---	---	Very brief	Occasional
		December	5.0-6.0	>6.0	---	---	---	---	None
			5.0-6.0	>6.0	---	---	---	---	None
Mg: Mangum-----	D								
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
Drummond-----	D								
		January	2.0-6.0	>6.0	---	---	---	---	None
		February	2.0-6.0	>6.0	---	---	---	---	None
		March	2.0-6.0	>6.0	---	---	---	---	Rare
		April	2.0-6.0	>6.0	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		November	2.0-6.0	>6.0	---	---	---	---	None
		December	2.0-6.0	>6.0	---	---	---	---	None
Mn: Minco-----	B		---	---	---	---	---	---	---
Na:									

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Naron-----	B		Ft	Ft	Ft				
Nb: Naron-----	B		---	---	---	---	---	---	---
Os: Ost-----	B		---	---	---	---	---	---	---
Ot: Ost-----	B		---	---	---	---	---	---	---
Pa: Pond Creek-----	B		---	---	---	---	---	---	---
Pd: Pond Creek-----	B		---	---	---	---	---	---	---
Ph: Dale-----	B	March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
Pk: Buttermilk-----	B	January	4.5-6.0	>6.0	---	---	---	---	None
		February	4.5-6.0	>6.0	---	---	---	---	None
		March	4.5-6.0	>6.0	---	---	---	---	Rare
		April	4.5-6.0	>6.0	---	---	---	---	Rare
		May	4.5-6.0	>6.0	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		November	4.5-6.0	>6.0	---	---	---	---	None
		December	4.5-6.0	>6.0	---	---	---	---	None
Ps: Pratt-----	A		---	---	---	---	---	---	---
Pt: Pratt-----	A		---	---	---	---	---	---	---
Tivoli-----	A		---	---	---	---	---	---	---
Qn: Quinlan-----	C		---	---	---	---	---	---	---
Qw: Quinlan-----	C		---	---	---	---	---	---	---
Woodward-----	B		---	---	---	---	---	---	---
Rb: Knoco-----	D		---	---	---	---	---	---	---
Rock Outcrop-----	D		---	---	---	---	---	---	---
Sb: Shellabarger-----	B		---	---	---	---	---	---	---
SBB: Shellabarger-----	B		---	---	---	---	---	---	---
Sc: Shellabarger-----	B		---	---	---	---	---	---	---
Tv: Tivoli-----	A		---	---	---	---	---	---	---
Vn: Vernon-----	D		---	---	---	---	---	---	---
Vr: Vernon-----	D		---	---	---	---	---	---	---
Vs: Knoco-----	D		---	---	---	---	---	---	---
Vernon-----	D		---	---	---	---	---	---	---
W: Water-----	---		---	---	---	---	---	---	---
Wa:			---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Waldeck-----	C		Ft	Ft	Ft				
		January	2.0-4.0	>6.0	---	---	---	---	None
		February	2.0-4.0	>6.0	---	---	---	---	None
		March	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		April	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		May	---	---	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		November	2.0-4.0	>6.0	---	---	---	---	None
		December	2.0-4.0	>6.0	---	---	---	---	None
Wo: Woodward-----	B		---	---	---	---	---	---	---
Quinlan-----	C		---	---	---	---	---	---	---
Ws: Woodward-----	B		---	---	---	---	---	---	---
Quinlan-----	C		---	---	---	---	---	---	---
Ya: Yahola-----	B		---	---	---	---	---	---	---
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		October	---	---	---	---	---	Very brief	Occasional
Ze: Zenda-----	C		---	---	---	---	---	---	---
		January	2.0-4.0	>6.0	---	---	---	---	None
		February	2.0-4.0	>6.0	---	---	---	---	None
		March	2.0-4.0	>6.0	---	---	---	---	None
		April	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		October	2.0-4.0	>6.0	---	---	---	---	None
		November	2.0-4.0	>6.0	---	---	---	---	None
		December	2.0-4.0	>6.0	---	---	---	---	None

The following table gives estimates of various soil features. The estimates are used in land use planning that involves engineering considerations.

A restrictive layer is a nearly continuous layer that has one or more physical, chemical, or thermal properties that significantly impede the movement of water and air through the soil or that restrict roots or otherwise provide an unfavorable root environment. Examples are bedrock, cemented layers, dense layers, and frozen layers. The table indicates the hardness and thickness of the restrictive layer, both of which significantly affect the ease of excavation. Depth to top is the vertical distance from the soil surface to the upper boundary of the restrictive layer.

Potential for frost action is the likelihood of upward or lateral expansion of the soil caused by the formation of segregated ice lenses (frost heave) and the subsequent collapse of the soil and loss of strength on thawing. Frost action occurs when moisture moves into the freezing zone of the soil. Temperature, texture, density, permeability, content of organic matter, and depth to the water table are the most important factors considered in evaluating the potential for frost action. It is assumed that the soil is not insulated by vegetation or snow and is not artificially drained. Silty and highly structured, clayey soils that have a high water table in winter are the most susceptible to frost action. Well drained, very gravelly, or very sandy soils are the least susceptible. Frost heave and low soil strength during thawing cause damage to pavements and other rigid structures.

Risk of corrosion pertains to potential soil-induced electrochemical or chemical action that corrodes or weakens uncoated steel or concrete. The rate of corrosion of uncoated steel is related to such factors as soil moisture, particle-size distribution, acidity, and electrical conductivity of the soil. The rate of corrosion of concrete is based mainly on the sulfate and sodium content, texture, moisture content, and acidity of the soil. Special site examination and design may be needed if the combination of factors results in a severe hazard of corrosion. The steel or concrete in installations that intersect soil boundaries or soil layers is more susceptible to corrosion than the steel or concrete in installations that are entirely within one kind of soil or within one soil layer.

For uncoated steel, the risk of corrosion, expressed as low, moderate, or high, is based on soil drainage class, total acidity, electrical resistivity near field capacity, and electrical conductivity of the saturation extract.

For concrete, the risk of corrosion also is expressed as low, moderate, or high. It is based on soil texture, acidity, and amount of sulfates in the saturation extract.

SOIL FEATURES--Continued
Barber County, Kansas

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness		Uncoated Steel	Concrete
033CP: Clairemont-----	---	In	In	---	None	Moderate	Low
033OA: Obaro-----	20-40	Bedrock (paralithic)	---	Weakly cemented	None	Low	Low
033OB: Obaro-----	20-40	Bedrock (paralithic)	---	Weakly cemented	None	Low	Low
Rock Outcrop----	---	---	---	---	None	---	---
033QR: Quinlan-----	10-20	Bedrock (paralithic)	---	Extremely weakly cemented	None	Moderate	Low
Woodward-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	Low	Low
033QT: Quinlan-----	10-20	Bedrock (paralithic)	---	Extremely weakly cemented	None	Moderate	Low
Woodward-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	Low	Low
033YE: Yahola-----	---	---	---	---	None	Low	Low
077BP: Woodward-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	Low	Low
Port-----	---	---	---	---	None	Moderate	Low
077FU: Farnum-----	---	---	---	---	None	Moderate	Low
077GE: Gerlane-----	---	---	---	---	None	Low	Low
077MC: Minco-----	---	---	---	---	None	Low	Low
077MN: Minco-----	---	---	---	---	None	Low	Low
077MO: Minco-----	---	---	---	---	None	Low	Low
077QU: Quinlan-----	10-20	Bedrock (paralithic)	---	Extremely weakly cemented	None	Moderate	Low
077SE: Shellabarger----	---	---	---	---	None	Low	Moderate
077WE: Quinlan-----	10-20	Bedrock (paralithic)	---	Extremely weakly cemented	None	Moderate	Low
Woodward-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	Low	Low
095AC: Albion-----	---	---	---	---	None	Low	Low
095DA: Dillwyn-----	---	---	---	---	Low	Low	Low
Plevna-----	---	---	---	---	Low	High	Low
095FA: Farnum-----	---	---	---	---	Low	Moderate	Low
095PB: Pratt-----	---	---	---	---	Low	Low	Moderate
095SB: Shellabarger----	---	---	---	---	None	Low	Moderate
095WA: Waldeck-----	---	---	---	---	Low	Moderate	Low
097LN: Lincoln-----	---	---	---	---	Low	Low	Low
097QW: Quinlan-----	10-20	Bedrock (paralithic)	---	Extremely weakly cemented	Low	Moderate	Low
Woodward-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	Low	Low	Low
097WA: Waldeck-----	---	---	---	---	Low	Moderate	Low
151AB: Albion-----	---	---	---	---	Low	Low	Low
151AO: Albion-----	---	---	---	---	Low	Low	Low
151AS: Albion-----	---	---	---	---	Low	Low	Low
151AS: Shellabarger----	---	---	---	---	Low	Low	Low
151BC: Blanket-----	---	---	---	---	Low	High	Low
151CA: Carwile-----	---	---	---	---	---	High	Moderate
151CK: Case-----	---	---	---	---	Low	Moderate	Low
Clark-----	---	---	---	---	Low	Moderate	Low
151CM: Clark-----	---	---	---	---	Low	Moderate	Low
151CN: Clark-----	---	---	---	---	Low	Moderate	Low

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top In	Thickness In	Hardness		Uncoated Steel	Concrete
151CO:							
Clark-----	---	---	---	---	Low	Moderate	Low
Ost-----	---	---	---	---	Low	Moderate	Low
151FA:							
Farnum-----	---	---	---	---	Low	Moderate	Low
151KP:							
Kanza-----	---	---	---	---	Low	High	Moderate
Plevna-----	---	---	---	---	Low	High	Low
151OS:							
Ost-----	---	---	---	---	Low	Moderate	Low
151PM:							
Pratt-----	---	---	---	---	Low	Low	Moderate
151PN:							
Pratt-----	---	---	---	---	Low	Low	Moderate
151PO:							
Pratt-----	---	---	---	---	Low	Low	Moderate
Carwile-----	---	---	---	---	Low	High	Moderate
151SA:							
Albion-----	---	---	---	---	Low	Low	Low
Kaski-----	---	---	---	---	Low	Low	Low
151SE:							
Shellabarger----	---	---	---	---	Low	Low	Moderate
1439:							
Crisfield-----	---	---	---	---	None	Low	Low
Ad:							
Albion-----	---	---	---	---	None	Low	Low
Shellabarger----	---	---	---	---	None	Low	Moderate
Ae:							
Albion-----	---	---	---	---	None	Low	Low
Shellabarger----	---	---	---	---	None	Low	Moderate
AED:							
Arents, Earthen Dam-----	---	---	---	---	---	---	---
AN:							
Albion-----	---	---	---	---	None	Low	Low
Shellabarger----	---	---	---	---	None	Low	Moderate
As:							
Clairemont-----	---	---	---	---	None	High	Low
At:							
Attica-----	---	---	---	---	None	Low	Low
Ba:							
Blanket-----	---	---	---	---	None	High	Low
Bb:							
Blanket-----	---	---	---	---	None	High	Low
Bc:							
Blanket-----	---	---	---	---	None	High	Low
Bf:							
Quinlan-----	10-20	Bedrock (paralithic)	---	Extremely weakly cemented	None	Moderate	Low
Clairemont-----	---	---	---	---	None	Moderate	Low
Ca:							
Canadian-----	---	---	---	---	None	Low	Low
Cc:							
Case-----	---	---	---	---	None	Moderate	Low
Clark-----	---	---	---	---	None	Moderate	Low
Cd:							
Clairemont-----	---	---	---	---	None	Moderate	Low
Cf:							
Clairemont-----	---	---	---	---	None	Moderate	Low
Ck:							
Clark-----	---	---	---	---	None	Moderate	Low
Fa:							
Farnum-----	---	---	---	---	None	Moderate	Low
Fm:							
Farnum-----	---	---	---	---	None	Moderate	Low
Fr:							
Farnum-----	---	---	---	---	None	Moderate	Low
Fu:							
Farnum-----	---	---	---	---	None	Moderate	Low
Ga:							
Grant-----	---	---	---	---	None	Moderate	Low
Gb:							
Grant-----	---	---	---	---	None	Moderate	Low
Gc:							
Grant-----	---	---	---	---	None	Moderate	Low
GRP:							
Gravel Pits-----	---	---	---	---	None	---	---
INT:							
Aquolls-----	---	---	---	---	Moderate	---	---
Ka:							
Kanza-----	---	---	---	---	None	High	Moderate
Kf:							
Kingfisher-----	20-40	Bedrock (paralithic)	---	Very weakly cemented	None	Moderate	Low

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness		Uncoated Steel	Concrete
Kv:		In	In				
Kingfisher-----	20-40	Bedrock (paralithic)	---	Very weakly cemented	None	Moderate	Low
Vernon-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	High	Low
Kz:							
Kingfisher-----	20-40	Bedrock (paralithic)	---	Very weakly cemented	None	Moderate	Low
Vernon-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	High	Low
Ln:							
Lincoln-----	---	---	---	---	None	Low	Low
LNN:							
Lincoln-----	---	---	---	---	None	Low	Low
Ma:							
Mangum-----	---	---	---	---	None	High	Low
Mg:							
Mangum-----	---	---	---	---	None	High	Low
Drummond-----	---	---	---	---	None	High	High
Mn:							
Minco-----	---	---	---	---	None	Low	Low
Na:							
Naron-----	---	---	---	---	None	Low	Low
Nb:							
Naron-----	---	---	---	---	None	Low	Low
Os:							
Ost-----	---	---	---	---	None	Moderate	Low
Ot:							
Ost-----	---	---	---	---	None	Moderate	Low
Pa:							
Pond Creek-----	---	---	---	---	None	Moderate	Moderate
Pd:							
Pond Creek-----	---	---	---	---	None	Moderate	Moderate
Ph:							
Dale-----	---	---	---	---	None	Moderate	Low
Pk:							
Buttermilk-----	---	---	---	---	None	High	Moderate
Ps:							
Pratt-----	---	---	---	---	None	Low	Moderate
Pt:							
Pratt-----	---	---	---	---	None	Low	Moderate
Tivoli-----	---	---	---	---	None	Low	Low
Qn:							
Quinlan-----	10-20	Bedrock (paralithic)	---	Extremely weakly cemented	None	Moderate	Low
Qw:							
Quinlan-----	10-20	Bedrock (paralithic)	---	Extremely weakly cemented	None	Moderate	Low
Woodward-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	Low	Low
Rb:							
Knoco-----	4-19	Bedrock (paralithic)	---	Extremely weakly cemented	None	High	Low
Rock Outcrop----	0-0	Bedrock (paralithic)	---	---	None	---	---
Sb:							
Shellabarger----	---	---	---	---	None	Low	Moderate
SBB:							
Shellabarger----	---	---	---	---	Low	Low	Moderate
Sc:							
Shellabarger----	---	---	---	---	None	Low	Moderate
Tv:							
Tivoli-----	---	---	---	---	None	Low	Low
Vn:							
Vernon-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	High	Low
Vr:							
Vernon-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	High	Low
Vs:							
Knoco-----	4-19	Bedrock (paralithic)	---	Extremely weakly cemented	None	High	Low
Vernon-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	High	Low
W:							
Water-----	---	---	---	---	None	---	Low
Wa:							
Waldeck-----	---	---	---	---	None	Moderate	Low
Wo:							
Woodward-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	Low	Low
Quinlan-----	10-20	Bedrock (paralithic)	---	Extremely weakly cemented	None	Moderate	Low

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness		Uncoated Steel	Concrete
Ws:		In	In				
Woodward-----	20-40	Bedrock (paralithic)	---	Extremely weakly cemented	None	Low	Low
Quinlan-----	10-20	Bedrock (paralithic)	---	Extremely weakly cemented	None	Moderate	Low
Ya:							
Yahola-----	---	---	---	---	None	Low	Low
Ze:							
Zenda-----	---	---	---	---	None	High	Low

WATER MANAGEMENT
Barber County, Kansas

The soils of the survey area are rated in the Water Management table according to limitations that affect their suitability for water management. Soils are rated for pond reservoir areas, drainage, irrigation, terraces and diversions, and grassed waterways. Restrictive features that affect each soil for the specified use is also provided in the table.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the specified use. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Moderately limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Limited indicates that the soil has one or more features that are significant limitations for the specified use. The limitations can be overcome, but generally require special design, soil reclamation, or installation procedures that may result in additional expense. Fair performance and moderate to high maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Limitation class terms, such as very limited or limited, etc., limitation ratings, and numerical ratings are shown for each soil feature listed. As many as three soil features may be listed for each soil component if applicable. The overall limitation rating for the soil component is based on the most severe limitation.

Pond reservoir areas hold water behind a dam or embankment. Soils best suited to this use have low seepage potential in the upper 60 inches. The seepage potential is determined by the permeability of the soil and the depth to fractured bedrock or other permeable material. Excessive slope can affect the storage capacity of the reservoir area.

Embankments, dikes, and levees are raised structures of soil material, generally less than 20 feet high, constructed to impound water or to protect land against overflow. In this table, the soils are rated as a source of material for embankment fill. The ratings apply to the soil material below the surface layer to a depth of about 5 feet. It is assumed that soil layers will be uniformly mixed and compacted during construction.

The ratings do not indicate the ability of the natural soil to support an embankment. Soil properties to a depth even greater than the height of the embankment can affect performance and safety of the embankment. Generally, deeper onsite investigation is needed to determine these properties.

Soil material in embankments must be resistant to seepage, piping, and erosion and have favorable compaction characteristics. Unfavorable features include less than 5 feet of suitable material and a high content of stones or boulders, organic matter, or salts or sodium. A high water table affects the amount of usable material. It also affects traffic ability.

Aquifer-fed excavated ponds are pits or dugouts that extend to a ground-water aquifer or to a depth below a permanent water table. Excluded are ponds that are fed only by surface runoff and embankment ponds that impound water 3 feet or more above the original surface. Excavated ponds are affected by depth to a permanent water table, permeability of the aquifer, and quality of the water as inferred from the salinity of the soil. Depth to bedrock and the content of large stones affect the ease of excavation.

Drainage is the removal of excess surface and subsurface water from the soil. How easily and effectively the soil is drained depends on the depth to bedrock, to a cemented pan, or to other layers that affect the rate of water movement; permeability; depth to a high water table or depth of standing water if the soil is subject to ponding; slope; susceptibility to flooding; subsidence of organic layers; and the potential for frost action. Excavating and grading and the stability of ditch banks are affected by depth to bedrock or to a cemented pan, large stones, slope, and the hazard of cutbanks caving. The productivity of the soil after drainage is adversely affected by extreme acidity or by toxic substances in the root zone, such as salts, sodium, and sulfur. Availability of drainage outlets is not considered in the ratings.

Irrigation is the controlled application of water to supplement rainfall and support plant growth. The design and management of an irrigation system are affected by depth to the water table, the need for drainage, flooding, available water capacity, intake rate, permeability, erosion hazard, and slope. The construction of a system is affected by large stones and depth to bedrock or to a cemented pan. The performance of a system is affected by the depth of the root zone, the amount of salts or sodium, and soil reaction.

Terraces and diversions are embankments or a combination of channels and ridges constructed across a slope to control erosion and conserve moisture by intercepting runoff. Slope, wetness, large stones, and depth to bedrock or to a cemented pan affect the construction of terraces and diversions. A restricted rooting depth, a very limited hazard of wind erosion or water erosion, an excessively coarse texture, and restricted permeability adversely affect maintenance.

Grassed waterways are natural or constructed channels, generally broad and shallow, which conduct surface water to outlets at a non-erosive velocity. Large stones, wetness, slope, and depth to bedrock or to a cemented pan affect the construction of grassed waterways. A hazard of wind erosion, low available water capacity, restricted rooting depth, toxic substances such as salts and sodium, and restricted permeability adversely affect the growth and maintenance of the grass after construction.

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
033CP: Clairemont-----	Limitation: deep to water	Limitation: erodes easily flooding	Limitation: erodes easily	Limitation: erodes easily
033OA: Obaro-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
033OB: Obaro-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
Rock Outcrop----	---	---	---	---
033QR: Quinlan-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
Woodward-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
033QT: Quinlan-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
Woodward-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
033YE: Yahola-----	Limitation: deep to water	Limitation: flooding soil blowing	Limitation: soil blowing	Favorable
077BP: Woodward-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
Port-----	Limitation: deep to water	Limitation: erodes easily flooding	Limitation: erodes easily	Limitation: erodes easily
077FU: Farnum-----	Limitation: deep to water	Limitation: slope	Favorable	Favorable
077GE: Gerlane-----	Limitation: flooding	Limitation: flooding wetness soil blowing	Limitation: wetness soil blowing	Favorable
077MC: Minco-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
077MN: Minco-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
077MO: Minco-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily	Limitation: erodes easily
077QU: Quinlan-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock
077SE: Shellabarger----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
077WE: Quinlan-----	Limitation: deep to water	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock
Woodward-----	Limitation: deep to water	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock
095AC: Albion-----	Limitation: deep to water	Limitation: slope soil blowing droughty	Limitation: too sandy soil blowing	Limitation: droughty

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
095DA: Dillwyn-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: wetness droughty
Plevna-----	Limitation: flooding	Limitation: flooding wetness soil blowing	Limitation: wetness soil blowing	Limitation: wetness
095FA: Farnum-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
095PB: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
095SB: Shellabarger---	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
095WA: Waldeck-----	Limitation: flooding	Limitation: flooding wetness soil blowing	Limitation: wetness soil blowing	Favorable
097LN: Lincoln-----	Limitation: deep to water	Limitation: flooding soil blowing droughty	Limitation: too sandy soil blowing	Limitation: droughty
097QW: Quinlan-----	Limitation: deep to water	Limitation: slope depth to rock	Limitation: slope depth to rock	Limitation: slope depth to rock
Woodward-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
097WA: Waldeck-----	Limitation: flooding	Limitation: flooding wetness	Limitation: wetness	Favorable
151AB: Albion-----	Limitation: deep to water	Limitation: soil blowing droughty	Limitation: too sandy soil blowing	Limitation: droughty
151AO: Albion-----	Limitation: deep to water	Limitation: slope soil blowing droughty	Limitation: too sandy soil blowing	Limitation: droughty
151AS: Albion-----	Limitation: deep to water	Limitation: slope soil blowing droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Shellabarger---	Limitation: deep to water	Limitation: slope soil blowing	Limitation: slope too sandy soil blowing	Limitation: slope
151BC: Blanket-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
151CA: Carwile-----	Limitation: percs slowly	Limitation: percs slowly wetness soil blowing	Limitation: erodes easily wetness soil blowing	Limitation: erodes easily percs slowly rooting depth
151CK: Case-----	Limitation: deep to water	Limitation: slope	Limitation: slope	Limitation: slope
Clark-----	Limitation: deep to water	Limitation: slope	Limitation: slope	Limitation: slope
151CM: Clark-----	Limitation: deep to water	Favorable	Favorable	Favorable
151CN: Clark-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
151CO: Clark-----	Limitation: deep to water	Favorable	Favorable	Favorable
Ost-----	Limitation: deep to water	Favorable	Favorable	Favorable

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
151FA: Farnum-----	Limitation: deep to water	Limitation: slope	Favorable	Favorable
151KP: Kanza-----	Limitation: flooding cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: wetness droughty
Plevna-----	Limitation: flooding	Limitation: flooding wetness soil blowing	Limitation: wetness soil blowing	Limitation: wetness
151OS: Ost-----	Limitation: deep to water	Favorable	Favorable	Favorable
151PM: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
151PN: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
151PO: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
Carwile-----	Limitation: percs slowly	Limitation: percs slowly wetness soil blowing	Limitation: erodes easily wetness soil blowing	Limitation: erodes easily percs slowly rooting depth
151SA: Albion-----	Limitation: deep to water	Limitation: slope soil blowing droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Kaski-----	Limitation: deep to water	Limitation: flooding	Favorable	Favorable
151SE: Shellabarger----	Limitation: deep to water	Limitation: soil blowing	Limitation: too sandy soil blowing	Favorable
1439: Crisfield-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
Ad: Albion-----	Limitation: deep to water	Limitation: slope soil blowing droughty	Limitation: too sandy soil blowing	Limitation: droughty
Shellabarger----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: soil blowing	Favorable
Ae: Albion-----	Limitation: deep to water	Limitation: slope soil blowing droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Shellabarger----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: slope soil blowing	Limitation: slope
AED: Arents, Earthen Dam-----	---	---	---	---
AN: Albion-----	Limitation: deep to water	Limitation: slope soil blowing droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Shellabarger----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: slope soil blowing	Limitation: slope
As: Clairemont-----	Limitation: deep to water	Limitation: erodes easily excess salt flooding	Limitation: erodes easily	Limitation: erodes easily excess salt
At: Attica-----	Limitation: deep to water	Limitation: fast intake slope soil blowing	Limitation: soil blowing	Favorable

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Ba: Blanket-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
Bb: Blanket-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
Bc: Blanket-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
Bf: Quinlan-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
Clairemont-----	Limitation: deep to water	Limitation: erodes easily flooding	Limitation: erodes easily	Limitation: erodes easily
Ca: Canadian-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
Cc: Case-----	Limitation: deep to water	Limitation: slope	Favorable	Favorable
Clark-----	Limitation: deep to water	Limitation: slope	Favorable	Favorable
Cd: Clairemont-----	Limitation: deep to water	Limitation: erodes easily flooding	Limitation: erodes easily	Limitation: erodes easily
Cf: Clairemont-----	Limitation: deep to water	Limitation: erodes easily flooding	Limitation: erodes easily	Limitation: erodes easily
Ck: Clark-----	Limitation: deep to water	Favorable	Favorable	Favorable
Fa: Farnum-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
Fm: Farnum-----	Limitation: deep to water	Favorable	Favorable	Favorable
Fr: Farnum-----	Limitation: deep to water	Favorable	Favorable	Favorable
Fu: Farnum-----	Limitation: deep to water	Favorable	Favorable	Favorable
Ga: Grant-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
Gb: Grant-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
Gc: Grant-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily	Limitation: erodes easily
GRP: Gravel Pits-----	---	---	---	---
INT: Aquolls-----	---	---	---	---
Ka: Kanza-----	Limitation: flooding cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: wetness droughty
Kf: Kingfisher-----	Limitation: deep to water	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock
Kv: Kingfisher-----	Limitation: deep to water	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock
Vernon-----	Limitation: deep to water	Limitation: erodes easily percs slowly depth to rock	Limitation: erodes easily percs slowly depth to rock	Limitation: erodes easily percs slowly depth to rock
Kz: Kingfisher-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Vernon-----	Limitation: deep to water	Limitation: percs slowly slope depth to rock	Limitation: erodes easily percs slowly depth to rock	Limitation: erodes easily percs slowly depth to rock
Ln: Lincoln-----	Limitation: deep to water	Limitation: flooding soil blowing droughty	Limitation: too sandy soil blowing	Limitation: droughty
LNN: Lincoln-----	Limitation: deep to water	Limitation: fast intake soil blowing droughty	Limitation: too sandy soil blowing	Limitation: droughty
Ma: Mangum-----	Limitation: deep to water	Limitation: flooding percs slowly slow intake	Limitation: percs slowly	Limitation: percs slowly
Mg: Mangum-----	Limitation: deep to water	Limitation: percs slowly slow intake	Limitation: percs slowly	Limitation: percs slowly
Drummond-----	Limitation: excess sodium percs slowly	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily excess sodium percs slowly
Mn: Minco-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
Na: Naron-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
Nb: Naron-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
Os: Ost-----	Limitation: deep to water	Favorable	Favorable	Favorable
Ot: Ost-----	Limitation: deep to water	Favorable	Favorable	Favorable
Pa: Pond Creek-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
Pd: Pond Creek-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
Ph: Dale-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
Pk: Buttermilk-----	Limitation: deep to water	Limitation: excess salt	Limitation: erodes easily	Limitation: erodes easily excess salt
Ps: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
Pt: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Tivoli-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Qn: Quinlan-----	Limitation: deep to water	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock
Qw: Quinlan-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
Woodward-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Rb: Knoco-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope too arid
Rock Outcrop----	Limitation: deep to water	Limitation: slope depth to rock	Limitation: slope depth to rock	Limitation: slope depth to rock
Sb: Shellabarger----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: soil blowing	Favorable
SBB: Shellabarger----	Limitation: deep to water	Limitation: soil blowing	Limitation: too sandy soil blowing	Favorable
Sc: Shellabarger----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: soil blowing	Favorable
Tv: Tivoli-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Vn: Vernon-----	Limitation: deep to water	Limitation: percs slowly slope depth to rock	Limitation: erodes easily percs slowly depth to rock	Limitation: erodes easily percs slowly depth to rock
Vr: Vernon-----	Limitation: deep to water	Limitation: percs slowly slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
Vs: Knoco-----	Limitation: deep to water	Limitation: slope slow intake droughty	Limitation: erodes easily percs slowly depth to rock	Limitation: erodes easily too arid droughty
Vernon-----	Limitation: deep to water	Limitation: percs slowly slope depth to rock	Limitation: erodes easily percs slowly depth to rock	Limitation: erodes easily percs slowly depth to rock
W: Water-----	---	---	---	---
Wa: Waldeck-----	Limitation: flooding	Limitation: flooding wetness soil blowing	Limitation: wetness soil blowing	Favorable
Wo: Woodward-----	Limitation: deep to water	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock
Quinlan-----	Limitation: deep to water	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock
Ws: Woodward-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock
Quinlan-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily depth to rock	Limitation: erodes easily depth to rock
Ya: Yahola-----	Limitation: deep to water	Limitation: flooding soil blowing	Limitation: soil blowing	Favorable
Ze: Zenda-----	Limitation: flooding	Limitation: flooding wetness	Limitation: wetness	Favorable

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
033CP: Clairemont-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.93	Very limited Deep to water	1.00
033OA: Obaro-----	100	Somewhat limited Seepage Depth to bedrock	0.70 0.04	Somewhat limited Piping Thin layer	0.76 0.70	Very limited Deep to water	1.00
033OB: Obaro-----	75	Somewhat limited Seepage Depth to bedrock Slope	0.70 0.04 0.02	Somewhat limited Piping Thin layer	0.76 0.70	Very limited Deep to water	1.00
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
033QR: Quinlan-----	55	Very limited Seepage Depth to bedrock Slope	1.00 0.69 0.00	Very limited Thin layer Piping	1.00 1.00	Very limited Deep to water	1.00
Woodward-----	45	Somewhat limited Seepage Depth to bedrock Slope	0.70 0.11 0.00	Very limited Piping Thin layer	1.00 0.86	Very limited Deep to water	1.00
033QT: Quinlan-----	55	Very limited Seepage Depth to bedrock Slope	1.00 0.69 0.21	Very limited Thin layer Piping	1.00 1.00	Very limited Deep to water	1.00
Woodward-----	45	Somewhat limited Seepage Depth to bedrock Slope	0.70 0.11 0.08	Very limited Piping Thin layer	1.00 0.86	Very limited Deep to water	1.00
033YE: Yahola-----	100	Very limited Seepage	1.00	Very limited Piping Seepage	1.00 0.07	Very limited Deep to water	1.00
077BP: Woodward-----	65	Somewhat limited Seepage Depth to bedrock Slope	0.70 0.11 0.00	Very limited Piping Thin layer	1.00 0.85	Very limited Deep to water	1.00
Port-----	35	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.82	Very limited Deep to water	1.00
077FU: Farnum-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.13	Very limited Deep to water	1.00
077GE: Gerlane-----	100	Very limited Seepage	1.00	Very limited Piping Seepage	1.00 0.09	Very limited Cutbanks cave Deep to water Slow refill	1.00 0.81 0.43
077MC: Minco-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
077MN: Minco-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
077MO: Minco-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
077QU: Quinlan-----	100	Very limited Seepage	1.00	Very limited Thin layer	1.00	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
		Depth to bedrock	0.84				
077SE: Shellabarger-----	100	Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
077WE: Quinlan-----	50	Very limited Seepage Depth to bedrock	1.00 0.84	Very limited Thin layer	1.00	Very limited Deep to water	1.00
Woodward-----	50	Somewhat limited Seepage Depth to bedrock	0.70 0.30	Very limited Piping Thin layer	1.00 0.98	Very limited Deep to water	1.00
095AC: Albion-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
095DA: Dillwyn-----	60	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.18	Very limited Cutbanks cave Deep to water	1.00 0.00
Plevna-----	40	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 1.00	Very limited Cutbanks cave	1.00
095FA: Farnum-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
095PB: Pratt-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
095SB: Shellabarger-----	100	Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
095WA: Waldeck-----	100	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.43	Very limited Cutbanks cave Deep to water	1.00 0.25
097LN: Lincoln-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.22	Very limited Deep to water	1.00
097QW: Quinlan-----	55	Very limited Seepage Depth to bedrock Slope	1.00 0.66 0.04	Very limited Thin layer Piping	1.00 1.00	Very limited Deep to water	1.00
Woodward-----	45	Somewhat limited Seepage Depth to bedrock Slope	0.70 0.19 0.01	Very limited Piping Thin layer	1.00 0.93	Very limited Deep to water	1.00
097WA: Waldeck-----	100	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.43	Very limited Cutbanks cave Deep to water	1.00 0.25
151AB: Albion-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
151AO: Albion-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
151AS: Albion-----	50	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Shellabarger-----	50	Slope	0.00				
		Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
		Slope	0.00				
151BC: Blanket-----	100	Somewhat limited Seepage	0.70	Somewhat limited Hard to pack	0.33	Very limited Deep to water	1.00
151CA: Carwile-----	100	Somewhat limited Seepage	0.57	Very limited Depth to saturated zone	1.00	Somewhat limited Slow refill	0.30
						Cutbanks cave	0.10
151CK: Case-----	60	Somewhat limited Seepage Slope	0.70 0.00	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
Clark-----	40	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
151CM: Clark-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
151CN: Clark-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping Seepage	0.12 0.08	Very limited Deep to water	1.00
151CO: Clark-----	70	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
Ost-----	30	Somewhat limited Seepage	0.05	Very limited Piping	1.00	Very limited Deep to water	1.00
151FA: Farnum-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping Seepage	0.21 0.06	Very limited Deep to water	1.00
151KP: Kanza-----	50	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.98	Very limited Cutbanks cave	1.00
Plevna-----	50	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 1.00	Very limited Cutbanks cave	1.00
151OS: Ost-----	100	Somewhat limited Seepage	0.05	Very limited Piping	1.00	Very limited Deep to water	1.00
151PM: Pratt-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
151PN: Pratt-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
151PO: Pratt-----	60	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
Carwile-----	40	Somewhat limited Seepage	0.57	Very limited Depth to saturated zone Seepage	1.00	Somewhat limited Slow refill	0.30
					0.02	Cutbanks cave	0.10
151SA: Albion-----	70	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
		Slope	0.00				
Kaski-----	30	Somewhat limited Seepage	0.70	Somewhat limited Piping Seepage	0.78 0.09	Very limited Deep to water	1.00
151SE: Shellabarger-----	100	Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
1439: Crisfield-----	100	Very limited Seepage	1.00	Somewhat limited Seepage Depth to saturated zone	0.78 0.18	Very limited Cutbanks cave Deep to water	1.00 0.44
Ad: Albion-----	60	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
Shellabarger-----	40	Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
Ae: Albion-----	55	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
Shellabarger-----	45	Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
AN: Albion-----	65	Very limited Seepage	1.00	Somewhat limited Seepage	0.98	Very limited Deep to water	1.00
Shellabarger-----	35	Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
As: Clairemont-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping Salinity	0.50 0.12	Very limited Deep to water	1.00
At: Attica-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.72	Very limited Deep to water	1.00
Ba: Blanket-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
Bb: Blanket-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
Bc: Blanket-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
Bf: Quinlan-----	60	Very limited Seepage Depth to bedrock Slope	1.00 0.69 0.36	Very limited Thin layer Piping	1.00 1.00	Very limited Deep to water	1.00
Clairemont-----	40	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.93	Very limited Deep to water	1.00
Ca: Canadian-----	100	Very limited Seepage	1.00	Very limited Piping Seepage	1.00 0.09	Very limited Deep to water	1.00
Cc: Case-----	50	Somewhat limited		Somewhat limited		Very limited	

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
		Seepage	0.70	Piping	0.12	Deep to water	1.00
Clark-----	50	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
Cd: Clairemont-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.93	Very limited Deep to water	1.00
Cf: Clairemont-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.93	Very limited Deep to water	1.00
Ck: Clark-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
Fa: Farnum-----	100	Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.07	Very limited Deep to water	1.00
Fm: Farnum-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
Fr: Farnum-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
Fu: Farnum-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
Ga: Grant-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.96	Very limited Deep to water	1.00
Gb: Grant-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.96	Very limited Deep to water	1.00
Gc: Grant-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.96	Very limited Deep to water	1.00
GRP: Gravel Pits-----	100	Not rated		Not rated		Not rated	
INT: Aquolls-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Somewhat limited Cutbanks cave	0.10
Ka: Kanza-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.65	Very limited Cutbanks cave	1.00
Kf: Kingfisher-----	100	Somewhat limited Depth to bedrock Seepage	0.17 0.05	Somewhat limited Thin layer Piping	0.91 0.57	Very limited Deep to water	1.00
Kv: Kingfisher-----	70	Somewhat limited Depth to bedrock Seepage	0.17 0.05	Somewhat limited Thin layer Piping	0.91 0.57	Very limited Deep to water	1.00
Vernon-----	30	Somewhat limited Depth to bedrock	0.37	Somewhat limited Thin layer Hard to pack	0.99 0.50	Very limited Deep to water	1.00
Kz: Kingfisher-----	60	Somewhat limited Depth to bedrock	0.17	Somewhat limited Thin layer	0.91	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Vernon-----	40	Seepage	0.05	Piping	0.57		
		Somewhat limited Depth to bedrock	0.37	Somewhat limited Thin layer Hard to pack	0.99 0.50	Very limited Deep to water	1.00
Ln: Lincoln-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.22	Very limited Deep to water	1.00
LNN: Lincoln-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.89	Very limited Deep to water	1.00
Ma: Mangum-----	100	Not limited		Somewhat limited Hard to pack	0.47	Very limited Deep to water	1.00
Mg: Mangum-----	65	Not limited		Somewhat limited Hard to pack	0.47	Very limited Deep to water	1.00
Drummond-----	35	Not limited		Not limited		Somewhat limited Deep to water Cutbanks cave Salty water	0.81 0.10 0.01
Mn: Minco-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
Na: Naron-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.08	Very limited Deep to water	1.00
Nb: Naron-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.08	Very limited Deep to water	1.00
Os: Ost-----	100	Somewhat limited Seepage	0.05	Very limited Piping	1.00	Very limited Deep to water	1.00
Ot: Ost-----	100	Somewhat limited Seepage	0.05	Very limited Piping	1.00	Very limited Deep to water	1.00
Pa: Pond Creek-----	100	Somewhat limited Seepage	0.05	Somewhat limited Piping	0.70	Very limited Deep to water	1.00
Pd: Pond Creek-----	100	Somewhat limited Seepage	0.05	Somewhat limited Piping	0.70	Very limited Deep to water	1.00
Ph: Dale-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.80	Very limited Deep to water	1.00
Pk: Buttermilk-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping Salinity	0.47 0.12	Very limited Deep to water	1.00
Ps: Pratt-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
Pt: Pratt-----	70	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
Tivoli-----	30	Very limited Seepage	1.00	Somewhat limited Seepage	0.99	Very limited Deep to water	1.00
Qn: Quinlan-----	100	Very limited Seepage Depth to bedrock	1.00 0.69	Very limited Thin layer Piping	1.00 1.00	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Qw: Quinlan-----	60	Very limited Seepage Depth to bedrock	1.00 0.69	Very limited Thin layer Piping	1.00 1.00	Very limited Deep to water	1.00
Woodward-----	40	Somewhat limited Seepage Depth to bedrock	0.70 0.01	Very limited Piping Thin layer	1.00 0.46	Very limited Deep to water	1.00
Rb: Knoco-----	70	Somewhat limited Depth to bedrock Slope	0.79 0.64	Very limited Thin layer Piping	1.00 1.00	Very limited Deep to water	1.00
Rock Outcrop-----	30	Not rated		Not rated		Not rated	
Sb: Shellabarger-----	100	Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
SBB: Shellabarger-----	100	Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
Sc: Shellabarger-----	100	Somewhat limited Seepage	0.70	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
Tv: Tivoli-----	100	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.99	Very limited Deep to water	1.00
Vn: Vernon-----	100	Somewhat limited Depth to bedrock	0.37	Somewhat limited Thin layer Hard to pack	0.99 0.50	Very limited Deep to water	1.00
Vr: Vernon-----	100	Somewhat limited Depth to bedrock	0.37	Somewhat limited Thin layer Hard to pack	0.99 0.50	Very limited Deep to water	1.00
Vs: Knoco-----	50	Somewhat limited Depth to bedrock	0.79	Very limited Thin layer Piping	1.00 1.00	Very limited Deep to water	1.00
Vernon-----	50	Somewhat limited Depth to bedrock	0.11	Somewhat limited Thin layer	0.85	Very limited Deep to water	1.00
W: Water-----	100	Not rated		Not rated		Not rated	
Wa: Waldeck-----	100	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.43	Very limited Cutbanks cave Deep to water	1.00 0.25
Wo: Woodward-----	70	Somewhat limited Seepage Depth to bedrock	0.70 0.01	Very limited Piping Thin layer	1.00 0.46	Very limited Deep to water	1.00
Quinlan-----	30	Very limited Seepage Depth to bedrock	1.00 0.69	Very limited Thin layer Piping	1.00 1.00	Very limited Deep to water	1.00
Ws: Woodward-----	65	Somewhat limited Seepage Depth to bedrock	0.70 0.01	Very limited Piping Thin layer	1.00 0.46	Very limited Deep to water	1.00
Quinlan-----	35	Very limited		Very limited		Very limited	

WATER MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ya: Yahola-----	100	Seepage Depth to bedrock	1.00 0.69	Thin layer Piping	1.00 1.00	Deep to water	1.00
		Very limited Seepage	1.00	Very limited Piping Seepage	1.00 0.07	Very limited Deep to water	1.00
Ze: Zenda-----	100	Somewhat limited Seepage	0.70	Somewhat limited Depth to saturated zone Piping	0.43 0.18	Somewhat limited Slow refill	0.30
						Deep to water Cutbanks cave	0.25 0.10

SANITARY FACILITIES
Barber County, Kansas

Sanitary Facilities

The following tables show the degree and kind of soil limitations that affect septic tank absorption fields, sewage lagoons, sanitary landfills, and daily cover for landfill. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Septic tank absorption fields are areas in which effluent from a septic tank is distributed into the soil through subsurface tiles or perforated pipe. Only that part of the soil between depths of 24 and 60 inches is evaluated. The ratings are based on the soil properties that affect absorption of the effluent, construction and maintenance of the system, and public health. Permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, and flooding affect absorption of the effluent. Stones and boulders, ice, and bedrock or a cemented pan interfere with installation. Subsidence interferes with installation and maintenance. Excessive slope may cause lateral seepage and surfacing of the effluent in downslope areas.

Some soils are underlain by loose sand and gravel or fractured bedrock at a depth of less than 4 feet below the distribution lines. In these soils the absorption field may not adequately filter the effluent, particularly when the system is new. As a result, the ground water may become contaminated.

Sewage lagoons are shallow ponds constructed to hold sewage while aerobic bacteria decompose the solid and liquid wastes. Lagoons should have a nearly level floor surrounded by cut slopes or embankments of compacted soil. Nearly impervious soil material for the lagoon floor and sides is required to minimize seepage and contamination of ground water. Considered in the ratings are slope, permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, flooding, large stones, and content of organic matter.

Soil permeability is a critical property affecting the suitability for sewage lagoons. Most porous soils eventually become sealed when they are used as sites for sewage lagoons. Until sealing occurs, however, the hazard of pollution is severe. Soils that have a permeability rate of more than 2 inches per hour are too porous for the proper functioning of sewage lagoons. In these soils, seepage of the effluent can result in contamination of the ground water. Ground-water contamination is also a hazard if fractured bedrock is within a depth of 40 inches, if the water table is high enough to raise the level of sewage in the lagoon, or if floodwater overtops the lagoon.

A high content of organic matter is detrimental to proper functioning of the lagoon because it inhibits aerobic activity. Slope, bedrock, and cemented pans can cause construction problems, and large stones can hinder compaction of the lagoon floor. If the lagoon is to be uniformly deep throughout, the slope must be gentle enough and the soil material must be thick enough over bedrock or a cemented pan to make land smoothing practical.

A trench sanitary landfill is an area where solid waste is placed in successive layers in an excavated trench. The waste is spread, compacted, and covered daily with a thin layer of soil excavated at the site. When the trench is full, a final cover of soil material at least 2 feet thick is placed over the landfill. The ratings in the table are based on the soil properties that affect the risk of pollution, the ease of excavation, trafficability, and revegetation. These properties include permeability, depth to bedrock or a cemented pan, depth to a water table, ponding, slope, flooding, texture, stones and boulders, highly organic layers, soil reaction, and content of salts and sodium. Unless otherwise stated, the ratings apply only to that part of the soil within a depth of about 6 feet. For deeper trenches, onsite investigation may be needed.

Hard, nonrippable bedrock, creviced bedrock, or highly permeable strata in or directly below the proposed trench bottom can affect the ease of excavation and the hazard of ground-water pollution. Slope affects construction of the trenches and the movement of surface water around the landfill. It also affects the construction and performance of roads in areas of the landfill.

Soil texture and consistence affect the ease with which the trench is dug and the ease with which the soil can be used as daily or final cover. They determine the workability of the soil when dry and when wet. Soils that are plastic and sticky when wet are difficult to excavate, grade, or compact and are difficult to place as a uniformly thick cover over a layer of refuse.

The soil material used as the final cover for a trench landfill should be suitable for plants. It should not have excess sodium or salts and should not be too acid. The surface layer generally has the best workability, the highest content of organic matter, and the best potential for plants. Material from the surface layer should be stockpiled for use as the final cover.

SANITARY FACILITIES
Barber County, Kansas

In an area sanitary landfill, solid waste is placed in successive layers on the surface of the soil. The waste is spread, compacted, and covered daily with a thin layer of soil from a source away from the site. A final cover of soil material at least 2 feet thick is placed over the completed landfill. The ratings in the table are based on the soil properties that affect trafficability and the risk of pollution. These properties include flooding, permeability, depth to a water table, ponding, slope, and depth to bedrock or a cemented pan.

Flooding is a serious problem because it can result in pollution in areas downstream from the landfill. If permeability is too rapid or if fractured bedrock, a fractured cemented pan, or the water table is close to the surface, the leachate can contaminate the water supply. Slope is a consideration because of the extra grading required to maintain roads in the steeper areas of the landfill. Also, leachate may flow along the surface of the soils in the steeper areas and cause difficult seepage problems.

Daily cover for landfill is the soil material that is used to cover compacted solid waste in an area sanitary landfill. The soil material is obtained offsite, transported to the landfill, and spread over the waste. The ratings in the table also apply to the final cover for a landfill. They are based on the soil properties that affect workability, the ease of digging, and the ease of moving and spreading the material over the refuse daily during wet and dry periods. These properties include soil texture, depth to a water table, ponding, rock fragments, slope, depth to bedrock or a cemented pan, reaction, and content of salts, sodium, or lime.

Loamy or silty soils that are free of large stones and excess gravel are the best cover for a landfill. Clayey soils may be sticky and difficult to spread; sandy soils are subject to wind erosion.

Slope affects the ease of excavation and of moving the cover material. Also, it can influence runoff, erosion, and reclamation of the borrow area.

After soil material has been removed, the soil material remaining in the borrow area must be thick enough over bedrock, a cemented pan, or the water table to permit revegetation. The soil material used as the final cover for a landfill should be suitable for plants. It should not have excess sodium, salts, or lime and should not be too acid.

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
033CP: Clairemont-----	100	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
033OA: Obaro-----	100	Very limited Depth to bedrock Restricted permeability Slope	1.00 0.50 0.04	Very limited Depth to soft bedrock Slope	1.00 1.00
033OB: Obaro-----	75	Very limited Depth to bedrock Slope Restricted permeability	1.00 0.96 0.50	Very limited Depth to soft bedrock Slope	1.00 1.00
Rock Outcrop-----	25	Not rated		Not rated	
033QR: Quinlan-----	55	Very limited Depth to bedrock Slope	1.00 0.37	Very limited Depth to soft bedrock Slope	1.00 1.00
Woodward-----	45	Very limited Depth to bedrock Restricted permeability Slope	1.00 0.50 0.37	Very limited Depth to soft bedrock Slope	1.00 1.00
033QT: Quinlan-----	55	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00
Woodward-----	45	Very limited Depth to bedrock Slope Restricted permeability	1.00 1.00 0.50	Very limited Depth to soft bedrock Slope	1.00 1.00
033YE: Yahola-----	100	Very limited Flooding	1.00	Very limited Flooding Seepage	1.00 1.00
077BP: Woodward-----	65	Very limited Depth to bedrock Restricted permeability Slope	1.00 0.50 0.37	Very limited Depth to soft bedrock Slope	1.00 1.00
Port-----	35	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage Slope	1.00 0.50 0.00
077FU: Farnum-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope Seepage	0.67 0.50
077GE: Gerlane-----	100	Very limited Flooding Depth to saturated zone Restricted permeability	1.00 1.00 0.68	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 0.71
077MC: Minco-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
077MN: Minco-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.00

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
077MO: Minco-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope Seepage	0.67 0.50
077QU: Quinlan-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock Slope	1.00 0.67
077SE: Shellabarger-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.00
077WE: Quinlan-----	50	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock Slope	1.00 0.00
Woodward-----	50	Very limited Depth to bedrock Restricted permeability	1.00 0.50	Very limited Depth to soft bedrock Slope	1.00 0.00
095AC: Albion-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.67
095DA: Dillwyn-----	60	Very limited Depth to saturated zone Filtering capacity	1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00
Plevna-----	40	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
095FA: Farnum-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
095PB: Pratt-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.91
095SB: Shellabarger-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.00
095WA: Waldeck-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
097LN: Lincoln-----	100	Very limited Flooding Filtering capacity Depth to saturated zone	1.00 1.00 0.08	Very limited Flooding Seepage	1.00 1.00
097QW: Quinlan-----	55	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00
Woodward-----	45	Very limited Depth to bedrock Slope	1.00 0.84	Very limited Depth to soft bedrock Slope	1.00 1.00

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
097WA: Waldeck-----	100	Restricted permeability	0.50		
		Very limited Flooding	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
151AB: Albion-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.09
151AO: Albion-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.67
151AS: Albion-----	50	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Slope	0.37	Slope	1.00
Shellabarger-----	50	Somewhat limited Restricted permeability	0.50	Very limited Slope	1.00
		Slope	0.37	Seepage	0.50
151BC: Blanket-----	100	Very limited Restricted permeability	1.00	Somewhat limited Seepage	0.50
				Slope	0.09
151CA: Carwile-----	100	Very limited Restricted permeability	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Seepage	0.32
151CK: Case-----	60	Somewhat limited Restricted permeability	0.50	Very limited Slope	1.00
		Slope	0.37	Seepage	0.50
Clark-----	40	Somewhat limited Restricted permeability	0.50	Very limited Slope	1.00
		Slope	0.16	Seepage	0.50
151CM: Clark-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
				Slope	0.09
151CN: Clark-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
				Slope	0.00
151CO: Clark-----	70	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Ost-----	30	Very limited Restricted permeability	1.00	Not limited	
151FA: Farnum-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope	0.67
				Seepage	0.50
151KP: Kanza-----	50	Very limited Flooding	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
Plevna-----	50	Very limited Flooding	1.00	Very limited Flooding	1.00

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
		Depth to saturated zone	1.00	Seepage	1.00
				Depth to saturated zone	1.00
151OS: Ost-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.09
151PM: Pratt-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.91
151PN: Pratt-----	100	Very limited Filtering capacity Slope	1.00 0.16	Very limited Slope	1.00
151PO: Pratt-----	60	Very limited Filtering capacity	1.00	Seepage	1.00
				Very limited Seepage	1.00
Carwile-----	40	Slope		Slope	0.91
		Very limited Restricted permeability	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Seepage	0.32
151SA: Albion-----	70	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Slope	0.37	Slope	1.00
Kaski-----	30	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
151SE: Shellabarger-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
				Slope	0.09
1439: Crisfield-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Flooding	0.40	Flooding	0.40
Ad: Albion-----	60	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.09
Shellabarger-----	40	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
				Slope	0.09
Ae: Albion-----	55	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Slope	0.16	Slope	1.00
Shellabarger-----	45	Somewhat limited Restricted permeability	0.50	Very limited Slope	1.00
		Slope	0.16	Seepage	0.50
AED: Arents, Earthen Dam-	100	Not rated		Not rated	
AN: Albion-----	65	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Slope	0.16	Slope	1.00
Shellabarger-----	35	Somewhat limited Restricted permeability	0.50	Very limited Slope	1.00
		Slope	0.16	Seepage	0.50

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
As: Clairemont-----	100	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
At: Attica-----	100	Not limited		Very limited Seepage Slope	1.00 0.09
Ba: Blanket-----	100	Very limited Restricted permeability	1.00	Somewhat limited Seepage	0.50
Bb: Blanket-----	100	Very limited Restricted permeability	1.00	Somewhat limited Seepage Slope	0.50 0.00
Bc: Blanket-----	100	Very limited Restricted permeability	1.00	Somewhat limited Seepage Slope	0.50 0.00
Bf: Quinlan-----	60	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00
Clairemont-----	40	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
Ca: Canadian-----	100	Very limited Filtering capacity Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40
Cc: Case-----	50	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.33
Clark-----	50	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.33
Cd: Clairemont-----	100	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
Cf: Clairemont-----	100	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
Ck: Clark-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Fa: Farnum-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Fm: Farnum-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Fr: Farnum-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.00
Fu: Farnum-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.00
Ga: Grant-----	100	Somewhat limited		Somewhat limited	

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Gb: Grant-----	100	Restricted permeability	0.50	Seepage	0.50
		Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Gc: Grant-----	100			Slope	0.00
		Somewhat limited Restricted permeability	0.50	Somewhat limited Slope	0.67
GRP: Gravel Pits-----	100			Seepage	0.50
		Not rated		Not rated	
INT: Aquolls-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Ponding	1.00	Ponding	1.00
Ka: Kanza-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Seepage	1.00
Kf: Kingfisher-----	100	Filtering capacity	1.00	Depth to saturated zone	1.00
		Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
Kv: Kingfisher-----	70	Restricted permeability	1.00	Slope	0.00
		Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
Vernon-----	30	Restricted permeability	1.00	Slope	0.00
		Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
Kz: Kingfisher-----	60			Slope	0.00
		Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
Vernon-----	40	Restricted permeability	1.00	Slope	0.67
		Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
Ln: Lincoln-----	100			Slope	0.67
		Very limited Flooding	1.00	Very limited Flooding	1.00
LNN: Lincoln-----	100	Filtering capacity	1.00	Seepage	1.00
		Depth to saturated zone	0.08		
Ma: Mangum-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00
		Restricted permeability	1.00		
Mg: Mangum-----	65	Very limited Restricted permeability	1.00	Somewhat limited Flooding	0.40
		Flooding	0.40		
Drummond-----	35	Very limited Restricted permeability	1.00	Somewhat limited Depth to saturated zone	0.71

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Mn: Minco-----	100	Depth to saturated zone	1.00	Flooding	0.40
		Flooding	0.40		
Na: Naron-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Nb: Naron-----	100	Somewhat limited Restricted permeability	0.50	Very limited Seepage	1.00
Os: Ost-----	100	Somewhat limited Restricted permeability	0.50	Very limited Seepage	1.00
				Slope	0.00
Ot: Ost-----	100	Very limited Restricted permeability	1.00	Not limited	
Pa: Pond Creek-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.00
Pd: Pond Creek-----	100	Very limited Restricted permeability	1.00	Not limited	
Ph: Dale-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.00
Pk: Buttermilk-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
		Flooding	0.40	Flooding	0.40
Ps: Pratt-----	100	Depth to saturated zone	0.22		
Pt: Pratt-----	70	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Tivoli-----	30	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
		Slope	0.16	Slope	1.00
Qn: Quinlan-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
Qw: Quinlan-----	60	Slope	0.16	Slope	1.00
		Depth to bedrock	1.00	Depth to soft bedrock	1.00
Woodward-----	40	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
		Restricted permeability	0.50	Slope	1.00
Rb: Knoco-----	70	Slope	0.16	Depth to soft bedrock	1.00
Rock Outcrop-----	30	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
		Slope	1.00	Slope	1.00
		Not rated		Not rated	

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Sb: Shellabarger-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope	0.67
				Seepage	0.50
SBB: Shellabarger-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Sc: Shellabarger-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope	0.67
				Seepage	0.50
Tv: Tivoli-----	100	Very limited Filtering capacity Slope	1.00	Very limited Seepage	1.00
			0.84	Slope	1.00
Vn: Vernon-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
				Slope	0.33
Vr: Vernon-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
		Slope	0.16	Slope	1.00
Vs: Knoco-----	50	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
				Slope	0.09
Vernon-----	50	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
				Slope	0.09
W: Water-----	100	Not rated		Not rated	
Wa: Waldeck-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
Wo: Woodward-----	70	Very limited Depth to bedrock	1.00	Somewhat limited Depth to soft bedrock	1.00
		Restricted permeability	0.50	Slope	0.00
Quinlan-----	30	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
				Slope	0.00
Ws: Woodward-----	65	Very limited Depth to bedrock	1.00	Somewhat limited Depth to soft bedrock	1.00
		Restricted permeability	0.50	Slope	0.67
Quinlan-----	35	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock	1.00
				Slope	0.67
Ya: Yahola-----	100	Very limited Flooding	1.00	Very limited Flooding Seepage	1.00
					1.00
Ze: Zenda-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Restricted permeability	0.50	Seepage	0.50

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit				

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
033CP: Clairemont-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Not limited	
033OA: Obaro-----	100	Very limited Depth to bedrock Seepage Too clayey Slope	1.00 1.00 0.50 0.04	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Too clayey Slope	1.00 0.50 0.04
033OB: Obaro-----	75	Very limited Depth to bedrock Seepage Slope Too clayey	1.00 1.00 0.96 0.50	Very limited Depth to bedrock Slope	1.00 0.96	Very limited Depth to bedrock Slope Too clayey	1.00 0.96 0.50
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
033QR: Quinlan-----	55	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.37	Very limited Depth to bedrock Slope	1.00 0.37	Very limited Depth to bedrock Slope	1.00 0.37
Woodward-----	45	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.37	Very limited Depth to bedrock Slope	1.00 0.37	Very limited Depth to bedrock Slope	1.00 0.37
033QT: Quinlan-----	55	Very limited Depth to bedrock Slope Seepage	1.00 1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00
Woodward-----	45	Very limited Depth to bedrock Slope Seepage	1.00 1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00
033YE: Yahola-----	100	Very limited Flooding Seepage	1.00 1.00	Very limited Flooding Seepage	1.00 1.00	Somewhat limited Seepage	0.50
077BP: Woodward-----	65	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.37	Very limited Depth to bedrock Slope	1.00 0.37	Very limited Depth to bedrock Slope	1.00 0.37
Port-----	35	Very limited Flooding Too clayey	1.00 0.50	Very limited Flooding	1.00	Somewhat limited Too clayey	0.50
077FU: Farnum-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
077GE: Gerlane-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Somewhat limited Seepage	0.50
077MC: Minco-----	100	Not limited		Not limited		Not limited	
077MN: Minco-----	100	Not limited		Not limited		Not limited	
077MO: Minco-----	100	Not limited		Not limited		Not limited	
077QU: Quinlan-----	100	Very limited Depth to bedrock Seepage	1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
077SE: Shellabarger-----	100	Not limited		Not limited		Not limited	
077WE: Quinlan-----	50	Very limited Depth to bedrock Seepage	1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
Woodward-----	50	Very limited Depth to bedrock Seepage	1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
095AC: Albion-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage Gravel content	1.00 1.00 0.00
095DA: Dillwyn-----	60	Very limited		Very limited		Very limited	

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Plevna-----	40	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
		Seepage	1.00	Seepage	1.00	Depth to saturated zone	0.86
		Too Sandy	1.00			Too Sandy	0.50
		Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	0.50
095FA: Farnum-----	100	Seepage	1.00	Seepage	1.00		
095PB: Pratt-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
095SB: Shellabarger-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Seepage Too Sandy	1.00 0.50
095WA: Waldeck-----	100	Not limited		Not limited		Not limited	
097LN: Lincoln-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Seepage	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Too Sandy	0.50
		Seepage	1.00	Seepage	1.00		
		Too Sandy	1.00				
097QW: Quinlan-----	55	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00
Woodward-----	45	Seepage	1.00				
097WA: Waldeck-----	100	Very limited Depth to bedrock Slope	1.00 1.00 0.84	Very limited Depth to bedrock Slope	1.00 1.00 0.84	Very limited Depth to bedrock Slope	1.00 1.00 0.84
		Seepage	1.00	Seepage	1.00		
151AB: Albion-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Somewhat limited Seepage	0.50
151AO: Albion-----	100	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	0.09
151AS: Albion-----	100	Seepage	1.00	Seepage	1.00		
Shellabarger-----	50	Too Sandy	1.00	Too Sandy	1.00	Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
151BC: Blanket-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Seepage	1.00
151CA: Carwile-----	100	Too Sandy	1.00	Too Sandy	1.00	Too Sandy	1.00
151CK: Case-----	60	Seepage	1.00	Seepage	1.00	Seepage	1.00
Clark-----	40	Too Sandy	1.00	Too Sandy	1.00	Too Sandy	1.00
		Seepage	1.00	Seepage	1.00	Seepage	1.00
Clark-----	40	Somewhat limited Too clayey	0.50	Somewhat limited Too clayey	0.50	Somewhat limited Too clayey	0.50
		Slope	0.37	Slope	0.37	Slope	0.37
Clark-----	40	Somewhat limited Too clayey	0.50	Somewhat limited Too clayey	0.50	Somewhat limited Too clayey	0.50
		Slope	0.16	Slope	0.16	Slope	0.16

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
151CM: Clark-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
151CN: Clark-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
151CO: Clark-----	70	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Ost-----	30	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
151FA: Farnum-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
151KP: Kanza-----	50	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Too Sandy Seepage Depth to saturated zone	1.00 1.00 1.00
Plevna-----	50	Too Sandy Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 0.50
151OS: Ost-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
151PM: Pratt-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Seepage Too Sandy	1.00 0.50
151PN: Pratt-----	100	Very limited Seepage Too Sandy Slope	1.00 1.00 0.16	Very limited Seepage Slope	1.00 0.16	Very limited Seepage Too Sandy Slope	1.00 0.50 0.16
151PO: Pratt-----	60	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Seepage Too Sandy	1.00 0.50
Carwile-----	40	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Hard to compact	1.00
151SA: Albion-----	70	Very limited Seepage Too Sandy Slope	1.00 1.00 0.37	Very limited Seepage Slope	1.00 0.37	Very limited Too Sandy Seepage Slope	1.00 1.00 0.37
Kaski-----	30	Very limited Flooding	1.00	Very limited Flooding	1.00	Not limited	
151SE: Shellabarger-----	100	Not limited		Not limited		Not limited	
1439: Crisfield-----	100	Very limited Depth to saturated zone Seepage Too Sandy Flooding	1.00 1.00 1.00 0.40	Very limited Depth to saturated zone Seepage Flooding	1.00 1.00 0.40	Very limited Too Sandy Seepage Depth to saturated zone	1.00 1.00 0.01
Ad: Albion-----	60	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage Gravel content	1.00 1.00 0.00
Shellabarger-----	40	Not limited		Not limited		Not limited	
Ae: Albion-----	55	Very limited Seepage Too Sandy Slope	1.00 1.00 0.16	Very limited Seepage Slope	1.00 0.16	Very limited Too Sandy Seepage Slope Gravel content	1.00 1.00 0.16 0.00
Shellabarger-----	45	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
AN: Albion-----	65	Very limited Seepage Too Sandy Slope	1.00 1.00 0.16	Very limited Seepage Slope	1.00 0.16	Very limited Too Sandy Seepage Slope	1.00 1.00 0.16
Shellabarger-----	35	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16
As: Clairemont-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Not limited	
At: Attica-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Somewhat limited Seepage Too Sandy	0.50 0.50
Ba: Blanket-----	100	Somewhat limited Too clayey	0.50	Not limited		Very limited Hard to compact Too clayey	1.00 0.50
Bb: Blanket-----	100	Somewhat limited Too clayey	0.50	Not limited		Very limited Hard to compact Too clayey	1.00 0.50
Bc: Blanket-----	100	Somewhat limited Too clayey	0.50	Not limited		Very limited Hard to compact Too clayey	1.00 0.50
Bf: Quinlan-----	60	Very limited Depth to bedrock Slope Seepage	1.00 1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00
Clairemont-----	40	Very limited Flooding	1.00	Very limited Flooding	1.00	Not limited	
Ca: Canadian-----	100	Very limited Seepage Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40	Somewhat limited Seepage	0.50
Cc: Case-----	50	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Clark-----	50	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Cd: Clairemont-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Not limited	
Cf: Clairemont-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Not limited	
Ck: Clark-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Fa: Farnum-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Fm: Farnum-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Fr: Farnum-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Fu: Farnum-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Ga: Grant-----	100	Not limited		Not limited		Not limited	
Gb: Grant-----	100	Not limited		Not limited		Not limited	
Gc: Grant-----	100	Not limited		Not limited		Not limited	
GRP: Gravel Pits-----	100	Not rated		Not rated		Not rated	
INT: Aquolls-----	100	Very limited Depth to saturated zone Ponding Seepage	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
Ka: Kanza-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Seepage	1.00

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Kf: Kingfisher-----	100	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00	Too Sandy	0.50
		Too Sandy	1.00				
Kv: Kingfisher-----	70	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Seepage	1.00			Too clayey	0.50
		Too clayey	0.50				
Vernon-----	30	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Seepage	1.00			Too clayey	0.50
		Too clayey	0.50				
Kz: Kingfisher-----	60	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Seepage	1.00			Too clayey	0.50
		Too clayey	0.50				
Vernon-----	40	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Seepage	1.00			Too clayey	1.00
		Too clayey	1.00			Hard to compact	1.00
Ln: Lincoln-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Too Sandy	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
		Seepage	1.00	Seepage	1.00		
LNN: Lincoln-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Too Sandy	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
		Seepage	1.00	Seepage	1.00		
Ma: Mangum-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Too clayey	1.00
		Too clayey	1.00			Hard to compact	1.00
Mg: Mangum-----	65	Very limited Too clayey	1.00	Somewhat limited Flooding	0.40	Very limited Too clayey	1.00
		Flooding	0.40			Hard to compact	1.00
Drummond-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Hard to compact	1.00
		Seepage	1.00	Flooding	0.40		
		Flooding	0.40				
Mn: Minco-----	100	Not limited		Not limited		Not limited	
Na: Naron-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Not limited	
Nb: Naron-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Not limited	
Os: Ost-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Ot: Ost-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Pa: Pond Creek-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Pd: Pond Creek-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Ph: Dale-----	100	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Not limited	
Pk: Buttermilk-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Flooding	0.40	Flooding	0.40		

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ps: Pratt-----	100	Very limited Seepage Too Sandy Slope	1.00 1.00 0.00	Very limited Seepage Slope	1.00 0.00	Very limited Seepage Too Sandy Slope	1.00 0.50 0.00
Pt: Pratt-----	70	Very limited Seepage Too Sandy Slope	1.00 1.00 0.16	Very limited Seepage Slope	1.00 0.16	Very limited Seepage Too Sandy Slope	1.00 0.50 0.16
Tivoli-----	30	Very limited Seepage Too Sandy Slope	1.00 1.00 0.16	Very limited Seepage Slope	1.00 0.16	Very limited Too Sandy Seepage Slope	1.00 1.00 0.16
Qn: Quinlan-----	100	Very limited Depth to bedrock Seepage	1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
Qw: Quinlan-----	60	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.16	Very limited Depth to bedrock Slope	1.00 0.16	Very limited Depth to bedrock Slope	1.00 0.16
Woodward-----	40	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.16	Somewhat limited Depth to bedrock Slope	1.00 0.16	Somewhat limited Depth to bedrock Slope	1.00 0.16
Rb: Knoco-----	70	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Slope	1.00	Very limited Depth to bedrock Slope Hard to compact	1.00 1.00 1.00
Rock Outcrop-----	30	Not rated		Not rated		Not rated	
Sb: Shellabarger-----	100	Not limited		Not limited		Not limited	
SBB: Shellabarger-----	100	Not limited		Not limited		Not limited	
Sc: Shellabarger-----	100	Not limited		Not limited		Not limited	
Tv: Tivoli-----	100	Very limited Seepage Too Sandy Slope	1.00 1.00 0.84	Very limited Seepage Slope	1.00 0.84	Very limited Too Sandy Seepage Slope	1.00 1.00 0.84
Vn: Vernon-----	100	Very limited Depth to bedrock Too clayey Seepage	1.00 1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Too clayey Hard to compact	1.00 1.00 1.00
Vr: Vernon-----	100	Very limited Depth to bedrock Too clayey Seepage Slope	1.00 1.00 1.00 0.16	Very limited Depth to bedrock Slope	1.00 0.16	Very limited Depth to bedrock Too clayey Hard to compact Slope	1.00 1.00 1.00 0.16
Vs: Knoco-----	50	Very limited Depth to bedrock	1.00	Not limited		Very limited Depth to bedrock Hard to compact	1.00 1.00
Vernon-----	50	Very limited Depth to bedrock Too clayey Seepage	1.00 1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Too clayey Hard to compact	1.00 1.00 1.00
W: Water-----	100	Not rated		Not rated		Not rated	
Wa: Waldeck-----	100	Very limited Flooding Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Seepage Too Sandy Depth to saturated zone	1.00 1.00 0.09
Wo: Woodward-----	70	Very limited Depth to bedrock Seepage	1.00 1.00	Somewhat limited Depth to bedrock	1.00	Somewhat limited Depth to bedrock	1.00
Quinlan-----	30	Very limited		Very limited		Very limited	

SANITARY FACILITIES--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ws: Woodward-----	65	Depth to bedrock Seepage	1.00 1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Very limited Depth to bedrock Seepage	1.00 1.00	Somewhat limited Depth to bedrock	1.00	Somewhat limited Depth to bedrock	1.00
Quinlan-----	35	Very limited Depth to bedrock Seepage	1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Very limited Depth to bedrock Seepage	1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
Ya: Yahola-----	100	Very limited Flooding Seepage	1.00 1.00	Very limited Flooding Seepage	1.00 1.00	Somewhat limited Seepage	0.50
		Very limited Flooding Seepage	1.00 1.00	Very limited Flooding Seepage	1.00 1.00	Somewhat limited Seepage	0.50
Ze: Zenda-----	100	Very limited Flooding Depth to saturated zone Too clayey	1.00 1.00 1.00 0.50	Very limited Flooding Depth to saturated zone	1.00 1.00 1.00	Somewhat limited Too clayey Depth to saturated zone	0.50 0.09
		Very limited Flooding Depth to saturated zone Too clayey	1.00 1.00 1.00 0.50	Very limited Flooding Depth to saturated zone	1.00 1.00 1.00	Somewhat limited Too clayey Depth to saturated zone	0.50 0.09

AGRICULTURAL WASTE MANAGEMENT Barber County, Kansas

The nature of the soil is also important in the application of organic wastes and wastewater to land as fertilizers and irrigation; it is also important when the soil is used as a medium for treatment and disposal of these wastes. Favorable soil properties are required to prevent environmental damage.

The use of organic wastes and wastewater as production resources will result in energy conservation, prevent the waste of these important resources, and prevent problems associated with their disposal. Where disposal is the goal, and a maximum amount is disposed in a minimum area to hold costs to a minimum, risk of environmental damage is the principal constraint. Where the reuse goal is pursued, and a minimum amount is applied to a maximum area to obtain the greatest benefit, environmental damage is unlikely.

Interpretations developed for waste management may include ratings for (1) manure and food processing wastes; (2) municipal sewage sludge; (3) irrigation use of wastewater; or (4) treatment of wastewater by the slow rate process, overland flow process, or rapid infiltration process. If available, these should be located in this subsection.

Soil properties are important considerations in areas where soils are used as sites for the treatment and disposal of organic waste and wastewater. Selection of soils with properties that favor waste management can help to prevent environmental damage.

The Ag-Waste tables show the degree and kind of soil limitations affecting the treatment of agricultural waste, including municipal and food-processing wastewater and effluent from lagoons or storage ponds. Municipal wastewater is the waste stream from a municipality. It contains domestic waste and may contain industrial waste. It may have received primary or secondary treatment. It is rarely untreated sewage. Food-processing wastewater results from the preparation of fruits, vegetables, milk, cheese, and meats for public consumption. In places it is high in content of sodium and chloride. In the context of these tables, the effluent in lagoons and storage ponds is from facilities used to treat or store food-processing wastewater or domestic or animal waste. Domestic and food-processing wastewater is very dilute, and the effluent from the facilities that treat or store it commonly is very low in content of carbonaceous and nitrogenous material; the content of nitrogen commonly ranges from 10 to 30 milligrams per liter. The wastewater from animal waste treatment lagoons or storage ponds, however, has much higher concentrations of these materials, mainly because the manure has not been diluted as much as the domestic waste. The content of nitrogen in this wastewater generally ranges from 50 to 2,000 milligrams per liter. When wastewater is applied, checks should be made to ensure that nitrogen, phosphorus, heavy metals, and salts are not added in excessive amounts.

The ratings in the tables are for waste management systems that not only dispose of and treat organic waste or wastewater but also are beneficial to crops (application of manure and food-processing waste, application of sewage sludge, and disposal of wastewater by irrigation) and for waste management systems that are designed only for the purpose of wastewater disposal and treatment (overland flow of wastewater, rapid infiltration of wastewater, and slow rate treatment of wastewater).

The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect agricultural waste management. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are generally favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Application of manure and food-processing waste not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. Manure is the excrement of livestock and poultry, and food-processing waste is damaged fruit and vegetables and the peelings, stems, leaves, pits, and soil particles removed in food preparation. The manure and food-processing waste are either solid, slurry, or liquid. Their nitrogen content varies. A high content of nitrogen limits the application rate. Toxic or otherwise dangerous wastes, such as those mixed with the lye used in food processing, are not considered in the ratings.

The ratings are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the waste is applied, and the method by which the waste is applied. The properties that affect absorption include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, and available water capacity. The properties that affect plant growth and microbial activity include reaction, the sodium adsorption ratio, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of waste. Permanently frozen soils are unsuitable for waste treatment.

Application of sewage sludge not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. In the context of this table, sewage sludge is the residual product of the treatment of municipal sewage. The solid component consists mainly of cell mass, primarily bacteria cells that developed during secondary treatment and have incorporated soluble organics into their own bodies. The sludge has small amounts of sand, silt, and other solid debris. The content of nitrogen varies. Some sludge has constituents that are toxic to plants or hazardous to the food chain, such as heavy metals and exotic organic compounds, and should be analyzed chemically prior to use.

AGRICULTURAL WASTE MANAGEMENT
Barber County, Kansas

The content of water in the sludge ranges from about 98 percent to less than 40 percent. The sludge is considered liquid if it is more than about 90 percent water, slurry if it is about 50 to 90 percent water, and solid if it is less than about 50 percent water.

The ratings in the table are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the sludge is applied, and the method by which the sludge is applied. The properties that affect absorption, plant growth, and microbial activity include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, available water capacity, reaction, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of sludge. Permanently frozen soils are unsuitable for waste treatment.

Disposal of wastewater by irrigation not only disposes of municipal wastewater and wastewater from food-processing plants, lagoons, and storage ponds but also can improve crop production by increasing the amount of water available to crops. The ratings in the table are based on the soil properties that affect the design, construction, management, and performance of the irrigation system. The properties that affect design and management include the sodium adsorption ratio, depth to a water table, ponding, available water capacity, permeability, slope, and flooding. The properties that affect construction include stones, cobbles, depth to bedrock or a cemented pan, depth to a water table, and ponding.

The properties that affect performance include depth to bedrock or a cemented pan, bulk density, the sodium adsorption ratio, salinity, reaction, and the cation-exchange capacity, which is used to estimate the capacity of a soil to adsorb heavy metals. Permanently frozen soils are not suitable for disposal of wastewater by irrigation.

See the National Soil Handbook, September 1992, Part 620, for criteria used in rating soils for sanitary facilities and waste management.

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
033CP: Clairemont-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
033OA: Obaro-----	100	Somewhat limited Depth to bedrock	0.10	Somewhat limited Depth to bedrock	0.10	Very limited Too steep for surface application	1.00
		Slope	0.04	Slope	0.04	Too steep for sprinkler application	0.22
						Depth to bedrock	0.10
033OB: Obaro-----	75	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Too steep for surface application	1.00
		Depth to bedrock	0.10	Depth to bedrock	0.10	Too steep for sprinkler application	0.97
Rock Outcrop-----	25	Not rated		Not rated		Depth to bedrock	0.10
						Not rated	
033QR: Quinlan-----	55	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Droughty	1.00	Droughty	1.00	Droughty	1.00
		Slope	0.37	Slope	0.37	Too steep for surface application	1.00
						Too steep for sprinkler application	0.59
Woodward-----	45	Somewhat limited Depth to bedrock	0.46	Somewhat limited Depth to bedrock	0.46	Very limited Too steep for surface application	1.00
		Slope	0.37	Slope	0.37	Too steep for sprinkler application	0.59
		Droughty	0.15	Droughty	0.15	Depth to bedrock	0.46
						Droughty	0.15
033QT: Quinlan-----	55	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Slope	1.00	Slope	1.00	Too steep for surface application	1.00
		Droughty	1.00	Droughty	1.00	Too steep for sprinkler application	1.00
						Droughty	1.00
Woodward-----	45	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Too steep for surface application	1.00
		Depth to bedrock	0.46	Depth to bedrock	0.46	Too steep for sprinkler application	1.00
		Droughty	0.15	Droughty	0.15	Depth to bedrock	0.46
						Droughty	0.15
033YE: Yahola-----	100	Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Somewhat limited Flooding	0.60
		Filtering capacity	0.00	Filtering capacity	0.00	Filtering capacity	0.00
077BP: Woodward-----	65	Somewhat limited Droughty	0.71	Somewhat limited Droughty	0.71	Very limited Too steep for surface application	1.00
		Depth to bedrock	0.42	Depth to bedrock	0.42	Droughty	0.71
		Slope	0.37	Slope	0.37	Too steep for sprinkler application	0.59
						Depth to bedrock	0.42

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Port-----	35	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
077FU: Farnum-----	100	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
077GE: Gerlane-----	100	Somewhat limited Flooding Filtering capacity	0.60 0.00	Very limited Flooding Filtering capacity	1.00 0.00	Somewhat limited Flooding Filtering capacity	0.60 0.00
077MC: Minco-----	100	Not limited		Not limited		Not limited	
077MN: Minco-----	100	Not limited		Not limited		Not limited	
077MO: Minco-----	100	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
077QU: Quinlan-----	100	Very limited Depth to bedrock Droughty	1.00 1.00	Very limited Droughty Depth to bedrock	1.00 1.00	Very limited Droughty Depth to bedrock Too steep for surface application	1.00 1.00 0.31
077SE: Shellabarger-----	100	Somewhat limited Too acid	0.11	Somewhat limited Too acid	0.42	Somewhat limited Too acid	0.42
077WE: Quinlan-----	50	Very limited Depth to bedrock Droughty	1.00 1.00	Very limited Droughty Depth to bedrock	1.00 1.00	Very limited Droughty Depth to bedrock	1.00 1.00
Woodward-----	50	Somewhat limited Depth to bedrock Droughty	0.90 0.71	Somewhat limited Depth to bedrock Droughty	0.90 0.71	Somewhat limited Depth to bedrock Droughty	0.90 0.71
095AC: Albion-----	100	Very limited Filtering capacity Too acid	1.00 0.03	Very limited Filtering capacity Too acid	1.00 0.14	Very limited Filtering capacity Too steep for surface application	1.00 0.31
		Droughty	0.00	Droughty	0.00	Too acid Droughty	0.14 0.00
095DA: Dillwyn-----	60	Very limited Filtering capacity Depth to saturated zone Leaching limitation Droughty	1.00 1.00 0.45 0.21	Very limited Filtering capacity Depth to saturated zone Droughty	1.00 1.00 0.21	Very limited Filtering capacity Depth to saturated zone Droughty	1.00 1.00 0.21
Plevna-----	40	Very limited Flooding Depth to saturated zone Runoff limitation Filtering capacity	1.00 1.00 0.40 0.00	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 0.00	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 0.00
095FA: Farnum-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00
095PB: Pratt-----	100	Very limited Filtering capacity Leaching limitation	1.00 0.45	Very limited Filtering capacity	1.00	Very limited Filtering capacity Too steep for surface application Too steep for sprinkler application	1.00 0.66 0.00

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
095SB: Shellabarger-----	100	Somewhat limited Too acid	0.11	Somewhat limited Too acid	0.42	Somewhat limited Too acid	0.42
095WA: Waldeck-----	100	Very limited Filtering capacity Flooding	1.00 0.60	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Filtering capacity Flooding	1.00 0.60
		Depth to saturated zone	0.43	Depth to saturated zone	0.43	Depth to saturated zone	0.43
097LN: Lincoln-----	100	Very limited Filtering capacity Droughty	1.00 0.90	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Filtering capacity Droughty	1.00 0.90
		Flooding Leaching limitation	0.60 0.45	Droughty	0.90	Flooding	0.60
097QW: Quinlan-----	55	Very limited Depth to bedrock Droughty Slope	1.00 1.00 1.00	Very limited Depth to bedrock Droughty Slope	1.00 1.00 1.00	Very limited Depth to bedrock Droughty Too steep for surface application	1.00 1.00 1.00
						Too steep for sprinkler application	1.00
Woodward-----	45	Somewhat limited Slope	0.84	Somewhat limited Slope	0.84	Very limited Too steep for surface application	1.00
		Depth to bedrock	0.71	Depth to bedrock	0.71	Too steep for sprinkler application	0.89
		Droughty	0.38	Droughty	0.38	Depth to bedrock Droughty	0.71 0.38
097WA: Waldeck-----	100	Very limited Filtering capacity Flooding	1.00 0.60	Very limited Flooding Filtering capacity	1.00 1.00	Very limited Filtering capacity Flooding	1.00 0.60
		Depth to saturated zone	0.43	Depth to saturated zone	0.43	Depth to saturated zone	0.43
151AB: Albion-----	100	Very limited Filtering capacity Too acid	1.00 0.03	Very limited Filtering capacity Too acid	1.00 0.14	Very limited Filtering capacity Too acid Too steep for surface application	1.00 0.14 0.00
151AO: Albion-----	100	Very limited Filtering capacity Too acid	1.00 0.03	Very limited Filtering capacity Too acid	1.00 0.14	Very limited Filtering capacity Too steep for surface application Too acid	1.00 0.31 0.14
151AS: Albion-----	50	Very limited Filtering capacity Slope Too acid	1.00 0.37 0.03	Very limited Filtering capacity Slope Too acid	1.00 0.37 0.14	Very limited Too steep for surface application Filtering capacity Too steep for sprinkler application Too acid	1.00 1.00 0.59 0.14
Shellabarger-----	50	Somewhat limited Slope	0.37	Somewhat limited Too acid	0.42	Very limited Too steep for surface application	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
151BC: Blanket-----	100	Too acid	0.11	Slope	0.37	Too steep for sprinkler application Too acid	0.59 0.42
		Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability Too steep for surface application	0.22 0.00
151CA: Carwile-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Restricted permeability	1.00	Restricted permeability	1.00	Restricted permeability	1.00
		Runoff limitation Too acid	0.40 0.02	Too acid	0.07	Too acid	0.07
151CK: Case-----	60	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Too steep for surface application Too steep for sprinkler application	1.00 0.59
Clark-----	40	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Too steep for surface application Too steep for sprinkler application	1.00 0.39
151CM: Clark-----	100	Not limited		Not limited		Somewhat limited Too steep for surface application	0.00
151CN: Clark-----	100	Not limited		Not limited		Not limited	
151CO: Clark-----	70	Not limited		Not limited		Not limited	
Ost-----	30	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
151FA: Farnum-----	100	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
151KP: Kanza-----	50	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Droughty Runoff limitation	0.82 0.40	Droughty Too acid	0.82 0.14	Droughty Too acid	0.82 0.14
Plevna-----	50	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
		Runoff limitation	0.40	Filtering capacity	0.00	Filtering capacity	0.00
		Filtering capacity	0.00				
151OS: Ost-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability Too steep for surface application	0.22 0.00
151PM: Pratt-----	100	Very limited		Very limited		Very limited	

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
151PN: Pratt-----	100	Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Leaching	0.45			Too steep for surface application	0.66
		Limitation				Too steep for sprinkler application	0.00
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
151PO: Pratt-----	60	Leaching	0.45	Slope	0.16	Filtering capacity	1.00
		Limitation	0.16			Too steep for sprinkler application	0.39
		Slope					
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
Carwile-----	40	Leaching	0.45			Too steep for surface application	0.66
		Limitation				Too steep for sprinkler application	0.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Restricted permeability	1.00	Restricted permeability	1.00	Restricted permeability	1.00
151SA: Albion-----	70	Runoff limitation	0.40	Too acid	0.07	Too acid	0.07
		Too acid	0.02				
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
		Slope	0.37	Slope	0.37	Filtering capacity	1.00
Kaski-----	30	Too acid	0.03	Too acid	0.14	Too steep for sprinkler application	0.59
		Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Too acid	0.14
						Somewhat limited Flooding	0.60
151SE: Shellabarger-----	100	Too acid	0.11	Somewhat limited Too acid	0.42	Somewhat limited Too acid	0.42
						Too steep for surface application	0.00
1439: Crisfield-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Too acid	0.37	Too acid	0.96	Too acid	0.96
		Depth to saturated zone	0.18	Flooding	0.40	Depth to saturated zone	0.18
		Droughty	0.11	Depth to saturated zone Droughty	0.18	Droughty	0.11
Ad: Albion-----	60				0.11		
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Too acid	0.03	Too acid	0.14	Too acid	0.14
						Too steep for surface application	0.00
Shellabarger-----	40	Somewhat limited Too acid	0.11	Somewhat limited Too acid	0.42	Somewhat limited Too acid	0.42
						Too steep for surface application	0.00

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ae: Albion-----	55	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
		Slope	0.16	Slope	0.16	Filtering capacity	1.00
		Too acid	0.03	Too acid	0.14	Too steep for sprinkler application	0.39
						Too acid	0.14
Shellabarger-----	45	Somewhat limited Slope	0.16	Somewhat limited Too acid	0.42	Very limited Too steep for surface application	1.00
		Too acid	0.11	Slope	0.16	Too acid	0.42
						Too steep for sprinkler application	0.39
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
AN: Albion-----	65	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
		Slope	0.16	Slope	0.16	Filtering capacity	1.00
		Too acid	0.03	Too acid	0.14	Too steep for sprinkler application	0.39
						Too acid	0.14
Shellabarger-----	35	Somewhat limited Slope	0.16	Somewhat limited Too acid	0.42	Very limited Too steep for surface application	1.00
		Too acid	0.11	Slope	0.16	Too acid	0.42
						Too steep for sprinkler application	0.39
As: Clairemont-----	100	Very limited Flooding Salinity	1.00 0.50	Very limited Flooding Salinity	1.00 1.00	Very limited Flooding Salinity	1.00 1.00
At: Attica-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity Too steep for surface application	0.00 0.00
Ba: Blanket-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
Bb: Blanket-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
Bc: Blanket-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
Bf: Quinlan-----	60	Very limited Depth to bedrock Droughty Slope	1.00 1.00 1.00	Very limited Depth to bedrock Droughty Slope	1.00 1.00 1.00	Very limited Depth to bedrock Droughty Too steep for surface application Too steep for sprinkler application	1.00 1.00 1.00 1.00
Clairemont-----	40	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ca: Canadian-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Flooding Filtering capacity	0.40 0.00	Somewhat limited Filtering capacity	0.00
Cc: Case-----	50	Not limited		Not limited		Somewhat limited Too steep for surface application	0.08
Clark-----	50	Not limited		Not limited		Somewhat limited Too steep for surface application	0.08
Cd: Clairemont-----	100	Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Somewhat limited Flooding	0.60
Cf: Clairemont-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Ck: Clark-----	100	Not limited		Not limited		Not limited	
Fa: Farnum-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00
Fm: Farnum-----	100	Not limited		Not limited		Not limited	
Fr: Farnum-----	100	Not limited		Not limited		Not limited	
Fu: Farnum-----	100	Not limited		Not limited		Not limited	
Ga: Grant-----	100	Not limited		Not limited		Not limited	
Gb: Grant-----	100	Not limited		Not limited		Not limited	
Gc: Grant-----	100	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
GRP: Gravel Pits-----	100	Not rated		Not rated		Not rated	
INT: Aquolls-----	100	Very limited Depth to saturated zone Low adsorption Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Low adsorption Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Low adsorption Ponding	1.00 1.00 1.00
Ka: Kanza-----	100	Very limited Flooding Depth to saturated zone Filtering capacity Droughty Runoff limitation	1.00 1.00 1.00 0.48 0.40	Very limited Flooding Depth to saturated zone Filtering capacity Droughty Too acid	1.00 1.00 1.00 0.48 0.14	Very limited Flooding Depth to saturated zone Filtering capacity Droughty Too acid	1.00 1.00 1.00 0.48 0.14
Kf: Kingfisher-----	100	Somewhat limited Depth to bedrock Restricted permeability Droughty	0.65 0.30 0.04	Somewhat limited Depth to bedrock Restricted permeability Droughty	0.65 0.22 0.04	Somewhat limited Depth to bedrock Restricted permeability Droughty	0.65 0.22 0.04
Kv: Kingfisher-----	70	Somewhat limited Depth to bedrock Restricted permeability Droughty	0.65 0.30 0.04	Somewhat limited Depth to bedrock Restricted permeability Droughty	0.65 0.22 0.04	Somewhat limited Depth to bedrock Restricted permeability Droughty	0.65 0.22 0.04
Vernon-----	30	Very limited Restricted permeability Depth to bedrock	1.00 1.00 0.97	Very limited Restricted permeability Depth to bedrock	1.00 1.00 0.97	Very limited Restricted permeability Depth to bedrock	1.00 1.00 0.97

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Kz: Kingfisher-----	60	Droughty Runoff limitation	0.93 0.40	Droughty	0.93	Droughty	0.93
		Somewhat limited Depth to bedrock	0.65	Somewhat limited Depth to bedrock	0.65	Somewhat limited Depth to bedrock	0.65
		Restricted permeability	0.30	Restricted permeability	0.22	Too steep for surface application	0.31
		Droughty	0.04	Droughty	0.04	Restricted permeability	0.22
Vernon-----	40	Droughty	0.04	Droughty	0.04	Droughty	0.04
		Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Depth to bedrock	0.97	Depth to bedrock	0.97	Depth to bedrock	0.97
		Droughty Runoff limitation	0.93 0.40	Droughty	0.93	Droughty Too steep for surface application	0.93 0.31
Ln: Lincoln-----	100	Droughty	0.04	Droughty	0.04	Droughty	0.04
		Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Droughty Leaching limitation	0.94 0.45	Droughty	0.94	Droughty	0.94
LNN: Lincoln-----	100	Droughty	0.94	Droughty	0.94	Droughty	0.94
		Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Droughty	0.96	Droughty	0.96	Droughty	0.96
		Flooding Leaching limitation	0.60 0.45	Flooding Droughty	0.96	Flooding	0.60
Ma: Mangum-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Flooding	0.60	Flooding	1.00	Flooding	0.60
		Runoff limitation	0.40				
Mg: Mangum-----	65	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Runoff limitation	0.40	Flooding	0.40		
		Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Droughty Runoff limitation Salinity	0.64 0.40 0.01	Droughty Flooding	0.64 0.40	Droughty	0.64
Drummond-----	35	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Droughty	0.64	Droughty	0.64	Droughty	0.64
		Runoff limitation	0.40	Flooding	0.40		
		Salinity	0.01				
Mn: Minco-----	100	Not limited		Not limited		Not limited	
Na: Naron-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00
Nb: Naron-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00
Os: Ost-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
Ot: Ost-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
Pa: Pond Creek-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
		Too acid	0.02	Too acid	0.07	Too acid	0.07
Pd: Pond Creek-----	100	Somewhat limited		Somewhat limited		Somewhat limited	

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ph: Dale-----	100	Restricted permeability	0.30	Restricted permeability	0.22	Restricted permeability	0.22
		Too acid	0.02	Too acid	0.07	Too acid	0.07
Pk: Buttermilk-----	100	Not limited		Somewhat limited Flooding	0.40	Not limited	
		Somewhat limited Salinity	0.01	Somewhat limited Flooding Salinity	0.40 0.13	Somewhat limited Salinity	0.13
Ps: Pratt-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
		Leaching limitation	0.45	Slope	0.00	Too steep for surface application	1.00
		Slope	0.00			Too steep for sprinkler application	0.10
Pt: Pratt-----	70	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
		Leaching limitation	0.45	Slope	0.16	Filtering capacity	1.00
		Slope	0.16			Too steep for sprinkler application	0.39
Tivoli-----	30	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
		Droughty	0.99	Droughty	0.99	Filtering capacity	1.00
		Leaching limitation	0.45	Slope	0.16	Droughty	0.99
		Slope	0.16			Too steep for sprinkler application	0.39
Qn: Quinlan-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Droughty	1.00	Droughty	1.00	Droughty	1.00
Qw: Quinlan-----	60	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
		Droughty	1.00	Droughty	1.00	Droughty	1.00
		Slope	0.16	Slope	0.16	Too steep for surface application	1.00
						Too steep for sprinkler application	0.39
Woodward-----	40	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Too steep for surface application	1.00
						Too steep for sprinkler application	0.39
Rb: Knoco-----	70	Very limited Restricted permeability	1.00	Very limited Droughty	1.00	Very limited Droughty	1.00
		Depth to bedrock	1.00	Restricted permeability	1.00	Restricted permeability	1.00
		Droughty	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Depth to dense layer	1.00	Slope	1.00	Too steep for surface application	1.00
		Slope	1.00	Sodium content	1.00	Too steep for sprinkler application	1.00
Rock Outcrop-----	30	Not rated		Not rated		Not rated	

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Sb: Shellabarger-----	100	Somewhat limited Too acid	0.11	Somewhat limited Too acid	0.42	Somewhat limited Too acid Too steep for surface application	0.42 0.31
SBB: Shellabarger-----	100	Somewhat limited Too acid	0.11	Somewhat limited Too acid	0.42	Somewhat limited Too acid	0.42
Sc: Shellabarger-----	100	Somewhat limited Too acid	0.11	Somewhat limited Too acid	0.42	Somewhat limited Too acid Too steep for surface application	0.42 0.31
Tv: Tivoli-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00	Very limited Too steep for surface application	1.00
		Droughty	1.00	Droughty	1.00	Filtering capacity	1.00
		Slope Leaching limitation	0.84 0.45	Slope	0.84	Droughty Too steep for sprinkler application	1.00 0.89
Vn: Vernon-----	100	Very limited Restricted permeability Depth to bedrock Droughty Runoff limitation	1.00 0.97 0.93 0.40	Very limited Restricted permeability Depth to bedrock Droughty	1.00 0.97 0.93	Very limited Restricted permeability Depth to bedrock Droughty Too steep for surface application	1.00 0.97 0.93 0.08
Vr: Vernon-----	100	Very limited Restricted permeability Depth to bedrock	1.00 0.97	Very limited Restricted permeability Depth to bedrock	1.00 0.97	Very limited Restricted permeability Too steep for surface application	1.00 1.00
		Droughty Runoff limitation Slope	0.93 0.40 0.16	Droughty Slope	0.93 0.16	Depth to bedrock Droughty Too steep for sprinkler application	0.97 0.93 0.39
Vs: Knoco-----	50	Very limited Restricted permeability Depth to bedrock	1.00 1.00	Very limited Droughty	1.00	Very limited Droughty	1.00
		Droughty Depth to dense layer Sodium content	1.00 1.00 1.00	Restricted permeability Depth to bedrock Sodium content	1.00 1.00 1.00	Restricted permeability Depth to bedrock Sodium content	1.00 1.00 1.00
						Too steep for surface application	0.00
Vernon-----	50	Very limited Restricted permeability Droughty Depth to bedrock Runoff limitation	1.00 0.93 0.42 0.40	Very limited Restricted permeability Droughty Depth to bedrock	1.00 0.93 0.42	Very limited Restricted permeability Droughty Depth to bedrock Too steep for surface application	1.00 0.93 0.42 0.00
W: Water-----	100	Not rated		Not rated		Not rated	
Wa: Waldeck-----	100	Very limited Filtering capacity	1.00	Very limited Flooding	1.00	Very limited Filtering capacity	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued
Barber County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Wo: Woodward----- Quinlan-----	70	Flooding	0.60	Filtering capacity	1.00	Flooding	0.60
		Depth to saturated zone	0.43	Depth to saturated zone	0.43	Depth to saturated zone	0.43
	30	Not limited		Not limited		Not limited	
		Very limited Depth to bedrock Droughty	1.00 1.00	Very limited Depth to bedrock Droughty	1.00 1.00	Very limited Depth to bedrock Droughty	1.00 1.00
Ws: Woodward-----	65	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
	35	Very limited Depth to bedrock Droughty	1.00 1.00	Very limited Depth to bedrock Droughty	1.00 1.00	Very limited Depth to bedrock Droughty Too steep for surface application	1.00 1.00 0.31
Ya: Yahola-----	100	Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Somewhat limited Flooding	0.60
		Filtering capacity	0.00	Filtering capacity	0.00	Filtering capacity	0.00
Ze: Zenda-----	100	Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Somewhat limited Flooding	0.60
		Depth to saturated zone	0.43	Depth to saturated zone	0.43	Depth to saturated zone	0.43

WIN-PST SPISP II
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL_KS Sort Order: MUSYM

Barber County, Kansas: KS007

							SPISP II Ratings		
							Leaching	Solution	Adsorbed
MUSYM/SEQ#	COMPONENT/TEXTURE/MU%	HYD	KFACT	SURFACE DEPTH	% OM	(SLP)	Runoff (SSRP)	Runoff (SARP)	
033CP 1	CLAIREMONT L 100%	B	0.43	15"	1.0%	I	I	I	
033OA 1	OBARO SICL 100%	B	0.43	35"	0.5%	I	I	I	
033OB 1	OBARO SICL 75%	B	0.43	35"	0.5%	I	I	I	
033OB 2	ROCK OUTCROP 25%	D	0.00	0"	0.0%	V	H	I (s)	
033QR 1	QUINLAN L 55%	C	0.37	14"	0.5%	L	H	H	
033QR 2	WOODWARD L 45%	B	0.37	30"	1.3%	I	I	I	
033QT 1	QUINLAN L 55%	C	0.37	14"	0.5%	L	H	H (s)	
033QT 2	WOODWARD L 45%	B	0.37	30"	1.3%	I	I	H (s)	
033YE 1	YAHOLA FSL 100%	B	0.20	12"	0.8%	H	I	I	
077BP 1	WOODWARD SIL 65%	B	0.37	24"	1.3%	I	I	H (s)	
077BP 2	PORT SIL 35%	B	0.37	27"	2.0%	L	I	I	
077CH 1	CRISFIELD FSL 100%	B	0.20	18"	1.5%	I	I	I	
077FU 1	FARNUM L 100%	B	0.28	11"	2.0%	I	I	I	
077GE 1	GERLANE FSL 100%	B	0.20	17"	2.0%	H (w)	I	I	
077MC 1	MINCO SIL 100%	B	0.37	42"	2.0%	L	I	I	
077MN 1	MINCO SIL 100%	B	0.37	42"	2.0%	L	I	I	
077MO 1	MINCO SIL 100%	B	0.37	42"	2.0%	L	I	I	
077QU 1	QUINLAN L 100%	C	0.37	9"	0.5%	L	H	H	
077SE 1	SHELLABARGER FSL 100%	B	0.20	13"	1.5%	I	I	I	
077WE 1	QUINLAN L 50%	C	0.37	9"	0.5%	L	H	H	
077WE 2	WOODWARD L 50%	B	0.37	24"	1.3%	I	I	I	
095AC 1	ALBION SL 100%	B	0.20	8"	1.5%	H	I	I	
095DA 1	DILLWYN LFS 60%	A	0.17	8"	1.0%	H (w)	L	L	
095DA 2	PLEVNA FSL 40%	D	0.20	11"	2.5%	H (w)	H	H	
095FA 1	FARNUM SL 100%	B	0.20	16"	1.5%	I	I	I	
095SB 1	SHELLABARGER SL 100%	B	0.20	10"	1.5%	H	I	I	
095WA 1	WALDECK FSL 100%	C	0.20	12"	1.5%	H (w)	H	I	
097LN 1	LINCOLN SL 100%	A	0.20	8"	0.5%	H	L	L	

WIN-PST SPISP II
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL_KS Sort Order: MUSYM

Barber County, Kansas: KS007

097QW 1	QUINLAN L 55%	C	0.37	15"	0.5% L	H	H (s)
097QW 2	WOODWARD L 45%	B	0.37	27"	1.3% I	I	H (s)
097WA 1	WALDECK L 100%	C	0.28	14"	2.0% H (w)	H	H
151AB 1	ALBION SL 100%	B	0.20	8"	1.5% H	I	I
151AO 1	ALBION SL 100%	B	0.20	8"	1.5% H	I	I
151AS 1	ALBION SL 50%	B	0.20	8"	1.5% H	I	I
151AS 2	SHELLABARGER FSL 50%	B	0.20	11"	1.5% I	I	I
151BC 1	BLANKET SICL 100%	C	0.37	13"	2.0% L	H	H
151CA 1	CARWILE FSL 100%	D	0.24	12"	2.0% H (w)	H	H
151CK 1	CASE CL 60%	B	0.32	6"	1.3% H	I	I
151CK 2	CLARK CL 40%	B	0.28	8"	1.5% I	I	I
151CM 1	CLARK CL 100%	B	0.28	8"	1.5% I	I	I
151CN 1	CLARK FSL 100%	B	0.20	8"	1.5% H	I	I
151CO 1	CLARK CL 70%	B	0.28	8"	1.5% I	I	I
151CO 2	OST CL 30%	B	0.32	9"	2.0% I	I	I
151FA 1	FARNUM CL 100%	B	0.28	7"	2.0% I	I	I
151KP 1	KANZA LFS 50%	D	0.17	11"	2.0% H (w)	H	H
151KP 2	PLEVNA FSL 50%	D	0.20	10"	2.5% H (w)	H	H
151OS 1	OST CL 100%	B	0.32	9"	2.0% I	I	I
151PM 1	PRATT LFS 100%	A	0.17	10"	0.8% H	L	L
151PN 1	PRATT LFS 100%	A	0.17	10"	0.8% H	L	L
151PO 1	PRATT LFS 60%	A	0.17	10"	0.8% H	L	L
151PO 2	CARWILE FSL 40%	D	0.24	12"	2.0% H (w)	H	H
151SA 1	ALBION FSL 70%	B	0.20	8"	1.5% H	I	I
151SA 2	KASKI L 30%	B	0.28	26"	2.0% L	I	I
151SE 1	SHELLABARGER FSL 100%	B	0.20	11"	1.5% I	I	I
Ad 1	ALBION SL 60%	B	0.20	8"	1.5% H	I	I
Ad 2	SHELLABARGER SL 40%	B	0.20	14"	1.5% I	I	I
Ae 1	ALBION SL 55%	B	0.20	8"	1.5% H	I	I
Ae 2	SHELLABARGER SL 45%	B	0.20	14"	1.5% I	I	I

WIN-PST SPISP II
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL_KS Sort Order: MUSYM

Barber County, Kansas: KS007

AED 1	ARENTS, EARTHEN DAM 100%		0.00	0"	0.0% ?	?	?
AN 1	ALBION SL 65%	B	0.20	8"	1.5% H	I	I
AN 2	SHELLABARGER SL 35%	B	0.20	11"	1.5% I	I	I
As 1	CLAIREMONT SIL 100%	B	0.43	8"	1.0% H	I	I
At 1	ATTICA LFS 100%	B	0.17	12"	0.8% H	I	I
Ba 1	BLANKET SIL 100%	C	0.37	9"	2.0% L	H	H
Bb 1	BLANKET SIL 100%	C	0.37	9"	2.0% L	H	H
Bc 1	BLANKET SICL 100%	C	0.37	9"	2.0% L	H	H
Bf 1	QUINLAN L 60%	C	0.37	14"	0.5% L	H	H (s)
Bf 2	CLAIREMONT SIL 40%	B	0.43	14"	1.0% I	I	I
Ca 1	CANADIAN FSL 100%	B	0.20	18"	2.0% I	I	I
Cc 1	CASE CL 50%	B	0.32	6"	1.3% H	I	I
Cc 2	CLARK CL 50%	B	0.28	10"	1.5% I	I	I
Cd 1	CLAIREMONT SIL 100%	B	0.43	14"	1.0% I	I	I
Cf 1	CLAIREMONT SIL 100%	B	0.43	14"	1.0% I	I	I
Ck 1	CLARK CL 100%	B	0.28	10"	1.5% I	I	I
Fa 1	FARNUM FSL 100%	B	0.20	9"	1.5% H	I	I
Fm 1	FARNUM L 100%	B	0.28	9"	2.0% I	I	I
Fr 1	FARNUM L 100%	B	0.28	9"	2.0% I	I	I
Fu 1	FARNUM CL 100%	B	0.28	9"	2.0% I	I	I
Ga 1	GRANT SIL 100%	B	0.37	13"	2.0% I	I	I
Gb 1	GRANT SIL 100%	B	0.37	13"	2.0% I	I	I
Gc 1	GRANT SIL 100%	B	0.37	13"	2.0% I	I	I
GRP 1	GRAVEL PITS 100%		0.00	0"	0.0% ?	?	?
INT 1	AQUOLLS VAR 100%	C	0.00	60"	0.0% ?	H	?
Ka 1	KANZA LFS 100%	D	0.17	7"	2.0% H (w)	H	H
Kf 1	KINGFISHER SIL 100%	B	0.37	9"	2.0% I	I	I
Kv 1	KINGFISHER SIL 70%	B	0.37	9"	2.0% I	I	I
Kv 2	VERNON CL 30%	D	0.37	5"	0.8% V	H	H
Kz 1	KINGFISHER SIL 60%	B	0.37	9"	2.0% I	I	I

WIN-PST SPISP II
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL_KS Sort Order: MUSYM

Barber County, Kansas: KS007

Kz 2	VERNON CL 40%	D	0.37	5"	0.8% V	H	H
Ln 1	LINCOLN FSL 100%	A	0.20	6"	0.5% H	L	L
LNN 1	LINCOLN LS 100%	A	0.17	10"	0.5% H	L	L
Ma 1	MANGUM C 100%	D	0.32	10"	1.5% V	H	H
Mg 1	MANGUM C 65%	D	0.32	10"	1.5% V	H	H
Mg 2	DRUMMOND SICL 35%	D	0.43	8"	0.8% H (w)	H	H
Mn 1	MINCO SIL 100%	B	0.37	27"	2.0% L	I	I
Na 1	NARON FSL 100%	B	0.20	12"	2.0% I	I	I
Nb 1	NARON FSL 100%	B	0.20	12"	2.0% I	I	I
Os 1	OST CL 100%	B	0.32	8"	2.0% I	I	I
Ot 1	OST CL 100%	B	0.32	8"	2.0% I	I	I
Pa 1	POND CREEK SIL 100%	B	0.37	11"	2.0% I	I	I
Pd 1	POND CREEK SIL 100%	B	0.37	11"	2.0% I	I	I
Ph 1	PORT SIL 100%	B	0.37	20"	2.0% I	I	I
Pk 1	BUTTERMILK SIL 100%	B	0.32	20"	2.0% I	I	I
PR 1	PRATT LFS 100%	A	0.17	12"	0.8% H	L	L
Ps 1	PRATT LFS 100%	A	0.17	7"	0.8% H	L	L
Pt 1	PRATT LFS 70%	A	0.17	7"	0.8% H	L	L
Pt 2	TIVOLI LFS 30%	A	0.17	5"	0.5% H	L	L
Qn 1	QUINLAN L 100%	C	0.37	14"	0.5% L	H	H
Qw 1	QUINLAN L 60%	C	0.37	14"	0.5% L	H	H
Qw 2	WOODWARD L 40%	B	0.37	40"	1.3% L	I	I
Rb 1	KNOCO C 70%	D	0.37	9"	1.0% V	H	H (s)
Rb 2	ROCK OUTCROP UWB 30%	D	0.00	10"	0.0% V	H	I (s)
Sb 1	SHELLABARGER SL 100%	B	0.20	14"	1.5% I	I	I
SBB 1	SHELLABARGER FSL 100%	B	0.20	11"	1.5% I	I	I
Sc 1	SHELLABARGER SL 100%	B	0.20	14"	1.5% I	I	I
Tv 1	TIVOLI FS 100%	A	0.17	5"	0.5% H	L	I (s)
Vn 1	VERNON CL 100%	D	0.37	5"	0.8% V	H	H
Vr 1	VERNON CL 100%	D	0.37	5"	0.8% V	H	H

WIN-PST SPISP II
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL_KS Sort Order: MUSYM

Barber County, Kansas: KS007

Vs 1	KNOCO C 50%	D	0.37	9"	1.0% V	H	H
Vs 2	VERNON CL 50%	D	0.37	5"	0.8% V	H	H
W 1	WATER 100%		0.00	0"	0.0% ?	?	?
Wa 1	WALDECK SL 100%	C	0.20	12"	1.5% H (w)	H	I
Wo 1	WOODWARD L 70%	B	0.37	40"	1.3% L	I	I
Wo 2	QUINLAN L 30%	C	0.37	14"	0.5% L	H	H
Ws 1	WOODWARD L 65%	B	0.37	40"	1.3% L	I	I
Ws 2	QUINLAN L 35%	C	0.37	14"	0.5% L	H	H
Ya 1	YAHOLA SL 100%	B	0.20	14"	0.8% H	I	I
Ze 1	ZENDA CL 100%	C	0.28	20"	2.0% H (w)	H	H

(.\REPORTS\SOILS.TXT generated on 12/12/01 at 12:11:15)

H -- High
I -- Intermediate
L -- Low
V -- Very Low

Conditions that affect ratings:

- m -- There are macropores in the surface horizon deeper than 24"
- w -- The high water table comes within 24" of the surface during the growing season
- s -- The field slope is greater than 15%

SPISP II S-Ratings:

- SLP -- Soil Leaching Potential
- SSRP -- Soil Solution Runoff Potential
- SARP -- Soil Adsorbed Runoff Potential

In this section, hydric soils are defined and described and the hydric soils in the survey area are listed. The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin and others, 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for each of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part (Federal Register, 1994). These soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 1995). These criteria are used to identify a phase of a soil series that normally is associated with wetlands. The criteria used are selected estimated soil properties that are described in "Soil Taxonomy" (USDA, 1999) and "Keys to Soil Taxonomy" (USDA, 1998) and in the "Soil Survey Manual" (USDA, 1993).

If soils are wet enough for a long enough period to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils in this survey area are specified in "Field Indicators of Hydric Soils in the United States" (Hurt and others, 1996).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

Map units in the Hydric Soil Interpretations table meet the definition of hydric soils and, in addition, have at least one of the hydric soil indicators. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; Hurt and others, 1996).

Map units that are made up of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units made up of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform.

These map units, in general, do not meet the definition of hydric soils because they do not have one of the hydric soil indicators. A portion of these map units, however, may include hydric soils. Onsite investigation is recommended to determine whether hydric soils occur and the location of the included hydric soils.

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Barber County, Kansas

PAGE 2 of 8

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
033CP: CLAIREMONT LOAM, CHANNELED	CLAIREMONT	No	flood plain	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3,4	YES	YES	NO
033OA: OBARO SILTY CLAY LOAM, 5 TO 12 PERCENT SLOPES	OBARO	No	hillslope	---	---	---	---
033OB: OBARO-ROCK OUTCROP COMPLEX, 10 TO 30 PERCENT SLOPES	OBARO	No	hillslope	---	---	---	---
	ROCK OUTCROP	Unranked	---	---	---	---	---
033QR: QUINLAN-WOODWARD LOAMS, 6 TO 15 PERCENT SLOPES	QUINLAN	No	hillslope	---	---	---	---
	WOODWARD	No	hillslope	---	---	---	---
033QT: QUINLAN-WOODWARD LOAMS, 15 TO 30 PERCENT SLOPES	QUINLAN	No	hillslope	---	---	---	---
	WOODWARD	No	hillslope	---	---	---	---
033YE: YAHOLA FINE SANDY LOAM, OCCASIONALLY FLOODED	YAHOLA	No	flood plain	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B2,2B3,2A	YES	NO	NO
077BP: WOODWARD-PORT COMPLEX, 0 TO 20 PERCENT SLOPES	WOODWARD	No	hillslope	---	---	---	---
	PORT Unnamed wet soils	Unranked Yes	terrace depression	--- 2B3,3,2A,4	--- YES	--- YES	--- YES
077FU: FARNUM LOAM, 3 TO 6 PERCENT SLOPES	FARNUM	No	paleoterrace	---	---	---	---
077GE: GERLANE FINE SANDY LOAM, OCCASIONALLY FLOODED	GERLANE	No	terrace	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3,2B2,2A	YES	NO	NO
077MC: MINCO SILT LOAM, 0 TO 1 PERCENT SLOPES	MINCO	No	hillslope	---	---	---	---
077MN: MINCO SILT LOAM, 1 TO 3 PERCENT SLOPES	MINCO	No	hillslope	---	---	---	---
077MO: MINCO SILT LOAM, 3 TO 6 PERCENT SLOPES	MINCO	No	hillslope	---	---	---	---
077QU: QUINLAN LOAM, 3 TO 6 PERCENT SLOPES	QUINLAN	No	hillslope	---	---	---	---
077SE: SHELLABARGER FINE SANDY LOAM, 1 TO 3 PERCENT SLOPES	SHELLABARGER	No	paleoterrace	---	---	---	---
077WE: WOODWARD-QUINLAN LOAMS, 1 TO 3 PERCENT SLOPES	QUINLAN	No	hillslope	---	---	---	---
	WOODWARD Unnamed wet soils	No Yes	hillslope drainageway	--- 2A,2B3	--- YES	--- NO	--- NO
095AC: ALBION SANDY LOAM, 3 TO 6 PERCENT SLOPES	ALBION	No	paleoterrace	---	---	---	---
095DA: DILLWYN-PLEVNA COMPLEX, OCCASIONALLY FLOODED	DILLWYN	No	interdune, dune, paleoterrace	---	---	---	---
	PLEVNA Unnamed wet soils	Yes Yes	flood plain depression	2B3,4 2B3,2A,3	YES YES	YES NO	NO YES

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Barber County, Kansas

PAGE 3 of 8

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
095FA: FARNUM SANDY LOAM, 0 TO 2 PERCENT SLOPES	FARNUM	No	paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
	Unnamed wet soils	Yes	depression	2B3,3,2A	YES	NO	YES
095PB: PRATT LOAMY FINE SAND, 1 TO 5 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
	Unnamed wet soils	Yes	depression	2A,2B3,3	YES	NO	YES
095SB: SHELLABARGER SANDY LOAM, 1 TO 3 PERCENT SLOPES	SHELLABARGER	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2B3,2A	YES	NO	NO
095WA: WALDECK FINE SANDY LOAM, OCCASIONALLY FLOODED	WALDECK	No	flood plain	---	---	---	---
	PLEVNA	Yes	flood plain	2B3	YES	NO	NO
	Unnamed wet soils	Yes	depression	3,2B3,2A,4	YES	YES	YES
097LN: LINCOLN SANDY LOAM, OCCASIONALLY FLOODED	LINCOLN	No	flood plain	---	---	---	---
	PLEVNA	Yes	depression	2B3,4	YES	YES	NO
	Unnamed wet soils	Yes	drainageway	2A,2B3,2B2	YES	NO	NO
097QW: QUINLAN-WOODWARD LOAMS, 6 TO 25 PERCENT SLOPES	QUINLAN	No	---	---	---	---	---
	WOODWARD	No	---	---	---	---	---
097WA: WALDECK LOAM, OCCASIONALLY FLOODED	WALDECK	No	flood plain	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B2,2B3,2A	YES	NO	NO
151AB: ALBION SANDY LOAM, 1 TO 4 PERCENT SLOPES	ALBION	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3,2A	YES	NO	NO
151AO: ALBION SANDY LOAM, 3 TO 7 PERCENT SLOPES, ERODED	ALBION	No	paleoterrace	---	---	---	---
151AS: ALBION AND SHELLABARGER SOILS, 7 TO 15 PERCENT SLOPES	ALBION	No	paleoterrace	---	---	---	---
	SHELLABARGER	No	paleoterrace	---	---	---	---
151BC: BLANKET SILTY CLAY LOAM, 1 TO 4 PERCENT SLOPES, ERODED	BLANKET	No	paleoterrace	---	---	---	---
151CA: CARWILE FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES	CARWILE	Yes	depression	2A	YES	NO	NO
151CK: CASE-CLARK COMPLEX, 7 TO 15 PERCENT SLOPES	CASE	No	paleoterrace	---	---	---	---
	CLARK	No	paleoterrace	---	---	---	---
151CM: CLARK CLAY LOAM, 1 TO 4 PERCENT SLOPES	CLARK	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	drainageway	4,3,2A	YES	YES	YES
151CN: CLARK FINE SANDY LOAM, 1 TO 3 PERCENT SLOPES	CLARK	No	paleoterrace	---	---	---	---
151CO: CLARK-OST CLAY LOAMS, 0 TO 1 PERCENT SLOPES	CLARK	No	paleoterrace	---	---	---	---
	OST	No	paleoterrace	---	---	---	---

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Barber County, Kansas

PAGE 4 of 8

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
151FA: FARNUM CLAY LOAM, 3 TO 6 PERCENT SLOPES, ERODED	FARNUM	No	paleoterrace	---	---	---	---
151KP: KANZA-PLEVNA COMPLEX, FREQUENTLY FLOODED	KANZA	Yes	flood plain	2B3	YES	NO	NO
	PLEVNA	Yes	flood plain	2B3,4	YES	YES	NO
151OS: OST CLAY LOAM, 1 TO 4 PERCENT SLOPES	OST	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	depression	3,2B3,2A,4	YES	YES	YES
151PM: PRATT LOAMY FINE SAND, 3 TO 8 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
	Unnamed wet soils	Yes	depression	3,2B3,2A	YES	NO	YES
151PN: PRATT LOAMY FINE SAND, 8 TO 12 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
151PO: PRATT-CARWILE COMPLEX, 0 TO 5 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	3,2A	YES	NO	YES
	Unnamed wet soils	Yes	depression	3,2B3	YES	NO	YES
151SA: ALBION-KASKI COMPLEX, 0 TO 15 PERCENT SLOPES	ALBION	No	paleoterrace	---	---	---	---
	KASKI	No	flood plain	---	---	---	---
151SE: SHELLABARGER FINE SANDY LOAM, 1 TO 4 PERCENT SLOPES	SHELLABARGER	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3,2A	YES	NO	NO
1439: CRISFIELD SANDY LOAM, RARELY FLOODED	CRISFIELD	No	terrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2B3,2A	YES	NO	NO
Ad: ALBION-SHELLABARGER SANDY LOAMS, 2 TO 4 PERCENT SLOPES	ALBION	No	paleoterrace	---	---	---	---
	SHELLABARGER	No	paleoterrace	---	---	---	---
Ae: ALBION AND SHELLABARGER SOILS, 4 TO 15 PERCENT SLOPES	ALBION	No	paleoterrace	---	---	---	---
	SHELLABARGER	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2A,2B3	YES	NO	NO
AED: ARENTS, EARTHEN DAM	ARENTS, EARTHEN DAM	Unranked	---	---	---	---	---
AN: ALBION-SHELLABARGER SANDY LOAMS, 4 TO 15 PERCENT SLOPES	ALBION	No	paleoterrace	---	---	---	---
	SHELLABARGER	No	paleoterrace	---	---	---	---
As: CLAIREMONT SOILS, SALINE, CHanneled	CLAIREMONT	No	flood plain	---	---	---	---
	Unnamed wet soils	Yes	depression	2A,2B3,4	YES	YES	NO
At: ATTICA LOAMY FINE SAND, 1 TO 5 PERCENT SLOPES	ATTICA	No	dune, paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2A,2B3,3	YES	NO	YES
Ba: BLANKET SILT LOAM, 0 TO 1 PERCENT SLOPES	BLANKET	No	paleoterrace	---	---	---	---
Bb: BLANKET SILT LOAM, 1 TO 3 PERCENT SLOPES	BLANKET	No	paleoterrace	---	---	---	---

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Barber County, Kansas

PAGE 5 of 8

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Bc: BLANKET SILTY CLAY LOAM, 1 TO 3 PERCENT SLOPES, ERODED	BLANKET	No	paleoterrace	---	---	---	---
Bf: QUINLAN-CLAIREMONT COMPLEX, 0 TO 50 PERCENT SLOPES	QUINLAN	No	hillslope	---	---	---	---
	CLAIREMONT Unnamed wet soils	No Yes	flood plain drainageway	--- 2A, 2B3	--- YES	--- NO	--- NO
Ca: CANADIAN FINE SANDY LOAM, RARELY FLOODED	CANADIAN	No	flood plain	---	---	---	---
Cc: CASE-CLARK CLAY LOAMS, 2 TO 6 PERCENT SLOPES	CASE	No	paleoterrace	---	---	---	---
	CLARK	No	paleoterrace	---	---	---	---
Cd: CLAIREMONT SILT LOAM, OCCASIONALLY FLOODED	CLAIREMONT	No	flood plain	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2A, 2B3, 4	YES	YES	NO
Cf: CLAIREMONT SOILS, CHANNELED	CLAIREMONT	No	flood plain	---	---	---	---
	Unnamed wet soils	Yes	drainageway	4, 2B3, 2A	YES	YES	NO
Ck: CLARK CLAY LOAM, 0 TO 2 PERCENT SLOPES	CLARK	No	paleoterrace	---	---	---	---
Fa: FARNUM FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES	FARNUM	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2A, 3, 2B3	YES	NO	YES
Fm: FARNUM LOAM, 0 TO 1 PERCENT SLOPES	FARNUM	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2A, 3, 2B3	YES	NO	YES
Fr: FARNUM LOAM, 1 TO 3 PERCENT SLOPES	FARNUM	No	paleoterrace	---	---	---	---
Fu: FARNUM CLAY LOAM, 1 TO 3 PERCENT SLOPES, ERODED	FARNUM	No	paleoterrace	---	---	---	---
Ga: GRANT SILT LOAM, 0 TO 1 PERCENT SLOPES	GRANT	No	terrace	---	---	---	---
Gb: GRANT SILT LOAM, 1 TO 3 PERCENT SLOPES	GRANT	No	terrace	---	---	---	---
Gc: GRANT SILT LOAM, 3 TO 6 PERCENT SLOPES	GRANT	No	terrace	---	---	---	---
GRP: GRAVEL PITS	GRAVEL PITS	Unranked	---	---	---	---	---
INT: AQUOLLS	AQUOLLS	Yes	depression, terrace	3, 2B3	YES	NO	YES
Ka: KANZA SOILS, FREQUENTLY FLOODED	KANZA	Yes	flood plain	2B3	YES	NO	NO
Kf: KINGFISHER SILT LOAM, 1 TO 3 PERCENT SLOPES	KINGFISHER	No	hillslope	---	---	---	---
Kv: KINGFISHER-VERNON COMPLEX, 1 TO 3 PERCENT SLOPES	KINGFISHER	No	hillslope	---	---	---	---
	VERNON	No	hillslope	---	---	---	---
Kz: KINGFISHER-VERNON COMPLEX, 3 TO 6 PERCENT SLOPES	KINGFISHER	No	hillslope	---	---	---	---
	VERNON	No	hillslope	---	---	---	---

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Barber County, Kansas

PAGE 6 of 8

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Ln: LINCOLN SOILS, FREQUENTLY FLOODED	LINCOLN	No	flood plain	---	---	---	---
	KANZA Unnamed wet soils	Yes Yes	flood plain depression	2B2 2A, 2B2, 3, 4, 2B3	YES YES	NO YES	NO YES
LNN: LINCOLN LOAMY SAND, OCCASIONALLY FLOODED	LINCOLN	No	flood plain	---	---	---	---
	KANZA	Yes	depression	2B2	YES	NO	NO
Ma: MANGUM CLAY, OCCASIONALLY FLOODED	MANGUM	No	flood plain	---	---	---	---
	Unnamed wet soils	Yes	depression	3, 2B3, 2A	YES	NO	YES
Mg: MANGUM-DRUMMOND COMPLEX, RARELY FLOODED	MANGUM	No	flood plain	---	---	---	---
	DRUMMOND Unnamed wet soils	No Yes	terrace depression	--- 2A, 3, 2B3	--- YES	--- NO	--- YES
Mn: MINCO SILT LOAM, 0 TO 2 PERCENT SLOPES	MINCO	No	hillslope	---	---	---	---
Na: NARON FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES	NARON	No	dune, paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2A, 3, 2B3	YES	NO	YES
Nb: NARON FINE SANDY LOAM, 1 TO 3 PERCENT SLOPES	NARON	No	dune, paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2A, 2B3, 3, 4	YES	YES	YES
Os: OST CLAY LOAM, 0 TO 1 PERCENT SLOPES	OST	No	paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
Ot: OST CLAY LOAM, 1 TO 3 PERCENT SLOPES	OST	No	paleoterrace	---	---	---	---
Pa: POND CREEK SILT LOAM, 0 TO 1 PERCENT SLOPES	POND CREEK	No	terrace	---	---	---	---
Pd: POND CREEK SILT LOAM, 1 TO 3 PERCENT SLOPES	POND CREEK	No	terrace	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2A, 2B3, 4	YES	YES	NO
Ph: DALE SILT LOAM, RARELY FLOODED	DALE	No	terrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2A, 2B3	YES	NO	NO
Pk: BUTTERMILK SILT LOAM, RARELY FLOODED	BUTTERMILK	No	flood plain	---	---	---	---
	Unnamed wet soils	Yes	depression	2A, 2B3, 4	YES	YES	NO
Ps: PRATT LOAMY FINE SAND, 5 TO 10 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
	Unnamed wet soils	Yes	depression	2A, 2B3, 3, 2B 2	YES	NO	YES
Pt: PRATT-TIVOLI LOAMY FINE SANDS, 5 TO 15 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
	TIVOLI	No	dune, paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2A, 2B3, 2B2, 3	YES	NO	YES
Qn: QUINLAN LOAM, 1 TO 3 PERCENT SLOPES	QUINLAN	No	hillslope	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3, 2A	YES	NO	NO

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Barber County, Kansas

PAGE 7 of 8

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Qw: QUINLAN-WOODWARD COMPLEX, 5 TO 15 PERCENT SLOPES	QUINLAN	No	hillslope	---	---	---	---
	WOODWARD Unnamed wet soils	No Yes	hillslope drainageway	--- 2B3,4,2A	--- YES	--- YES	--- NO
Rb: KNOCO-SHALE OUTCROP COMPLEX, 15 TO 80 PERCENT SLOPES	KNOCO	No	hillslope	---	---	---	---
	ROCK OUTCROP Unnamed wet soils	Unranked Yes	--- drainageway	--- 2A,2B3	--- YES	--- NO	--- NO
Sb: SHELLABARGER SANDY LOAM, 3 TO 6 PERCENT SLOPES	SHELLABARGER	No	paleoterrace	---	---	---	---
SBB: SHELLABARGER FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES	SHELLABARGER	No	paleoterrace	---	---	---	---
Sc: SHELLABARGER SANDY LOAM, 3 TO 6 PERCENT SLOPES, ERODED	SHELLABARGER	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2A,2B3,4	YES	YES	NO
Tv: TIVOLI FINE SAND, 5 TO 20 PERCENT SLOPES	TIVOLI	No	dune, paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2A,2B3,3,2B 2	YES	NO	YES
Vn: VERNON CLAY LOAM, 3 TO 5 PERCENT SLOPES	VERNON	No	hillslope	---	---	---	---
Vr: VERNON CLAY LOAM, 5 TO 15 PERCENT SLOPES	VERNON	No	hillslope	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2A,2B3	YES	NO	NO
Vs: VERNON-KNOCO COMPLEX, 1 TO 5 PERCENT SLOPES	KNOCO	No	hillslope	---	---	---	---
	VERNON	No	hillslope	---	---	---	---
W: WATER	WATER	Unranked	---	---	---	---	---
Wa: WALDECK SANDY LOAM, OCCASIONALLY FLOODED	WALDECK	No	flood plain	---	---	---	---
	KANZA	Yes	flood plain	2B3	YES	NO	NO
Wo: WOODWARD-QUINLAN LOAMS, 0 TO 3 PERCENT SLOPES	WOODWARD	No	hillslope	---	---	---	---
	QUINLAN Unnamed wet soils	No Yes	hillslope drainageway	--- 2A,2B3	--- YES	--- NO	--- NO
Ws: WOODWARD-QUINLAN LOAMS, 3 TO 6 PERCENT SLOPES	WOODWARD	No	hillslope	---	---	---	---
	QUINLAN Unnamed wet soils	No Yes	hillslope drainageway	--- 2A,2B3	--- YES	--- NO	--- NO
Ya: YAHOLA SANDY LOAM, OCCASIONALLY FLOODED	YAHOLA	No	flood plain	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3,3,2A	YES	NO	YES
Ze: ZENDA CLAY LOAM, OCCASIONALLY FLOODED	ZENDA	No	dune, paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2A,2B3	YES	NO	NO

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Barber County, Kansas

PAGE 8 of 8

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria

FOOTNOTE: There may be small areas of included soils or miscellaneous areas that are significant to use and management of the soil; yet are too small to delineate on the soil map at the map's original scale. These may be designated as spot symbols and are defined in the published Soil Survey Report or the USDA-NRCS Technical Guide, Part II.

Areas mapped as water or any map unit that contains one of the following conventional symbols is considered a hydric soil map unit: marshes or swamps; wet spots; depressions; streams, lakes and ponds.

1. All Histosols except Folists, or
2. Soils in Aquic suborders, great groups, or subgroups, Albolls suborder, Aquisalids, Pachic subgroups, or Cumulic subgroups that are:
 - a. Somewhat poorly drained with a water table equal to 0.0 foot (ft) from the surface during the growing season, or
 - b. poorly drained or very poorly drained and have either:
 - (1) water table equal to 0.0 ft during the growing season if textures are coarse sand, sand, or fine sand in all layers within 20 inches (in), or for other soils
 - (2) water table at less than or equal to 0.5 ft from the surface during the growing season if permeability is equal to or greater than 6.0 in/hour (h) in all layers within 20 in, or
 - (3) water table at less than or equal to 1.0 ft from the surface during the growing season if permeability is less than 6.0 in/h in any layer within 20 in, or
3. Soils that are frequently ponded for long duration or very long duration during the growing season, or
4. Soils that are frequently flooded for long duration or very long duration during the growing season.