

<b>Table of Contents</b> <b>Section II – Soil and Site Information</b>
---

	<b>Issue Date</b>	<b>Date of Last Review</b>	<b>Responsible Staff</b>
<i>Use and Explanation of Soil Interpretations</i>			
<i>Explanation of Key Phrases Used in Soil Interpretations</i>			
<b>Soils Legends</b>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<i>* Acreage and Proportionate Extent of the Soils</i>			
<b>Soil Descriptions - Nontechnical</b>	<b>1/02</b>	<b>1/93</b>	<b>SOI</b>
<i>Use and Explanation of Nontechnical Descriptions</i>			
<i>*Nontechnical Soils Description Report</i>			
<b>Soil Descriptions - Technical</b>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<i>*Map Unit Description Report</i>			
<b>Cropland Interpretations - Technical</b>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<i>*Prime Farmland Report</i>			
<i>* Kansas Soil Rating for Plant Growth Index</i>			
<i>*Soil Properties for Conservation Planning</i>			
<b>Rangeland, Grazed Forestland, Native Pastureland Interpretations</b>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<i>*Rangeland Productivity Report</i>			
<i>*Range Site Descriptions</i>			
<b>Forestland Interpretations</b>	<b>1/93</b>	<b>1/93</b>	<b>SOI</b>
<i>Use and Explanation of Forestland Interpretations</i>			
<i>*Woodland Management and Productivity</i>			
<b>Nonagricultural Interpretations</b>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<i>*Building Site Development Report</i>			
<i>*Construction Materials Report</i>			
<b>Recreation Interpretations</b>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<i>*Recreational Interpretations</i>			
<b>Wildlife Interpretations</b>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<i>*Wildlife Interpretations Report</i>			
<b>Pastureland and Hayland Interpretations</b>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<i>*Yields Per Acre of Pasture and Hayland</i>			

	<b>Issue Date</b>	<b>Date of Last Review</b>	<b>Responsible Staff</b>
<b>Mined Land Interpretations</b> <i>Use and Explanation of Mined Land Interpretations</i>	<b>1/93</b>	<b>1/93</b>	<b>SOI</b>
<b>Windbreak Interpretations</b> <i>*Conservation Tree and Shrub Management Report</i>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<b>Engineering Interpretations</b> <i>*Engineering Index Properties</i> <i>*Physical Properties of the Soils</i> <i>*Chemical Properties of the Soils</i> <i>*Water Features</i> <i>*Soil Features</i> <i>*Water Management Report</i>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<b>Waste Disposal Interpretations</b> <i>*Sanitary Facilities Report</i> <i>*Agricultural Waste Management Report</i>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<b>Water Quantity and Quality Interpretations</b> <i>Use and Explanation of Water Quantity and Quality Interpretations</i> <i>*Appendix A – Soils Potential For Surface Loss and Leaching</i> <i>*Appendix B – Pesticide Selected Properties Database</i> <i>*Appendix C – Herbicide Selected Properties Database</i> <i>*Soil-Pesticide Interaction Screening Procedure Worksheet (Blank)</i> <i>*WIN-PST SPISP II Soil Sensitivity to Pesticide Loss Rating Report</i>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<b>Hydric Soil Interpretations</b> <i>Use and Explanation of Hydric Soil Interpretations</i> <i>*Hydric Soils List</i>	<b>1/02</b>	<b>1/02</b>	<b>SOI</b>
<b>HEL Interpretations</b> <i>Use and Explanation of Highly Erodible Land Interpretations</i> <i>*Highly Erodible Lands Report</i> <i>*LS and Supporting Data for 1990 Frozen HEL List</i> <i>*CRP 20 Soil Supporting Data for 1990 Frozen HEL List</i>	<b>7/95</b>	<b>1/00</b>	<b>SOI</b>

*\*County specific computer generated reports.*

## ACREAGE AND PROPORTIONATE EXTENT OF THE SOILS

Gray County, Kansas: Maintenance needed

Map symbol	Soil name	Acres	Percent
055BP	Bridgeport Clay Loam, Rarely Flooded-----	40	*
055LD	Las-Las Animas Complex, Occasionally Flooded-----	9	*
055TF	Valent Fine Sand, 6 To 20 Percent Slopes-----	1,972	0.4
055TV	Valent-Vona Loamy Fine Sands, 3 To 8 Percent Slopes-----	976	0.2
055VO	Vona Loamy Fine Sand, 1 To 5 Percent Slopes-----	271	*
081LU	Lubbock Silty Clay Loam, 0 To 1 Percent Slopes-----	128	*
083UM	Uly-Coly Silt Loams, 3 To 6 Percent Slopes, Eroded-----	58	*
119HC	Harney Silty Clay Loam, 1 To 3 Percent Slopes, Eroded-----	58	*
119RX	Roxbury Silt Loam, Rarely Flooded-----	2,021	0.4
119UA	Uly Silt Loam, 0 To 1 Percent Slopes-----	183	*
119UT	Uly Silt Loam, 3 To 6 Percent Slopes, Eroded-----	56	*
An	Bridgeport Silt Loam, Channeled-----	3,637	0.7
ARR	Arkansas River-----	811	0.1
Bc	Bridgeport Silty Clay Loam, 0 To 1 Percent Slopes-----	1,436	0.3
Bd	Bridgeport Silty Clay Loam, 1 To 3 Percent Slopes-----	2,388	0.4
Bo	Optima Fine Sand, 0 To 10 Percent Slopes-----	4,801	0.9
Da	Dale Silt Loam, Rarely Flooded-----	2,204	0.4
Gr	Schamber Gravelly Sandy Loam, 5 To 25 Percent Slopes-----	1,364	0.2
GRP	Gravel Pit-----	38	*
Ha	Harney Silt Loam, 0 To 1 Percent Slopes-----	69,891	12.6
HB	Harney Silt Loam, 1 To 3 Percent Slopes-----	736	0.1
INL	Aquolls-----	179	*
La	Las Animas-Lesho Complex, Alkali, Occasionally Flooded-----	3,248	0.6
Lc	Las Animas Soils, Occasionally Flooded-----	4,170	0.7
Ld	Las Animas Sandy Loam, Occasionally Flooded-----	973	0.2
Le	Las Clay Loam, Occasionally Flooded-----	1,342	0.2
Lh	Lesho Clay Loam, Occasionally Flooded-----	1,917	0.3
Lk	Lesho-Sweetwater Complex, Occasionally Flooded-----	2,203	0.4
Ll	Lincoln Soils, Frequently Flooded-----	887	0.2
Lo	Lofton Silty Clay Loam, 0 To 1 Percent Slopes-----	1,615	0.3
Lu	Lubbock Loam, 0 To 1 Percent Slopes-----	1,717	0.3
Md	Penden Clay Loam, 3 To 6 Percent Slopes-----	10,760	1.9
Me	Penden Clay Loam, 6 To 15 Percent Slopes-----	11,431	2.1
Mf	Penden Clay Loam, 3 To 6 Percent Slopes, Severely Eroded-----	3,621	0.7
Mh	Penden Complex, 6 To 15 Percent Slopes, Severely Eroded-----	1,043	0.2
Mm	Manter Fine Sandy Loam, 0 To 1 Percent Slopes-----	11,057	2.0
Mn	Manter Fine Sandy Loam, 1 To 3 Percent Slopes-----	3,256	0.6
Mo	Manter Fine Sandy Loam, 1 To 6 Percent Slopes, Eroded-----	14,650	2.6
Mp	Campus-Canlon Complex, 6 To 15 Percent Slopes-----	595	0.1
Mu	Manter-Ulysses Complex, 1 To 6 Percent Slopes-----	8,262	1.5
Pa	Pratt Loamy Fine Sand, 1 To 6 Percent Slopes-----	43,077	7.7
Pg	Pratt Loamy Fine Sand, Gravel Substratum, 0 To 1 Percent Slopes-----	525	*
Pt	Pratt-Optima Loamy Fine Sands, 3 To 9 Percent Slopes-----	24,351	4.4
Ra	Ness Clay-----	10,653	1.9
Rm	Richfield Silt Loam, 0 To 1 Percent Slopes-----	72,792	13.1
Rn	Richfield Silt Loam, 1 To 3 Percent Slopes-----	19,849	3.6
Ro	Richfield Silty Clay Loam, 1 To 3 Percent Slopes, Eroded-----	1,023	0.2
Rs	Richfield-Spearville Complex, 0 To 1 Percent Slopes-----	24,937	4.5
Sa	Satanta Loam, 0 To 1 Percent Slopes-----	13,749	2.5
SAP	Sand Pit-----	33	*
Sb	Satanta Loam, 1 To 3 Percent Slopes-----	2,083	0.4
Sp	Spearville Silty Clay Loam, 0 To 1 Percent Slopes-----	120,920	21.7
Sr	Spearville Complex, 1 To 3 Percent Slopes, Eroded-----	2,466	0.4
Sw	Sweetwater Soils, Occasionally Flooded-----	827	0.1
Tf	Optima Fine Sand, 6 To 20 Percent Slopes-----	9,841	1.8
Ua	Ulysses Silt Loam, 0 To 1 Percent Slopes-----	8,811	1.6
Ub	Ulysses Silt Loam, 1 To 3 Percent Slopes-----	11,746	2.1
UBB	Uly Silt Loam, 1 To 3 Percent Slopes-----	918	0.2
Uc	Ulysses Silt Loam, 3 To 6 Percent Slopes-----	6,387	1.1
UCC	Uly Silt Loam, 3 To 6 Percent Slopes-----	594	0.1
Um	Ulysses-Colby Silt Loams, 3 To 6 Percent Slopes, Eroded-----	4,396	0.8
W	Water-----	63	*
	Total-----	556,045	100.0

\* Less than 0.1 percent.

Nontechnical Soil Descriptions  
Gray County, Kansas

Nontechnical soil descriptions describe soil properties or management considerations specific to a soil map unit or group of map units, shown in the Nontechnical Descriptions report. These descriptions are written in terminology that Non-technical users of soil survey information can understand. Nontechnical soil descriptions are a powerful tool for creating reports. These high quality, easy to read reports can be generated by conservation planners and other NRCS employees for distribution to land users. Soil map unit descriptions and National Soil Information System records are the basis for these descriptions.

---

055BP Bridgeport Clay Loam, Rarely Flooded

Bridgeport, rarely flooded, soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is low. The parent material consists of silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Loamy Terrace (pe16-20) range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

055LD Las-Las Animas Complex, Occasionally Flooded

Las, occasionally flooded, soil makes up 55 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 30 inches. The soil contains a maximum amount of 15 percent calcium carbonate. This soil contains a very slightly saline horizon, it has a horizon that is slightly sodic. This soil is in the Saline Subirrigated (pe16-20) range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 3w.

Las Animas, occasionally flooded, soil makes up 45 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy and/or sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a very slightly saline horizon. This soil is in the Saline Subirrigated (pe16-20) range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 4w.

055TF Valent Fine Sand, 6 To 20 Percent Slopes

Valent soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a strongly sloping to steep dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Choppy Sands (pe16-20) range site. This soil is in the irrigated land capability class 6e. It is in the nonirrigated land capability classification 7e.

055TV Valent-Vona Loamy Fine Sands, 3 To 8 Percent Slopes

Valent soil makes up 50 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sands (pe16-20) range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

Vona soil makes up 50 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on dune field on paleoterrace. The runoff class is very low. The parent material consists of eolian sands. This soil is somewhat excessively drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil contains a very slightly saline horizon. This soil is in the Sands (pe16-20) range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

055VO Vona Loamy Fine Sand, 1 To 5 Percent Slopes

Vona soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping dune on dune field on paleoterrace. The runoff class is very low. The parent material consists of eolian sands. This soil is somewhat excessively drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil contains a very slightly saline horizon. This soil is in the Sands (pe16-20) range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

Nontechnical Soil Descriptions--Continued  
Gray County, Kansas

081LU Lubbock Silty Clay Loam, 0 To 1 Percent Slopes

Lubbock soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level paleoterrace. The runoff class is low. The parent material consists of silty alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 3c.

083UM Uly-Coly Silt Loams, 3 To 6 Percent Slopes, Eroded

Uly soil makes up 70 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping plain on tableland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe20-26) range site. It is in the nonirrigated land capability classification 4e.

Coly soil makes up 30 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Limy Upland (pe20-26) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

119HC Harney Silty Clay Loam, 1 To 3 Percent Slopes, Eroded

Harney, eroded, soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a gently sloping plain on tableland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe20-26) range site. It is in the nonirrigated land capability classification 3e.

119RX Roxbury Silt Loam, Rarely Flooded

Roxbury soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of calcareous fine-silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Terrace (pe20-26) range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

119UA Uly Silt Loam, 0 To 1 Percent Slopes

Uly soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level plain on tableland. The runoff class is negligible. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Loamy Upland (pe20-26) range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

119UT Uly Silt Loam, 3 To 6 Percent Slopes, Eroded

Uly soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland (pe20-26) range site. It is in the nonirrigated land capability classification 4e.

An Bridgeport Silt Loam, Channeled

Bridgeport, channeled, soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is low. The parent material consists of silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Lowland (pe16-20) range site. It is in the nonirrigated land capability classification 5w.

Nontechnical Soil Descriptions--Continued  
Gray County, Kansas

**Bc Bridgeport Silty Clay Loam, 0 To 1 Percent Slopes**

Bridgeport soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Loamy Terrace (pe16-20) range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

**Bd Bridgeport Silty Clay Loam, 1 To 3 Percent Slopes**

Bridgeport soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a gently sloping flood plain on river valley. The runoff class is low. The parent material consists of silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Loamy Terrace (pe16-20) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 3e.

**Bo Optima Fine Sand, 0 To 10 Percent Slopes**

Optima soil makes up 70 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level to strongly sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sands (pe17-20) range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

**Da Dale Silt Loam, Rarely Flooded**

Dale soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Terrace (pe16-20) range site. It is in the nonirrigated land capability classification 2c.

**Ha Harney Silt Loam, 0 To 1 Percent Slopes**

Harney soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe20-26) range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

**HB Harney Silt Loam, 1 To 3 Percent Slopes**

Harney soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a gently sloping plain on tableland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe20-26) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

**INL Aquolls**

Aquolls soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level depression on terrace on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is very poorly drained. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is occasional ponded. The top of the seasonal high water table is at 0 inches. It is in the nonirrigated land capability classification 5w.

**La Las Animas-Lesho Complex, Alkali, Occasionally Flooded**

Las Animas, occasionally flooded, soil makes up 50 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy and/or sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil contains a slightly saline horizon. This soil is in the Saline Subirrigated (pe16-20) range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 4w.

Nontechnical Soil Descriptions--Continued  
Gray County, Kansas

---

Lesho, occasionally flooded, soil makes up 40 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level river valley on flood plain. The runoff class is low. The parent material consists of loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a moderately saline horizon. This soil is in the Saline Subirrigated (pe16-20) range site. This soil is in the irrigated land capability class 3s. It is in the nonirrigated land capability classification 4s.

Lc Las Animas Soils, Occasionally Flooded

Las Animas, occasionally flooded, soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of loamy and/or sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a very slightly saline horizon. This soil is in the Saline Subirrigated (pe16-20) range site. It is in the nonirrigated land capability classification 6s.

Ld Las Animas Sandy Loam, Occasionally Flooded

Las Animas soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy and/or sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a very slightly saline horizon. This soil is in the Saline Subirrigated (pe16-20) range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Le Las Clay Loam, Occasionally Flooded

Las, occasionally flooded, soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a low available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 30 inches. The soil contains a maximum amount of 15 percent calcium carbonate. This soil contains a very slightly saline horizon, it has a horizon that is slightly sodic. This soil is in the Saline Lowland (pe16-20) range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 4w.

Lh Lesho Clay Loam, Occasionally Flooded

Lesho, occasionally flooded, soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level river valley on flood plain. The runoff class is low. The parent material consists of loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a very slightly saline horizon. This soil is in the Saline Subirrigated (pe16-20) range site. It is in the nonirrigated land capability classification 3w.

Lk Lesho-Sweetwater Complex, Occasionally Flooded

Lesho soil makes up 50 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level to gently sloping river valley on flood plain. The runoff class is low. The parent material consists of loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a very slightly saline horizon. This soil is in the Saline Subirrigated (pe16-20) range site. It is in the nonirrigated land capability classification 6s.

Sweetwater soil makes up 25 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is poorly drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 21 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a very slightly saline horizon, it has a horizon that is slightly sodic. This soil is in the Saline Subirrigated (pe16-20) range site. It is in the nonirrigated land capability classification 5w.

Nontechnical Soil Descriptions--Continued  
Gray County, Kansas

Ll Lincoln Soils, Frequently Flooded

Lincoln, occasionally flooded, soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of loamy and/or sandy alluvium. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 66 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy Lowland (pe16-20) range site. It is in the nonirrigated land capability classification 7w.

Lu Lubbock Loam, 0 To 1 Percent Slopes

Lubbock soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level paleoterrace. The runoff class is medium. The parent material consists of silty alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

Md Penden Clay Loam, 3 To 6 Percent Slopes

Penden soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping plain on tableland. The runoff class is medium. The parent material consists of residuum. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland (pe20-26) range site. It is in the nonirrigated land capability classification 3e.

Me Penden Clay Loam, 6 To 15 Percent Slopes

Penden soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep plain on tableland. The runoff class is medium. The parent material consists of residuum. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland (pe16-20) range site. It is in the nonirrigated land capability classification 6e.

Mf Penden Clay Loam, 3 To 6 Percent Slopes, Severely Eroded

Penden, Severely eroded, soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping plain on tableland. The runoff class is low. The parent material consists of residuum. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland (pe16-20) range site. It is in the nonirrigated land capability classification 6e.

Mh Penden Complex, 6 To 15 Percent Slopes, Severely Eroded

Penden, Severely Eroded, soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep plain on tableland. The runoff class is medium. The parent material consists of residuum. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland (pe16-20) range site. It is in the nonirrigated land capability classification 6e.

Mm Manter Fine Sandy Loam, 0 To 1 Percent Slopes

Manter soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level sand sheet on paleoterrace on tableland. The runoff class is negligible. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy (pe16-20) range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2e.

Mn Manter Fine Sandy Loam, 1 To 3 Percent Slopes

Manter soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a gently sloping sand sheet on paleoterrace on tableland. The runoff class is very low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy (pe16-20) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.



Nontechnical Soil Descriptions--Continued  
Gray County, Kansas

Mo Manter Fine Sandy Loam, 1 To 6 Percent Slopes, Eroded

Manter soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping sand sheet on paleoterrace on tableland. The runoff class is very low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy (pel6-20) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Mp Campus-Canlon Complex, 6 To 15 Percent Slopes

Campus soil makes up 75 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping upland. The runoff class is medium. The parent material consists of old calcareous fine-loamy alluvium and/or calcareous fine-loamy residuum. The soil is 19 to 40 inches deep to bedrock (lithic). This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 30 percent calcium carbonate. This soil is in the Limy Upland (pel6-20) range site. It is in the nonirrigated land capability classification 4e.

Canlon soil makes up 25 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep upland. The runoff class is medium. The parent material consists of calcareous loamy residuum weathered from sandstone. The soil is 8 to 20 inches deep to bedrock (lithic). This soil is well drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Shallow Limy (pel6-20) range site. It is in the nonirrigated land capability classification 6e.

Mu Manter-Ulysses Complex, 1 To 6 Percent Slopes

Manter soil makes up 55 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping sand sheet on paleoterrace on tableland. The runoff class is very low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy (pel6-20) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Ulysses soil makes up 45 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Loamy Upland (pel6-20) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Pa Pratt Loamy Fine Sand, 1 To 6 Percent Slopes

Pratt soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping dune on paleoterrace on river valley. The runoff class is negligible. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sands (pel6-20) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Pg Pratt Loamy Fine Sand, Gravel Substratum, 0 To 1 Percent Slopes

Pratt soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level dune on paleoterrace on river valley. The runoff class is negligible. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sands (pel6-20) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

Pt Pratt-Optima Loamy Fine Sands, 3 To 9 Percent Slopes

Pratt soil makes up 60 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sands (pel6-20) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

Nontechnical Soil Descriptions--Continued  
Gray County, Kansas

Optima soil makes up 40 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sands (pe17-20) range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

**Ra Ness Clay**

Ness soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level playa on tableland. The runoff class is negligible. The parent material consists of clayey alluvium and/or eolian deposits. This soil is poorly drained. The slowest permeability is very slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is frequent ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Lakebed (pe20-26) range site. It is in the nonirrigated land capability classification 6w.

**Rm Richfield Silt Loam, 0 To 1 Percent Slopes**

Richfield soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

**Rn Richfield Silt Loam, 1 To 3 Percent Slopes**

Richfield soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a gently sloping plain on tableland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

**Ro Richfield Silty Clay Loam, 1 To 3 Percent Slopes, Eroded**

Richfield soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a gently sloping plain on tableland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

**Rs Richfield-Spearville Complex, 0 To 1 Percent Slopes**

Richfield soil makes up 60 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

Spearville soil makes up 40 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level plain on tableland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Clay Upland (pe16-20) range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2s.

**Sa Satanta Loam, 0 To 1 Percent Slopes**

Satanta soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level sand sheet on paleoterrace on tableland. The runoff class is low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

Nontechnical Soil Descriptions--Continued  
Gray County, Kansas

---

Sb Satanta Loam, 1 To 3 Percent Slopes

Satanta soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a gently sloping sand sheet on paleoterrace on tableland. The runoff class is low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Sp Spearville Silty Clay Loam, 0 To 1 Percent Slopes

Spearville soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a nearly level plain on tableland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Clay Upland (pe16-20) range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2s.

Sr Spearville Complex, 1 To 3 Percent Slopes, Eroded

Spearville soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a gently sloping plain on tableland. The runoff class is high. The parent material consists of loess. This soil is well drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Clay Upland (pe16-20) range site. It is in the nonirrigated land capability classification 3e.

Sw Sweetwater Soils, Occasionally Flooded

Sweetwater soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of loamy alluvium. This soil is poorly drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 21 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a very slightly saline horizon, it has a horizon that is slightly sodic. This soil is in the Saline Subirrigated (pe16-20) range site. It is in the nonirrigated land capability classification 5w.

Tf Optima Fine Sand, 6 To 20 Percent Slopes

Optima soil makes up 90 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sands (pe17-20) range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

Ua Ulysses Silt Loam, 0 To 1 Percent Slopes

Ulysses soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a nearly level plain on tableland. The runoff class is negligible. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 2c.

Ub Ulysses Silt Loam, 1 To 3 Percent Slopes

Ulysses soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a gently sloping plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

UBB Uly Silt Loam, 1 To 3 Percent Slopes

Uly soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a gently sloping plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe20-26) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

NONTECHNICAL SOIL DESCRIPTIONS--Continued  
Gray County, Kansas

## Uc Ulysses Silt Loam, 3 To 6 Percent Slopes

Ulysses soil makes up 100 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a moderately sloping plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

## UCC Uly Silt Loam, 3 To 6 Percent Slopes

Uly soil makes up 100 percent of the map unit. This map unit is in the Rolling Plains and Breaks Major Land Resource Area. This soil occurs on a moderately sloping plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Upland (pe20-26) range site. It is in the nonirrigated land capability classification 3e.

## Um Ulysses-Colby Silt Loams, 3 To 6 Percent Slopes, Eroded

Ulysses soil makes up 55 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a moderately sloping plain on tableland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Loamy Upland (pe16-20) range site. It is in the nonirrigated land capability classification 4e.

Colby soil makes up 45 percent of the map unit. This map unit is in the Central High Tableland Major Land Resource Area. This soil occurs on a moderately sloping hillslope on tableland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Limy Upland (pe16-20) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

**055BP—Bridgeport clay loam,  
rarely flooded****Map Unit Composition**

Bridgeport: 100 percent

**Component Descriptions****Bridgeport**

*MLRA:* 72 - Central High Tableland

*Landform:* Flood plain on river valley

*Parent material:* Silty alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 11.6 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* Rare

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Loamy Terrace (pe16-20)

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 2c

*Typical Profile:*

H1—0 to 11 inches; clay loam

H2—11 to 60 inches; loam

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Moderate (About 6.2 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 24 to 36 inches

*Runoff class:* Low

*Ecological site:* Saline Subirrigated (pe16-20)

*Land capability (irrigated):* 2w

*Land capability (nonirrigated):* 3w

*Typical Profile:*

H1—0 to 8 inches; loam

H2—8 to 31 inches; clay loam

H3—31 to 60 inches; coarse sand

**Las Animas**

*MLRA:* 72 - Central High Tableland

*Landform:* Flood plain on river valley

*Parent material:* Loamy and/or sandy alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Somewhat poorly drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 6.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 18 to 36 inches

*Runoff class:* Negligible

*Ecological site:* Saline Subirrigated (pe16-20)

*Land capability (irrigated):* 3w

*Land capability (nonirrigated):* 4w

*Typical Profile:*

H1—0 to 14 inches; fine sandy loam

H2—14 to 30 inches; stratified very fine sandy loam to loamy fine sand

H3—30 to 60 inches; fine sand

**055LD—Las-Las Animas complex,  
occasionally flooded****Map Unit Composition**

Las: 55 percent

Las Animas: 45 percent

**Component Descriptions****Las**

*MLRA:* 72 - Central High Tableland

*Landform:* Flood plain on river valley

*Parent material:* Loamy alluvium over sandy and gravelly alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Somewhat poorly drained

**Minor Components  
Sweetwater****055TF—Valent fine sand, 6 to 20  
percent slopes****Map Unit Composition**

Valent: 100 percent

### Component Descriptions

#### Valent

*MLRA:* 72 - Central High Tableland  
*Landform:* Dune on paleoterrace on river valley  
*Parent material:* Sandy eolian deposits  
*Slope:* 8 to 30 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 4.0 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Choppy Sands (pe16-20)  
*Land capability (irrigated):* 6e  
*Land capability (nonirrigated):* 7e

#### Typical Profile:

H1—0 to 4 inches; fine sand  
 H2—4 to 50 inches; fine sand

### 055TV—Valent-Vona loamy fine sands, 3 to 8 percent slopes

### Map Unit Composition

Valent: 50 percent  
 Vona: 50 percent

### Component Descriptions

#### Valent

*MLRA:* 72 - Central High Tableland  
*Landform:* Dune on paleoterrace on river valley  
*Parent material:* Sandy eolian deposits  
*Slope:* 5 to 8 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Rapid (About 5.95 in/hr)  
*Available water capacity:* Low (About 4.8 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Sands (pe16-20)  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 6e

#### Typical Profile:

H1—0 to 4 inches; loamy fine sand  
 H2—4 to 60 inches; fine sand

#### Vona

*MLRA:* 72 - Central High Tableland  
*Landform:* Dune on dune field on paleoterrace  
*Parent material:* Eolian sands  
*Slope:* 3 to 8 percent  
*Drainage class:* Somewhat excessively drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Moderate (About 7.1 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Sands (pe16-20)  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 6e

#### Typical Profile:

H1—0 to 8 inches; loamy fine sand  
 H2—8 to 22 inches; fine sandy loam  
 H3—22 to 60 inches; fine sandy loam

### Minor Components Unnamed Hydric Soils

### 055VO—Vona loamy fine sand, 1 to 5 percent slopes

### Map Unit Composition

Vona: 100 percent

### Component Descriptions

#### Vona

*MLRA:* 72 - Central High Tableland  
*Landform:* Dune on dune field on paleoterrace  
*Parent material:* Eolian sands  
*Slope:* 1 to 5 percent  
*Drainage class:* Somewhat excessively drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Moderate (About 7.1 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Sands (pe16-20)  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 8 inches; loamy fine sand  
 H2—8 to 22 inches; fine sandy loam  
 H3—22 to 60 inches; sandy loam

**Minor Components**  
**Unnamed Hydric Soils**

**081LU—Lubbock silty clay loam, 0 to 1 percent slopes**

**Map Unit Composition**

Lubbock: 100 percent

**Component Descriptions**

**Lubbock**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Paleoterrace  
*Parent material:* Silty alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 10.0 inches)  
*Shrink-swell potential:* High (About 7.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Upland (pe16-20)  
*Land capability (irrigated):* 1  
*Land capability (nonirrigated):* 3c

*Typical Profile:*

H1—0 to 9 inches; silty clay loam  
 H2—9 to 42 inches; silty clay loam  
 H3—42 to 72 inches; silt loam

**083UM—Uly-Coly silt loams, 3 to 6 percent slopes, eroded**

**Map Unit Composition**

Uly: 70 percent  
 Coly: 30 percent

**Component Descriptions**

**Uly**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 3 to 6 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* High (About 12.0 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Medium  
*Ecological site:* Loamy Upland (pe20-26)  
*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 6 inches; silt loam  
 H2—6 to 19 inches; silty clay loam  
 H3—19 to 60 inches; silt loam

**Coly**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 3 to 6 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* High (About 12.0 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Limy Upland (pe20-26)  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 6 inches; silt loam  
 H2—6 to 60 inches; silt loam

**119HC—Harney silty clay loam, 1 to 3 percent slopes, eroded**

**Map Unit Composition**

Harney: 100 percent

## Component Descriptions

### Harney

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Plain on tableland

*Parent material:* Loess

*Slope:* 1 to 3 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.4 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Loamy Upland (pe20-26)

*Land capability (nonirrigated):* 3e

#### Typical Profile:

H1—0 to 6 inches; silty clay loam

H2—6 to 20 inches; silty clay loam

H3—20 to 60 inches; silt loam

## 119RX—Roxbury silt loam, rarely flooded

### Map Unit Composition

Roxbury: 100 percent

## Component Descriptions

### Roxbury

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Flood plain on river valley

*Parent material:* Calcareous fine-silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.4 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* Rare

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Loamy Terrace (pe20-26)

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 2c

#### Typical Profile:

H1—0 to 20 inches; silt loam

H2—20 to 60 inches; silty clay loam

## 119UA—Uly silt loam, 0 to 1 percent slopes

### Map Unit Composition

Uly: 100 percent

## Component Descriptions

### Uly

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Plain on tableland

*Parent material:* Loess

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 12.0 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Loamy Upland (pe20-26)

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 2c

#### Typical Profile:

A—0 to 6 inches; silt loam

Bw—6 to 12 inches; silt loam

Bk—12 to 18 inches; silt loam

Bck—18 to 32 inches; silt loam

C—32 to 62 inches; silt loam

### Minor Components

#### Ness

## 119UT—Uly silt loam, 3 to 6 percent slopes, eroded

### Map Unit Composition

Uly: 100 percent



### Component Descriptions

#### Uly

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Plain on tableland

*Parent material:* Loess

*Slope:* 3 to 6 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.2 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Limy Upland (pe20-26)

*Land capability (nonirrigated):* 4e

#### Typical Profile:

H1—0 to 6 inches; silt loam

H2—6 to 14 inches; silty clay loam

H3—14 to 60 inches; silt loam

### An—Bridgeport silt loam, channeled

#### Map Unit Composition

Bridgeport: 100 percent

### Component Descriptions

#### Bridgeport

*MLRA:* 72 - Central High Tableland

*Landform:* Flood plain on river valley

*Parent material:* Silty alluvium

*Slope:* 0 to 2 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 11.9 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Frequent

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Loamy Lowland (pe16-20)

*Land capability (nonirrigated):* 5w

#### Typical Profile:

H1—0 to 12 inches; loam

H2—12 to 60 inches; loam

### ARR—Arkansas River

### Bc—Bridgeport silty clay loam, 0 to 1 percent slopes

#### Map Unit Composition

Bridgeport: 100 percent

### Component Descriptions

#### Bridgeport

*MLRA:* 72 - Central High Tableland

*Landform:* Flood plain on river valley

*Parent material:* Silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.0 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* Rare

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Loamy Terrace (pe16-20)

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 2c

#### Typical Profile:

H1—0 to 10 inches; silty clay loam

H2—10 to 65 inches; silty clay loam

### Bd—Bridgeport silty clay loam, 1 to 3 percent slopes

#### Map Unit Composition

Bridgeport: 100 percent

### Component Descriptions

#### Bridgeport

*MLRA:* 72 - Central High Tableland

*Landform:* Flood plain on river valley

*Parent material:* Silty alluvium

*Slope:* 1 to 3 percent

*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* Very high (About 12.0 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* Rare  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Terrace (pe16-20)  
*Land capability (irrigated):* 2e  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 10 inches; silty clay loam  
 H2—10 to 65 inches; silty clay loam

## **Bo—Optima fine sand, 0 to 10 percent slopes**

### **Map Unit Composition**

Optima: 70 percent  
 Minor components: 30 percent

### **Component Descriptions**

#### **Optima**

*MLRA:* 72 - Central High Tableland  
*Landform:* Dune on paleoterrace on river valley  
*Parent material:* Sandy eolian deposits  
*Slope:* 0 to 10 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Rapid (About 6.00 in/hr)  
*Available water capacity:* Low (About 4.5 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Sands (pe17-20)  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 6e

*Typical Profile:*

A—0 to 8 inches; loamy fine sand  
 AC—8 to 17 inches; fine sand  
 C—17 to 80 inches; fine sand

#### **Minor Components**

##### **Eva**

*Composition:* About 30 percent  
*Slope:* 2 to 6 percent

*Drainage class:* Somewhat excessively drained  
*Ecological site:* Sands (pe17-20)

## **Da—Dale silt loam, rarely flooded**

### **Map Unit Composition**

Dale: 100 percent

### **Component Descriptions**

#### **Dale**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Flood plain on river valley  
*Parent material:* Silty alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* Very high (About 12.6 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* Rare  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Negligible  
*Ecological site:* Loamy Terrace (pe16-20)  
*Land capability (nonirrigated):* 2c

*Typical Profile:*

H1—0 to 25 inches; silt loam  
 H2—25 to 60 inches; silt loam

## **Gr—Schamber gravelly sandy loam, 5 to 25 percent slopes**

### **Map Unit Composition**

Schamber: 100 percent

### **Component Descriptions**

#### **Schamber**

*MLRA:* 72 - Central High Tableland  
*Landform:* Fan remnant on paleoterrace on river valley  
*Parent material:* Sandy and/or gravelly alluvium  
*Slope:* 5 to 25 percent  
*Drainage class:* Somewhat excessively drained

*Slowest permeability:* Rapid (About 6.00 in/hr)  
*Available water capacity:* Very low (About 3.0 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Gravelly Hills (pe16-20)  
*Land capability (nonirrigated):* 7s

## GRP—Gravel Pit

### Ha—Harney silt loam, 0 to 1 percent slopes

#### Map Unit Composition

Harney: 100 percent

#### Component Descriptions

##### Harney

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 0 to 1 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 11.8 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Upland (pe20-26)  
*Land capability (irrigated):* 1  
*Land capability (nonirrigated):* 2c

##### Typical Profile:

H1—0 to 9 inches; silt loam  
H2—9 to 25 inches; silty clay loam  
H3—25 to 65 inches; silty clay loam

#### Minor Components

##### Ness

### HB—Harney silt loam, 1 to 3 percent slopes

#### Map Unit Composition

Harney: 100 percent

#### Component Descriptions

##### Harney

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 1 to 3 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* High (About 11.4 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Medium  
*Ecological site:* Loamy Upland (pe20-26)  
*Land capability (irrigated):* 2e  
*Land capability (nonirrigated):* 2e

##### Typical Profile:

H1—0 to 6 inches; silt loam  
H2—6 to 20 inches; silty clay loam  
H3—20 to 60 inches; silt loam

### INL—Aquolls

#### Map Unit Composition

Aquolls: 100 percent

#### Component Descriptions

##### Aquolls

*MLRA:* 72 - Central High Tableland  
*Landform:* Depression on terrace on river valley  
*Parent material:* Alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Very poorly drained  
*Flooding hazard:* None  
*Ponding hazard:* Occasional  
*Depth to seasonal water saturation:* About 0 to 0 inches

*Runoff class:* Negligible  
*Land capability (nonirrigated):* 5w

*Typical Profile:*  
 H1—0 to 72 inches; variable

*General Considerations:* This map unit was formerly labeled as an Intermittent Water spot symbol. These depressional areas contain soils that are occasionally ponded for long duration.

## **La—Las Animas-Lesho complex, Alkali, occasionally flooded**

### **Map Unit Composition**

Las Animas: 50 percent  
 Lesho: 40 percent  
 Minor components: 10 percent

### **Component Descriptions**

#### **Las Animas**

*MLRA:* 72 - Central High Tableland  
*Landform:* Flood plain on river valley  
*Parent material:* Loamy and/or sandy alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Moderate (About 8.3 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 24 to 48 inches  
*Runoff class:* Negligible  
*Ecological site:* Saline Subirrigated (pe16-20)  
*Land capability (irrigated):* 3w  
*Land capability (nonirrigated):* 4w

*Typical Profile:*  
 H1—0 to 11 inches; sandy loam  
 H2—11 to 60 inches; sand

#### **Lesho**

*MLRA:* 72 - Central High Tableland  
*Landform:* River valley on flood plain  
*Parent material:* Loamy alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Moderate (About 6.2 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 24 to 48 inches  
*Runoff class:* Low  
*Ecological site:* Saline Subirrigated (pe16-20)  
*Land capability (irrigated):* 3s  
*Land capability (nonirrigated):* 4s

*Typical Profile:*  
 H1—0 to 14 inches; clay loam  
 H2—14 to 25 inches; clay loam  
 H3—25 to 60 inches; coarse sand

### **Minor Components**

#### **Slickspots**

*Composition:* About 10 percent  
*Slope:* 0 to 1 percent  
*Drainage class:* Moderately well drained  
*Ecological site:* Clay Pan (pe21-28)

### **Sweetwater**

## **Lc—Las Animas Soils, occasionally flooded**

### **Map Unit Composition**

Las Animas: 100 percent

### **Component Descriptions**

#### **Las Animas**

*MLRA:* 72 - Central High Tableland  
*Landform:* Flood plain on river valley  
*Parent material:* Loamy and/or sandy alluvium  
*Slope:* 0 to 2 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Low (About 5.6 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 18 to 36 inches  
*Runoff class:* Negligible  
*Ecological site:* Saline Subirrigated (pe16-20)  
*Land capability (nonirrigated):* 6s

*Typical Profile:*  
 H1—0 to 11 inches; loamy sand

H2—11 to 30 inches; stratified very fine  
sandy loam to loamy fine sand  
H3—30 to 60 inches; fine sand

**Minor Components**  
**Sweetwater**

**Ld—Las Animas sandy loam,  
occasionally flooded**

**Map Unit Composition**

Las Animas: 100 percent

**Component Descriptions**

**Las Animas**

*MLRA:* 72 - Central High Tableland  
*Landform:* Flood plain on river valley  
*Parent material:* Loamy and/or sandy alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Moderately rapid (About 2.00 in/hr)  
*Available water capacity:* Low (About 5.9 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 18 to 36 inches  
*Runoff class:* Negligible  
*Ecological site:* Saline Subirrigated (pe16-20)  
*Land capability (irrigated):* 3w  
*Land capability (nonirrigated):* 3w

*Typical Profile:*

H1—0 to 11 inches; sandy loam  
H2—11 to 30 inches; stratified very fine  
sandy loam to loamy fine sand  
H3—30 to 60 inches; fine sand

**Minor Components**  
**Sweetwater**

**Le—Las clay loam, occasionally  
flooded**

**Map Unit Composition**

Las: 100 percent

**Component Descriptions**

**Las**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Flood plain on river valley  
*Parent material:* Loamy alluvium over sandy and gravelly alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* Low (About 6.0 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 24 to 36 inches  
*Runoff class:* Low  
*Ecological site:* Saline Lowland (pe16-20)  
*Land capability (irrigated):* 2w  
*Land capability (nonirrigated):* 4w

*Typical Profile:*

H1—0 to 8 inches; clay loam  
H2—8 to 31 inches; clay loam  
H3—31 to 60 inches; coarse sand

**Minor Components**  
**Sweetwater**

**Lh—Lesho clay loam, occasionally  
flooded**

**Map Unit Composition**

Lesho: 100 percent

**Component Descriptions**

**Lesho**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* River valley on flood plain  
*Parent material:* Loamy alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* Moderate (About 6.1 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 24 to 48 inches

*Runoff class:* Low  
*Ecological site:* Saline Subirrigated (pe16-20)  
*Land capability (nonirrigated):* 3w

*Typical Profile:*

H1—0 to 14 inches; clay loam  
 H2—14 to 25 inches; clay loam  
 H3—25 to 60 inches; coarse sand

**Minor Components**  
**Sweetwater**

**Lk—Lesho-Sweetwater complex, occasionally flooded**

**Map Unit Composition**

Lesho: 50 percent  
 Sweetwater: 25 percent  
 Minor components: 25 percent

**Component Descriptions**

**Lesho**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* River valley on flood plain  
*Parent material:* Loamy alluvium  
*Slope:* 0 to 2 percent  
*Drainage class:* Somewhat poorly drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)  
*Available water capacity:* Moderate (About 6.5 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* Occasional  
*Depth to seasonal water saturation:* About 24 to 48 inches  
*Runoff class:* Low  
*Ecological site:* Saline Subirrigated (pe16-20)  
*Land capability (nonirrigated):* 6s

*Typical Profile:*

H1—0 to 14 inches; loam  
 H2—14 to 25 inches; clay loam  
 H3—25 to 60 inches; coarse sand

**Sweetwater**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Flood plain on river valley  
*Parent material:* Loamy alluvium  
*Slope:* 0 to 1 percent  
*Drainage class:* Poorly drained  
*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Moderate (About 6.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 6 to 36 inches

*Runoff class:* Low

*Ecological site:* Saline Subirrigated (pe16-20)

*Land capability (nonirrigated):* 5w

*Typical Profile:*

H1—0 to 17 inches; clay loam  
 H2—17 to 60 inches; coarse sand

**Minor Components**

**Las Animas**

*Phase:* Occasionally Flooded  
*Composition:* About 20 percent  
*Slope:* 0 to 1 percent  
*Drainage class:* Somewhat poorly drained  
*Ecological site:* Saline Subirrigated (pe16-20)

**Optima**

*Composition:* About 5 percent  
*Slope:* 5 to 8 percent  
*Drainage class:* Excessively drained  
*Ecological site:* Sands (pe17-20)

**LI—Lincoln Soils, frequently flooded**

**Map Unit Composition**

Lincoln: 100 percent

**Component Descriptions**

**Lincoln**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Flood plain on river valley  
*Parent material:* Loamy and/or sandy alluvium  
*Slope:* 0 to 3 percent  
*Drainage class:* Somewhat excessively drained  
*Slowest permeability:* Rapid (About 6.00 in/hr)  
*Available water capacity:* Moderate (About 7.5 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* Frequent  
*Depth to seasonal water saturation:* About 60 to 72 inches  
*Runoff class:* Negligible  
*Ecological site:* Sandy Lowland (pe16-20)  
*Land capability (nonirrigated):* 7w

*Typical Profile:*

- H1—0 to 8 inches; fine sandy loam
- H2—8 to 60 inches; stratified fine sand to clay loam

**Minor Components**  
**Sweetwater**

**Lo—Lofton silty clay loam, 0 to 1 percent slopes**

**Map Unit Composition**

Lofton: 100 percent

**Component Descriptions**

**Lofton**

*MLRA:* 72 - Central High Tableland

*Landform:* Bench on playa on tableland

*Parent material:* Clayey alluvium and/or eolian deposits

*Slope:* 0 to 1 percent

*Slowest permeability:* Very slow (About 0.00 in/hr)

*Available water capacity:* High (About 10.4 inches)

*Shrink-swell potential:* High (About 7.5 LEP)

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Typical Profile:*

- H1—0 to 6 inches; silty clay loam
- H2—6 to 36 inches; silty clay
- H3—36 to 60 inches; silty clay loam

**Lu—Lubbock loam, 0 to 1 percent slopes**

**Map Unit Composition**

Lubbock: 100 percent

**Component Descriptions**

**Lubbock**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Paleoterrace

*Parent material:* Silty alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 10.9 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Loamy Upland (pe16-20)

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 2c

*Typical Profile:*

- H1—0 to 15 inches; loam
- H2—15 to 36 inches; silty clay loam
- H3—36 to 60 inches; clay loam

**Md—Penden clay loam, 3 to 6 percent slopes**

**Map Unit Composition**

Penden: 100 percent

**Component Descriptions**

**Penden**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Plain on tableland

*Parent material:* Residuum

*Slope:* 3 to 6 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 10.4 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Limy Upland (pe20-26)

*Land capability (nonirrigated):* 3e

*Typical Profile:*

- H1—0 to 14 inches; clay loam
- H2—14 to 32 inches; clay loam
- H3—32 to 60 inches; clay loam

## **Me—Penden clay loam, 6 to 15 percent slopes**

### **Map Unit Composition**

Penden: 100 percent

### **Component Descriptions**

#### **Penden**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Plain on tableland

*Parent material:* Residuum

*Slope:* 6 to 15 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 10.4 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Limy Upland (pe16-20)

*Land capability (nonirrigated):* 6e

#### *Typical Profile:*

H1—0 to 15 inches; clay loam

H2—15 to 34 inches; clay loam

H3—34 to 60 inches; clay loam

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 10.3 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Limy Upland (pe16-20)

*Land capability (nonirrigated):* 6e

#### *Typical Profile:*

H1—0 to 4 inches; clay loam

H2—4 to 22 inches; clay loam

H3—22 to 30 inches; clay loam

H4—30 to 80 inches; clay loam

## **Mh—Penden complex, 6 to 15 percent slopes, Severely eroded**

### **Map Unit Composition**

Penden: 100 percent

### **Component Descriptions**

#### **Penden**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Plain on tableland

*Parent material:* Residuum

*Slope:* 6 to 15 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* High (About 10.3 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Limy Upland (pe16-20)

*Land capability (nonirrigated):* 6e

#### *Typical Profile:*

H1—0 to 4 inches; clay loam

H2—4 to 22 inches; clay loam

H3—22 to 30 inches; clay loam

H4—30 to 80 inches; clay loam

## **Mf—Penden clay loam, 3 to 6 percent slopes, Severely eroded**

### **Map Unit Composition**

Penden: 100 percent

### **Component Descriptions**

#### **Penden**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Plain on tableland

*Parent material:* Residuum

*Slope:* 3 to 6 percent

*Drainage class:* Well drained



## **Mm—Manter fine sandy loam, 0 to 1 percent slopes**

### **Map Unit Composition**

Manter: 100 percent

### **Component Descriptions**

#### **Manter**

*MLRA:* 72 - Central High Tableland

*Landform:* Sand sheet on paleoterrace on tableland

*Parent material:* Loamy eolian deposits

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 8.1 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Negligible

*Ecological site:* Sandy (pe16-20)

*Land capability (irrigated):* 2s

*Land capability (nonirrigated):* 2e

#### *Typical Profile:*

H1—0 to 18 inches; fine sandy loam

H2—18 to 30 inches; fine sandy loam

H3—30 to 62 inches; loam

#### **Minor Components**

#### **Unnamed Hydric Soils**

## **Mn—Manter fine sandy loam, 1 to 3 percent slopes**

### **Map Unit Composition**

Manter: 100 percent

### **Component Descriptions**

#### **Manter**

*MLRA:* 72 - Central High Tableland

*Landform:* Sand sheet on paleoterrace on tableland

*Parent material:* Loamy eolian deposits

*Slope:* 1 to 3 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 8.1 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sandy (pe16-20)

*Land capability (irrigated):* 2e

*Land capability (nonirrigated):* 2e

#### *Typical Profile:*

H1—0 to 18 inches; fine sandy loam

H2—18 to 30 inches; fine sandy loam

H3—30 to 62 inches; sandy loam

## **Mo—Manter fine sandy loam, 1 to 6 percent slopes, eroded**

### **Map Unit Composition**

Manter: 100 percent

### **Component Descriptions**

#### **Manter**

*MLRA:* 72 - Central High Tableland

*Landform:* Sand sheet on paleoterrace on tableland

*Parent material:* Loamy eolian deposits

*Slope:* 1 to 6 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 8.1 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sandy (pe16-20)

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

#### *Typical Profile:*

H1—0 to 18 inches; fine sandy loam

H2—18 to 30 inches; fine sandy loam

H3—30 to 62 inches; sandy loam

**Minor Components**  
**Unnamed Hydric Soils**

**Mp—Campus-Canlon complex, 6 to 15 percent slopes**

**Map Unit Composition**

Campus: 75 percent  
 Canlon: 25 percent

**Component Descriptions**

**Campus**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Upland

*Parent material:* Old calcareous fine-loamy alluvium and/or calcareous fine-loamy residuum

*Slope:* 6 to 8 percent

*Depth to restrictive feature:* 19 to 40 inches to bedrock (lithic)

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Low (About 3.6 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Limy Upland (pe16-20)

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 9 inches; loam

H2—9 to 19 inches; loam

R—19 to 19 inches; unweathered bedrock

**Canlon**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Upland

*Parent material:* Calcareous loamy residuum weathered from sandstone

*Slope:* 6 to 15 percent

*Depth to restrictive feature:* 8 to 20 inches to bedrock (lithic)

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very low (About 1.8 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Shallow Limy (pe16-20)

*Land capability (nonirrigated):* 6e

*Typical Profile:*

H1—0 to 8 inches; silt loam

R—8 to 8 inches; unweathered bedrock

**Mu—Manter-Ulysses complex, 1 to 6 percent slopes**

**Map Unit Composition**

Manter: 55 percent  
 Ulysses: 45 percent

**Component Descriptions**

**Manter**

*MLRA:* 72 - Central High Tableland

*Landform:* Sand sheet on paleoterrace on tableland

*Parent material:* Loamy eolian deposits

*Slope:* 1 to 6 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately rapid (About 2.00 in/hr)

*Available water capacity:* Moderate (About 8.1 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sandy (pe16-20)

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 18 inches; fine sandy loam

H2—18 to 30 inches; fine sandy loam

H3—30 to 62 inches; sandy loam

**Ulysses**

*MLRA:* 72 - Central High Tableland

*Landform:* Plain on tableland

*Parent material:* Loess

*Slope:* 1 to 6 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderate (About 0.60 in/hr)

*Available water capacity:* Very high (About 12.2 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Upland (pe16-20)  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 5 inches; silt loam  
 H2—5 to 14 inches; silty clay loam  
 H3—14 to 60 inches; silt loam

## **Pa—Pratt loamy fine sand, 1 to 6 percent slopes**

### **Map Unit Composition**

Pratt: 100 percent

### **Component Descriptions**

#### **Pratt**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Dune on paleoterrace on river valley  
*Parent material:* Sandy eolian deposits  
*Slope:* 1 to 6 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Rapid (About 6.00 in/hr)  
*Available water capacity:* Low (About 5.8 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Negligible  
*Ecological site:* Sands (pe16-20)  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

H1—0 to 11 inches; loamy fine sand  
 H2—11 to 26 inches; loamy fine sand  
 H3—26 to 60 inches; loamy fine sand

#### **Minor Components** **Unnamed Hydric Soils**

## **Pg—Pratt loamy fine sand, Gravel Substratum, 0 to 1 percent slopes**

### **Map Unit Composition**

Pratt: 100 percent

### **Component Descriptions**

#### **Pratt**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Dune on paleoterrace on river valley  
*Parent material:* Sandy eolian deposits  
*Slope:* 0 to 1 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Rapid (About 6.00 in/hr)  
*Available water capacity:* Moderate (About 6.5 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Negligible  
*Ecological site:* Sands (pe16-20)  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 11 inches; loamy fine sand  
 H2—11 to 40 inches; loamy fine sand  
 H3—40 to 60 inches; loamy fine sand

## **Pt—Pratt-Optima loamy fine sands, 3 to 9 percent slopes**

### **Map Unit Composition**

Pratt: 60 percent  
 Optima: 40 percent

### **Component Descriptions**

#### **Pratt**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Dune on paleoterrace on river valley  
*Parent material:* Sandy eolian deposits  
*Slope:* 3 to 9 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Rapid (About 6.00 in/hr)  
*Available water capacity:* Low (About 5.8 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sands (pe16-20)

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 4e

*Typical Profile:*

H1—0 to 11 inches; loamy fine sand  
H2—11 to 26 inches; loamy fine sand  
H3—26 to 60 inches; loamy fine sand

**Optima**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Dune on paleoterrace on river valley

*Parent material:* Sandy eolian deposits

*Slope:* 5 to 8 percent

*Drainage class:* Excessively drained

*Slowest permeability:* Rapid (About 6.00 in/hr)

*Available water capacity:* Low (About 4.5 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Very low

*Ecological site:* Sands (pe17-20)

*Land capability (irrigated):* 4e

*Land capability (nonirrigated):* 6e

*Typical Profile:*

A—0 to 8 inches; loamy fine sand  
AC—8 to 17 inches; fine sand  
C—17 to 80 inches; fine sand

**Ra—Ness clay**

**Map Unit Composition**

Ness: 100 percent

**Component Descriptions**

**Ness**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Playa on tableland

*Parent material:* Clayey alluvium and/or eolian deposits

*Slope:* 0 to 1 percent

*Drainage class:* Poorly drained

*Slowest permeability:* Very slow (About 0.00 in/hr)

*Available water capacity:* High (About 9.0 inches)

*Shrink-swell potential:* High (About 7.5 LEP)

*Flooding hazard:* None

*Ponding hazard:* Frequent

*Depth to seasonal water saturation:* About 0 to 0 inches

*Runoff class:* Negligible

*Ecological site:* Lakebed (pe20-26)

*Land capability (nonirrigated):* 6w

*Typical Profile:*

H1—0 to 31 inches; clay  
H2—31 to 60 inches; silt loam

**Rm—Richfield silt loam, 0 to 1 percent slopes**

**Map Unit Composition**

Richfield: 100 percent

**Component Descriptions**

**Richfield**

*MLRA:* 72 - Central High Tableland

*Landform:* Plain on tableland

*Parent material:* Loess

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 11.5 inches)

*Shrink-swell potential:* Moderate (About 4.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Loamy Upland (pe16-20)

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 2c

*Typical Profile:*

H1—0 to 5 inches; silt loam  
H2—5 to 17 inches; silty clay loam  
H3—17 to 60 inches; silty clay loam

**Minor Components**

**Ness**

## **Rn—Richfield silt loam, 1 to 3 percent slopes**

### **Map Unit Composition**

Richfield: 100 percent

### **Component Descriptions**

#### **Richfield**

*MLRA:* 72 - Central High Tableland

*Landform:* Plain on tableland

*Parent material:* Loess

*Slope:* 1 to 3 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 10.7 inches)

*Shrink-swell potential:* High (About 7.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Loamy Upland (pe16-20)

*Land capability (irrigated):* 2e

*Land capability (nonirrigated):* 2e

#### *Typical Profile:*

H1—0 to 6 inches; silt loam

H2—6 to 19 inches; silty clay loam

H3—19 to 60 inches; silt loam

#### **Minor Components**

#### **Ness**

## **Ro—Richfield silty clay loam, 1 to 3 percent slopes, eroded**

### **Map Unit Composition**

Richfield: 100 percent

### **Component Descriptions**

#### **Richfield**

*MLRA:* 72 - Central High Tableland

*Landform:* Plain on tableland

*Parent material:* Loess

*Slope:* 1 to 3 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 10.6 inches)

*Shrink-swell potential:* High (About 7.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Loamy Upland (pe16-20)

*Land capability (irrigated):* 3e

*Land capability (nonirrigated):* 3e

#### *Typical Profile:*

H1—0 to 6 inches; silty clay loam

H2—6 to 19 inches; silty clay loam

H3—19 to 60 inches; silt loam

## **Rs—Richfield-Spearville complex, 0 to 1 percent slopes**

### **Map Unit Composition**

Richfield: 60 percent

Spearville: 40 percent

### **Component Descriptions**

#### **Richfield**

*MLRA:* 72 - Central High Tableland

*Landform:* Plain on tableland

*Parent material:* Loess

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* High (About 10.7 inches)

*Shrink-swell potential:* High (About 7.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Low

*Ecological site:* Loamy Upland (pe16-20)

*Land capability (irrigated):* 1

*Land capability (nonirrigated):* 2c

#### *Typical Profile:*

H1—0 to 6 inches; silt loam

H2—6 to 19 inches; silty clay loam

H3—19 to 60 inches; silt loam

#### **Spearville**

*MLRA:* 72 - Central High Tableland

*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 0 to 1 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Slow (About 0.06 in/hr)  
*Available water capacity:* High (About 10.7 inches)  
*Shrink-swell potential:* High (About 7.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Medium  
*Ecological site:* Clay Upland (pe16-20)  
*Land capability (irrigated):* 2s  
*Land capability (nonirrigated):* 2s

*Typical Profile:*

H1—0 to 7 inches; silty clay loam  
 H2—7 to 20 inches; silty clay  
 H3—20 to 26 inches; silty clay loam  
 H4—26 to 60 inches; silt loam

**Minor Components**  
**Ness**

**Sa—Satanta loam, 0 to 1 percent slopes**

**Map Unit Composition**

Satanta: 100 percent

**Component Descriptions**

**Satanta**

*MLRA:* 72 - Central High Tableland  
*Landform:* Sand sheet on paleoterrace on tableland  
*Parent material:* Loamy eolian deposits  
*Slope:* 0 to 1 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* Moderate (About 8.9 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Upland (pe16-20)  
*Land capability (irrigated):* 1  
*Land capability (nonirrigated):* 2c

*Typical Profile:*

H1—0 to 10 inches; loam  
 H2—10 to 32 inches; clay loam  
 H3—32 to 50 inches; silt loam

**Minor Components**  
**Ness**

**SAP—sand Pit**

**Sb—Satanta loam, 1 to 3 percent slopes**

**Map Unit Composition**

Satanta: 100 percent

**Component Descriptions**

**Satanta**

*MLRA:* 72 - Central High Tableland  
*Landform:* Sand sheet on paleoterrace on tableland  
*Parent material:* Loamy eolian deposits  
*Slope:* 1 to 3 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* Moderate (About 8.9 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Upland (pe16-20)  
*Land capability (irrigated):* 2e  
*Land capability (nonirrigated):* 2e

*Typical Profile:*

H1—0 to 10 inches; loam  
 H2—10 to 32 inches; clay loam  
 H3—32 to 50 inches; silt loam

## **Sp—Spearville silty clay loam, 0 to 1 percent slopes**

### **Map Unit Composition**

Spearville: 100 percent

### **Component Descriptions**

#### **Spearville**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Plain on tableland

*Parent material:* Loess

*Slope:* 0 to 1 percent

*Drainage class:* Well drained

*Slowest permeability:* Slow (About 0.06 in/hr)

*Available water capacity:* High (About 10.7 inches)

*Shrink-swell potential:* High (About 7.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* Medium

*Ecological site:* Clay Upland (pe16-20)

*Land capability (irrigated):* 2s

*Land capability (nonirrigated):* 2s

#### *Typical Profile:*

H1—0 to 7 inches; silty clay loam

H2—7 to 20 inches; silty clay

H3—20 to 26 inches; silty clay loam

H4—26 to 60 inches; silt loam

#### **Minor Components**

**Ness**

## **Sr—Spearville complex, 1 to 3 percent slopes, eroded**

### **Map Unit Composition**

Spearville: 100 percent

### **Component Descriptions**

#### **Spearville**

*MLRA:* 73 - Rolling Plains and Breaks

*Landform:* Plain on tableland

*Parent material:* Loess

*Slope:* 1 to 3 percent

*Drainage class:* Well drained

*Slowest permeability:* Slow (About 0.06 in/hr)

*Available water capacity:* High (About 10.7 inches)

*Shrink-swell potential:* High (About 7.5 LEP)

*Flooding hazard:* None

*Depth to seasonal water saturation:* More than 6 feet

*Runoff class:* High

*Ecological site:* Clay Upland (pe16-20)

*Land capability (nonirrigated):* 3e

#### *Typical Profile:*

H1—0 to 7 inches; silty clay loam

H2—7 to 20 inches; silty clay

H3—20 to 26 inches; silty clay loam

H4—26 to 60 inches; silt loam

## **Sw—SweetWater, occasionally flooded**

### **Map Unit Composition**

Sweetwater: 100 percent

### **Component Descriptions**

#### **Sweetwater**

*MLRA:* 72 - Central High Tableland

*Landform:* Flood plain on river valley

*Parent material:* Loamy alluvium

*Slope:* 0 to 1 percent

*Drainage class:* Poorly drained

*Slowest permeability:* Moderately slow (About 0.20 in/hr)

*Available water capacity:* Moderate (About 6.4 inches)

*Shrink-swell potential:* Low (About 1.5 LEP)

*Flooding hazard:* Occasional

*Depth to seasonal water saturation:* About 6 to 36 inches

*Runoff class:* Low

*Ecological site:* Saline Subirrigated (pe16-20)

*Land capability (nonirrigated):* 5w

#### *Typical Profile:*

H1—0 to 17 inches; clay loam

H2—17 to 60 inches; very fine sandy loam

## **Tf—Optima fine sand, 6 to 20 percent slopes**

### **Map Unit Composition**

Optima: 90 percent  
Minor components: 10 percent

### **Component Descriptions**

#### **Optima**

*MLRA:* 72 - Central High Tableland  
*Landform:* Dune on paleoterrace on river valley  
*Parent material:* Sandy eolian deposits  
*Slope:* 6 to 20 percent  
*Drainage class:* Excessively drained  
*Slowest permeability:* Rapid (About 6.00 in/hr)  
*Available water capacity:* Low (About 4.5 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Very low  
*Ecological site:* Sands (pe17-20)  
*Land capability (irrigated):* 4e  
*Land capability (nonirrigated):* 6e

#### *Typical Profile:*

A—0 to 8 inches; fine sand  
AC—8 to 17 inches; fine sand  
C—17 to 80 inches; fine sand

### **Minor Components**

#### **Eva**

*Composition:* About 10 percent  
*Slope:* 3 to 8 percent  
*Drainage class:* Somewhat excessively drained  
*Ecological site:* Sands (pe17-20)

### **Unnamed Hydric Soils**

## **Ua—Ulysses silt loam, 0 to 1 percent slopes**

### **Map Unit Composition**

Ulysses: 100 percent

### **Component Descriptions**

#### **Ulysses**

*MLRA:* 72 - Central High Tableland

*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 0 to 1 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* High (About 12.0 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Negligible  
*Ecological site:* Loamy Upland (pe16-20)  
*Land capability (irrigated):* 1  
*Land capability (nonirrigated):* 2c

#### *Typical Profile:*

H1—0 to 5 inches; silt loam  
H2—5 to 14 inches; silty clay loam  
H3—14 to 60 inches; silt loam

### **Minor Components**

#### **Ness**

## **Ub—Ulysses silt loam, 1 to 3 percent slopes**

### **Map Unit Composition**

Ulysses: 100 percent

### **Component Descriptions**

#### **Ulysses**

*MLRA:* 72 - Central High Tableland  
*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 1 to 3 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* High (About 12.0 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Upland (pe16-20)  
*Land capability (irrigated):* 2e



*Land capability (nonirrigated): 2e*

*Typical Profile:*

- H1—0 to 5 inches; silt loam
- H2—5 to 14 inches; silty clay loam
- H3—14 to 60 inches; silt loam

## **UBB—Uly silt loam, 1 to 3 percent slopes**

### **Map Unit Composition**

Uly: 100 percent

### **Component Descriptions**

#### **Uly**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 1 to 3 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* Very high (About 12.0 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Upland (pe20-26)  
*Land capability (irrigated):* 2e  
*Land capability (nonirrigated):* 2e

*Typical Profile:*

- H1—0 to 8 inches; silt loam
- H2—8 to 55 inches; silt loam
- H3—55 to 60 inches; silt loam

#### **Minor Components**

#### **Ness**

## **Uc—Ulysses silt loam, 3 to 6 percent slopes**

### **Map Unit Composition**

Ulysses: 100 percent

### **Component Descriptions**

#### **Ulysses**

*MLRA:* 72 - Central High Tableland  
*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 3 to 6 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* High (About 12.0 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Upland (pe16-20)  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

- H1—0 to 6 inches; silt loam
- H2—6 to 16 inches; silty clay loam
- H3—16 to 60 inches; silt loam

## **UCC—Uly silt loam, 3 to 6 percent slopes**

### **Map Unit Composition**

Uly: 100 percent

### **Component Descriptions**

#### **Uly**

*MLRA:* 73 - Rolling Plains and Breaks  
*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 3 to 6 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* Very high (About 12.2 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Upland (pe20-26)  
*Land capability (nonirrigated):* 3e

*Typical Profile:*

- H1—0 to 6 inches; silt loam

H2—6 to 14 inches; silty clay loam  
H3—14 to 60 inches; silt loam

### **Um—Ulysses-Colby silt loams, 3 to 6 percent slopes, eroded**

#### **Map Unit Composition**

Ulysses: 55 percent  
Colby: 45 percent

#### **Component Descriptions**

##### **Ulysses**

*MLRA:* 72 - Central High Tableland  
*Landform:* Plain on tableland  
*Parent material:* Loess  
*Slope:* 3 to 6 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* High (About 12.0 inches)  
*Shrink-swell potential:* Moderate (About 4.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Low  
*Ecological site:* Loamy Upland (pe16-20)

*Land capability (nonirrigated):* 4e

##### *Typical Profile:*

H1—0 to 5 inches; silt loam  
H2—5 to 14 inches; silty clay loam  
H3—14 to 60 inches; silt loam

##### **Colby**

*MLRA:* 72 - Central High Tableland  
*Landform:* Hillslope on tableland  
*Parent material:* Loess  
*Slope:* 3 to 6 percent  
*Drainage class:* Well drained  
*Slowest permeability:* Moderate (About 0.60 in/hr)  
*Available water capacity:* High (About 12.0 inches)  
*Shrink-swell potential:* Low (About 1.5 LEP)  
*Flooding hazard:* None  
*Depth to seasonal water saturation:* More than 6 feet  
*Runoff class:* Medium  
*Ecological site:* Limy Upland (pe16-20)  
*Land capability (irrigated):* 3e  
*Land capability (nonirrigated):* 4e

##### *Typical Profile:*

H1—0 to 5 inches; silt loam  
H2—5 to 60 inches; silt loam

### **W—Water**

Prime farmland is one of several kinds of important farmland defined by the U.S. Department of Agriculture. It is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil qualities, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. It is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

A recent trend in land use in some parts of the survey area has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

The map units in the survey area that are considered prime farmland are listed in the following table. This list does not constitute a recommendation for a particular land use. On some soils included in the list, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures. The extent of each listed map unit is shown in the "Acres and Proportionate Extent of Soils" table. The location is shown on the detailed soil maps. The soil qualities that affect use and management are described in other tables in this document."

Map symbol	Mapunit name	Farmland Classification
119HC	Harney silty clay loam, 1 to 3 percent slopes, eroded	All areas are prime farmland
119RX	Roxbury silt loam, rarely flooded	All areas are prime farmland
119UA	Uly silt loam, 0 to 1 percent slopes	All areas are prime farmland
119UT	Uly silt loam, 3 to 6 percent slopes, eroded	All areas are prime farmland
Da	Dale silt loam, rarely flooded	All areas are prime farmland
Ha	Harney silt loam, 0 to 1 percent slopes	All areas are prime farmland
HB	Harney silt loam, 1 to 3 percent slopes	All areas are prime farmland
Lu	Lubbock loam, 0 to 1 percent slopes	All areas are prime farmland
Rm	Richfield silt loam, 0 to 1 percent slopes	All areas are prime farmland
Sa	Satanta loam, 0 to 1 percent slopes	All areas are prime farmland
Sb	Satanta loam, 1 to 3 percent slopes	All areas are prime farmland
Sp	Spearville silty clay loam, 0 to 1 percent slopes	All areas are prime farmland
Sr	Spearville complex, 1 to 3 percent slopes, eroded	All areas are prime farmland
UBB	Uly silt loam, 1 to 3 percent slopes	All areas are prime farmland
UCC	Uly silt loam, 3 to 6 percent slopes	All areas are prime farmland
081LU	Lubbock silty clay loam, 0 to 1 percent slopes	Prime farmland if irrigated
Rn	Richfield silt loam, 1 to 3 percent slopes	Prime farmland if irrigated
Ro	Richfield silty clay loam, 1 to 3 percent slopes, eroded	Prime farmland if irrigated
Rs	Richfield-spearville complex, 0 to 1 percent slopes	Prime farmland if irrigated
Ua	Ulysses silt loam, 0 to 1 percent slopes	Prime farmland if irrigated
Ub	Ulysses silt loam, 1 to 3 percent slopes	Prime farmland if irrigated
Uc	Ulysses silt loam, 3 to 6 percent slopes	Prime farmland if irrigated

SOIL RATING FOR PLANT GROWTH, modified 1998  
Gray County, Kansas

PAGE 1 of 1

The "Soil Rating for Plant Growth, modified 1998" (SRPG) is a relative rating of the capacity of a soil to produce a specific plant under a defined management system. The index is determined from yield data on a few benchmark soils and is used to calculate yields, the net returns from crops, land assessment values, and taxes and to perform risk analysis when land management decisions are made. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

Map symbol	Soil name	Crop Index
055BP	Bridgeport Clay Loam, Rarely Flooded-----	49
055LD	Las-Las Animas Complex, Occasionally Flooded-----	33
055TF	Valent Fine Sand, 6 To 20 Percent Slopes-----	15
055TV	Valent-Vona Loamy Fine Sands, 3 To 8 Percent Slopes-----	29
055VO	Vona Loamy Fine Sand, 1 To 5 Percent Slopes-----	37
081LU	Lubbock Silty Clay Loam, 0 To 1 Percent Slopes-----	58
083UM	Uly-Coly Silt Loams, 3 To 6 Percent Slopes, Eroded-----	55
119HC	Harney Silty Clay Loam, 1 To 3 Percent Slopes, Eroded-----	63
119RX	Roxbury Silt Loam, Rarely Flooded-----	56
119UA	Uly Silt Loam, 0 To 1 Percent Slopes-----	48
119UT	Uly Silt Loam, 3 To 6 Percent Slopes, Eroded-----	59
ARR	Arkansas River-----	0
An	Bridgeport Silt Loam, Channeled-----	40
Bc	Bridgeport Silty Clay Loam, 0 To 1 Percent Slopes-----	54
Bd	Bridgeport Silty Clay Loam, 1 To 3 Percent Slopes-----	54
Bo	Optima Fine Sand, 0 To 10 Percent Slopes-----	23
Da	Dale Silt Loam, Rarely Flooded-----	62
GRP	Gravel Pit-----	18
Gr	Schamber Gravelly Sandy Loam, 5 To 25 Percent Slopes-----	14
HB	Harney Silt Loam, 1 To 3 Percent Slopes-----	65
Ha	Harney Silt Loam, 0 To 1 Percent Slopes-----	68
INL	Aquolls-----	12
La	Las Animas-Lesho Complex, Alkali, Occasionally Flooded-----	31
Lc	Las Animas Soils, Occasionally Flooded-----	29
Ld	Las Animas Sandy Loam, Occasionally Flooded-----	32
Le	Las Clay Loam, Occasionally Flooded-----	34
Lh	Lesho Clay Loam, Occasionally Flooded-----	37
Lk	Lesho-Sweetwater Complex, Occasionally Flooded-----	35
Ll	Lincoln Soils, Frequently Flooded-----	28
Lo	Lofton Silty Clay Loam, 0 To 1 Percent Slopes-----	55
Lu	Lubbock Loam, 0 To 1 Percent Slopes-----	62
Md	Penden Clay Loam, 3 To 6 Percent Slopes-----	30
Me	Penden Clay Loam, 6 To 15 Percent Slopes-----	26
Mf	Penden Clay Loam, 3 To 6 Percent Slopes, Severely Eroded-----	30
Mh	Penden Complex, 6 To 15 Percent Slopes, Severely Eroded-----	26
Mm	Manter Fine Sandy Loam, 0 To 1 Percent Slopes-----	49
Mn	Manter Fine Sandy Loam, 1 To 3 Percent Slopes-----	48
Mo	Manter Fine Sandy Loam, 1 To 6 Percent Slopes, Eroded-----	47
Mp	Campus-Canlon Complex, 6 To 15 Percent Slopes-----	10
Mu	Manter-Ulysses Complex, 1 To 6 Percent Slopes-----	48
Pa	Pratt Loamy Fine Sand, 1 To 6 Percent Slopes-----	34
Pg	Pratt Loamy Fine Sand, Gravel Substratum, 0 To 1 Percent Slopes-----	40
Pt	Pratt-Optima Loamy Fine Sands, 3 To 9 Percent Slopes-----	27
Ra	Ness Clay-----	10
Rm	Richfield Silt Loam, 0 To 1 Percent Slopes-----	58
Rn	Richfield Silt Loam, 1 To 3 Percent Slopes-----	59
Ro	Richfield Silty Clay Loam, 1 To 3 Percent Slopes, Eroded-----	58
Rs	Richfield-Spearville Complex, 0 To 1 Percent Slopes-----	59
SAP	Sand Pit-----	0
Sa	Satanta Loam, 0 To 1 Percent Slopes-----	64
Sb	Satanta Loam, 1 To 3 Percent Slopes-----	63
Sp	Spearville Silty Clay Loam, 0 To 1 Percent Slopes-----	59
Sr	Spearville Complex, 1 To 3 Percent Slopes, Eroded-----	58
Sw	Sweetwater Soils, Occasionally Flooded-----	32
Tf	Optima Fine Sand, 6 To 20 Percent Slopes-----	19
UBB	Uly Silt Loam, 1 To 3 Percent Slopes-----	64
UCC	Uly Silt Loam, 3 To 6 Percent Slopes-----	59
Ua	Ulysses Silt Loam, 0 To 1 Percent Slopes-----	52
Ub	Ulysses Silt Loam, 1 To 3 Percent Slopes-----	51
Uc	Ulysses Silt Loam, 3 To 6 Percent Slopes-----	47
Um	Ulysses-Colby Silt Loams, 3 To 6 Percent Slopes, Eroded-----	48
W	Water-----	0

Gray County, Kansas: Maintenance needed  
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
055BP:BRIDGEPORT	100	1-	2c	Not prime farmland	B	Loamy Terrace (pe16-20)	5	.28	.28	5	4L	86
055LD:LAS-----	55	2w-	3w	Not prime farmland	C	Saline Subirrigated (pe16-20)	5	.32	.32	4	4L	86
055LD:LAS ANIMAS	45	3w-	4w	Not prime farmland	C	Saline Subirrigated (pe16-20)	3	.24	.24	5	3	86
055TF:VALENT----	100	6e-	7e	Not prime farmland	A	Choppy Sands (pe16-20)	1	.15	.15	5	1	250
055TF:OPTIMA----	90	4e-	6e	Not prime farmland	A	Sands (pe17-20)	2	.17	.17	5	2	134
055TV:VALENT----	50	4e-	6e	Not prime farmland	A	Sands (pe16-20)	2	.17	.17	5	2	134
055TV:VONA-----	50	4e-	6e	Not prime farmland	B	Sands (pe16-20)	2	.17	.17	4	2	134
055VO:VONA-----	100	4e-	4e	Not prime farmland	B	Sands (pe16-20)	2	.17	.17	4	2	134
081LU:LUBBOCK---	100	1-	3c	Prime farmland if irrigated	B	Loamy Upland (pe16-20)	8	.32	.32	5	7	38
083UM:ULY-----	70	N/A	4e	Not prime farmland	B	Loamy Upland (pe20-26)	7	.32	.32	5	6	48
083UM:COLY-----	30	3e-	3e	Not prime farmland	B	Limy Upland (pe20-26)	5	.43	.43	5	4L	86
119HC:HARNEY----	100	N/A	3e	All areas are prime farmland	B	Loamy Upland (pe20-26)	8	.32	.32	5	7	38
119RX:ROXBURY---	100	1-	2c	All areas are prime farmland	B	Loamy Terrace (pe20-26)	5	.32	.32	5	4L	86
119UA:ULY-----	100	1-	2c	All areas are prime farmland	B	Loamy Upland (pe20-26)	7	.32	.32	5	6	48
119UT:ULY-----	100	N/A	4e	All areas are prime farmland	B	Limy Upland (pe20-26)	7	.32	.32	5	6	48
ARR:ARKANSAS RIVER-----	100	N/A	8w	Not prime farmland	D	Unspecified		---	---	-	---	0
An:BRIDGEPORT---	100	N/A	5w	Not prime farmland	B	Loamy Lowland (pe16-20)	5	.28	.28	5	4L	86
Bc:BRIDGEPORT---	100	1-	2c	Not prime farmland	B	Loamy Terrace (pe16-20)	5	.32	.32	5	4L	86

Gray County, Kansas: Maintenance needed  
Field Office Thunderbook: Soils Properties for Conservation Planning--Continued

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Bd:BRIDGEPORT---	100	2e-	3e	Not prime farmland	B	Loamy Terrace (pe16-20)	5	.32	.32	5	4L	86
Bo:OPTIMA-----	70	4e-	6e	Not prime farmland	A	Sands (pe17-20)	2	.17	.17	5	2	134
Da:DALE-----	100	N/A	2c	All areas are prime farmland	B	Loamy Terrace (pe16-20)	6	.32	.32	5	5	56
GRP:PITS-----	100	N/A	8s	Not prime farmland	A	Unspecified	9	.10	.17	2	8	0
Gr:SCHAMBER-----	100	N/A	7s	Not prime farmland	A	Gravelly Hills (pe16-20)	9	.17	.28	5	8	0
HB:HARNEY-----	100	2e-	2e	All areas are prime farmland	B	Loamy Upland (pe20-26)	7	.32	.32	5	6	48
Ha:HARNEY-----	100	1-	2c	All areas are prime farmland	B	Loamy Upland (pe20-26)	7	.32	.32	5	6	48
INL:AQUOLLS-----	100	N/A	5w	Not prime farmland	C	Unspecified		---	---	-	---	0
La:LAS ANIMAS---	50	3w-	4w	Not prime farmland	C	Saline Subirrigated (pe16-20)	3	.24	.24	5	3	86
La:LESHO-----	40	3s-	4s	Not prime farmland	C	Saline Subirrigated (pe16-20)	5	.28	.28	4	4L	86
Lc:LAS ANIMAS---	100	N/A	6s	Not prime farmland	C	Saline Subirrigated (pe16-20)	2	.17	.17	5	2	134
Ld:LAS ANIMAS---	100	3w-	3w	Not prime farmland	C	Saline Subirrigated (pe16-20)	3	.24	.24	5	3	86
Le:LESHO-----	100	N/A	3w	Not prime farmland	C	Saline Subirrigated (pe16-20)	5	.28	.28	5	4L	86
Le:LAS-----	100	2w-	4w	Not prime farmland	C	Saline Lowland (pe16-20)	5	.32	.32	4	4L	86
Lh:LESHO-----	100	N/A	3w	Not prime farmland	C	Saline Subirrigated (pe16-20)	5	.28	.28	5	4L	86
Lk:LESHO-----	50	N/A	6s	Not prime farmland	C	Saline Subirrigated (pe16-20)	5	.28	.28	5	4L	86
Lk:SWEETWATER---	25	N/A	5w	Not prime farmland	D	Saline Subirrigated (pe16-20)	5	.28	.28	3	4L	86
Ll:LINCOLN-----	100	N/A	7w	Not prime farmland	A	Sandy Lowland (pe16-20)	3	.24	.24	5	3	86

Gray County, Kansas: Maintenance needed  
Field Office Thunderbook: Soils Properties for Conservation Planning--Continued

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Lo:LOFTON-----	100	N/A	N/A	Not prime farmland	D	Unspecified	4	.37	.37	5	4	86
Lu:LUBBOCK-----	100	1-	2c	All areas are prime farmland	B	Loamy Upland (pe16-20)	7	.28	.28	5	6	48
Md:PENDEN-----	100	N/A	3e	Not prime farmland	B	Limy Upland (pe20-26)	5	.28	.28	5	4L	86
Me:PENDEN-----	100	N/A	6e	Not prime farmland	B	Limy Upland (pe16-20)	5	.28	.28	5	4L	86
Mf:PENDEN-----	100	N/A	6e	Not prime farmland	B	Limy Upland (pe16-20)	5	.28	.28	4	4L	86
Mh:PENDEN-----	100	N/A	6e	Not prime farmland	B	Limy Upland (pe16-20)	5	.28	.28	4	4L	86
Mm:MANTER-----	100	2s-	2e	Not prime farmland	B	Sandy (pe16-20)	3	.20	.20	5	3	86
Mn:MANTER-----	100	2e-	2e	Not prime farmland	B	Sandy (pe16-20)	3	.20	.20	5	3	86
Mo:MANTER-----	100	3e-	3e	Not prime farmland	B	Sandy (pe16-20)	3	.20	.20	5	3	86
Mp:CAMPUS-----	75	N/A	4e	Not prime farmland	B	Limy Upland (pe16-20)	5	.28	.28	2	4L	86
Mp:CANLON-----	25	N/A	6e	Not prime farmland	D	Shallow Limy (pe16-20)	5	.37	.37	1	4L	86
Mu:MANTER-----	55	3e-	3e	Not prime farmland	B	Sandy (pe16-20)	3	.20	.20	5	3	86
Mu:ULYSSES-----	45	3e-	3e	Not prime farmland	B	Loamy Upland (pe16-20)	7	.32	.32	5	6	48
Pa:PRATT-----	100	3e-	3e	Not prime farmland	A	Sands (pe16-20)	2	.17	.17	5	2	134
Pg:PRATT-----	100	3e-	4e	Not prime farmland	A	Sands (pe16-20)	2	.17	.17	5	2	134
Pt:PRATT-----	60	3e-	4e	Not prime farmland	A	Sands (pe16-20)	2	.17	.17	5	2	134
Pt:OPTIMA-----	40	4e-	6e	Not prime farmland	A	Sands (pe17-20)	2	.17	.17	5	2	134
Ra:NESS-----	100	N/A	6w	Not prime farmland	D	Lakebed (pe20- 26)	4	.28	.28	5	4	86
Rm:RICHFIELD----	100	1-	2c	All areas are prime farmland	B	Loamy Upland (pe16-20)	7	.32	.32	5	6	48
Rn:RICHFIELD----	100	2e-	2e	Prime farmland if irrigated	B	Loamy Upland (pe16-20)	7	.32	.32	5	6	48



Gray County, Kansas: Maintenance needed  
Field Office Thunderbook: Soils Properties for Conservation Planning--Continued

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Ro:RICHFIELD----	100	3e-	3e	Prime farmland if irrigated	B	Loamy Upland (pe16-20)	8	.32	.32	5	7	38
Rs:RICHFIELD----	60	1-	2c	Prime farmland if irrigated	B	Loamy Upland (pe16-20)	7	.32	.32	5	6	48
Rs:SPEARVILLE---	40	2s-	2s	Prime farmland if irrigated	C	Clay Upland (pe16-20)	8	.37	.37	5	7	38
SAP:SAND PIT----	100	N/A	N/A	Not prime farmland		Unspecified		---	---	-	---	---
Sa:SATANTA-----	100	1-	2c	All areas are prime farmland	B	Loamy Upland (pe16-20)	7	.28	.28	5	6	48
Sb:SATANTA-----	100	2e-	2e	All areas are prime farmland	B	Loamy Upland (pe16-20)	7	.28	.28	5	6	48
Sp:SPEARVILLE---	100	2s-	2s	All areas are prime farmland	C	Clay Upland (pe16-20)	8	.32	.32	5	7	38
Sr:SPEARVILLE---	100	N/A	3e	All areas are prime farmland	C	Clay Upland (pe16-20)	8	.32	.32	5	7	38
Sw:SWEETWATER---	100	N/A	5w	Not prime farmland	D	Saline Subirrigated (pe16-20)	5	.28	.28	3	4L	86
Tf:OPTIMA-----	90	4e-	6e	Not prime farmland	A	Sands (pe17-20)	2	.17	.17	5	2	134
UBB:ULY-----	100	2e-	2e	All areas are prime farmland	B	Loamy Upland (pe20-26)	7	.32	.32	5	6	48
UCC:ULY-----	100	N/A	3e	All areas are prime farmland	B	Loamy Upland (pe20-26)	7	.32	.32	5	6	48
Ua:ULYSSES-----	100	1-	2c	Prime farmland if irrigated	B	Loamy Upland (pe16-20)	7	.32	.32	5	6	48
Ub:ULYSSES-----	100	2e-	2e	Prime farmland if irrigated	B	Loamy Upland (pe16-20)	7	.32	.32	5	6	48
Uc:ULYSSES-----	100	3e-	3e	Prime farmland if irrigated	B	Loamy Upland (pe16-20)	7	.32	.32	5	6	48
Um:ULYSSES-----	55	N/A	4e	Not prime farmland	B	Loamy Upland (pe16-20)	7	.32	.32	5	6	48
Um:COLBY-----	45	3e-	4e	Not prime farmland	B	Limy Upland (pe16-20)	5	.43	.43	5	4L	86
W:WATER-----	100	N/A	N/A			Unspecified		---	---	-	---	---

RANGELAND PRODUCTIVITY  
Gray County, Kansas

### Use and Explanation of Rangeland, Grazed Forest Land, Native Pastureland Interpretations

Information in this subsection can be used to plan the use and management of soils for rangeland, grazed forest land, and native pasture. Different kinds of soils vary in their capacity to produce native grasses and other plants suitable for grazing. Information in this subsection provides groupings of similar soils and estimates of potential forage production, which can be used to determine livestock stocking rates.

**Rangeland.** Range is land on which the native vegetation (climax or natural potential plant community) is predominantly grasses, grasslike plants, forbs, and shrubs suitable for grazing and browsing. Range includes natural grasslands, savannas, many wetlands, some deserts, tundra, and certain shrub and forb communities. Rangeland receives no regular or frequent cultural treatment. The composition and production of the plant community are determined by soil, climate, topography, overstory canopy, and grazing management.

**Grazed Forest Land.** Includes land on which the understory includes, as an integral part of the forest plant community, plants that can be grazed without significantly impairing other forest values.

**Native Pasture.** Includes land on which the native vegetation (climax or natural potential plant community) is forest but which is used and managed primarily for production of native plants for forage. Native pasture includes cut-over forest land and forest land cleared and now managed for native or naturalized forage plants.

#### Rangeland

In areas that have similar climate and topography, differences in the kind and amount of vegetation produced on rangeland are closely related to the kind of soil. Effective management based on the relationship between the soils and vegetation and water.

The Rangeland, Grazed Forest land, Native Pastureland Interpretations shows, for each soil that supports rangeland vegetation, the ecological site and the potential annual production of vegetation in favorable, normal, unfavorable years. An explanation of the column headings in this table follows.

An ecological site is the product of all the environmental factors responsible for its development. It has characteristic soils that have developed over time throughout the soil development process; a characteristic hydrology, particularly infiltration and runoff, that has developed over time; and a characteristic plant community (kind and amount of vegetation). The hydrology of a site is influenced by development of the soil and plant community. The vegetation, soils, and hydrology are all interrelated. Each is influenced by the others and influences the development of the others. The plant community on an ecological site is typified by an association of species that differs from that of other ecological sites in the kind and/or proportion of species or in total production. Descriptions of ecological sites are provided in the Field Office Technical Guide, which is available in local offices of the Natural Resources Conservation Service.

Total dry-weight production is the amount of vegetation that can be expected to grow annually on well managed rangeland that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, average, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are about average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Yields are adjusted to a common percent of air-dry moisture content.

Range management requires a knowledge of the kinds of soil and of the potential natural plant community. It also requires an evaluation of the present range similarity index and rangeland trend. Range similarity index is determined by comparing the present plant community with the potential natural plant community on a particular rangeland ecological site. The more closely the existing community resembles the potential community, the higher the range similarity index. Rangeland trend is defined as the direction of change in an existing plant community relative to the potential natural plant community. Further information about the range similarity index and rangeland trend is available in chapter 4 of the National Range and Pasture Handbook, which is available in local offices of the Natural Resources Conservation Service. The objective in range management is to control grazing so that the plants growing on a site are about the same in kind and amount as the potential natural plant community for that site. Such management generally results in the optimum production of vegetation, control of undesirable brush species, conservation of water, and control of erosion. Sometimes, however, an area with a range similarity index somewhat below the potential meets grazing needs, provides wildlife habitat, and protects soil and water resources.

RANGELAND PRODUCTIVITY--Continued  
Gray County, Kansas

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
055BP: Bridgeport, rarely flooded-----	Loamy Terrace (pe16-20)	4,000	3,000	2,000
055LD: Las, occasionally flooded-----	Saline Subirrigated (pe16-20)	6,000	5,000	4,000
Las Animas, occasionally flooded--	Saline Subirrigated (pe16-20)	3,000	2,500	1,000
055TF: Valent-----	Choppy Sands (pe16-20)	2,500	2,000	1,200
055TV: Valent-----	Sands (pe16-20)	2,500	2,000	1,200
Vona-----	Sands (pe16-20)	2,200	1,700	900
055VO: Vona-----	Sands (pe16-20)	2,200	1,700	900
081LU: Lubbock-----	Loamy Upland (pe16-20)	2,800	2,000	1,200
083UM: Uly-----	Loamy Upland (pe20-26)	3,700	3,200	2,700
Coly-----	Limy Upland (pe20-26)	3,300	3,000	2,700
119HC: Harney, eroded-----	Loamy Upland (pe20-26)	4,000	2,200	1,000
119RX: Roxbury-----	Loamy Terrace (pe20-26)	4,000	3,000	2,000
119UA: Uly-----	Loamy Upland (pe20-26)	3,700	3,200	2,700
119UT: Uly-----	Limy Upland (pe20-26)	3,700	3,200	2,700
An: Bridgeport, channeled-----	Loamy Lowland (pe16-20)	6,000	4,500	3,500
ARR: Arkansas River-----	---	---	---	---
Bc: Bridgeport-----	Loamy Terrace (pe16-20)	4,000	3,000	2,000
Bd: Bridgeport-----	Loamy Terrace (pe16-20)	4,000	3,000	2,000
Bo: Optima-----	Sands (pe17-20)	2,250	1,250	900
Da: Dale-----	Loamy Terrace (pe16-20)	8,500	6,100	4,500
Gr: Schamber-----	Gravelly Hills (pe16-20)	1,800	1,300	700
GRP: Pits-----	---	---	---	---
Ha: Harney-----	Loamy Upland (pe20-26)	4,000	2,200	1,000
HB: Harney-----	Loamy Upland (pe20-26)	4,000	2,200	1,000
INL: Aquolls-----	---	---	---	---
La: Las Animas, occasionally flooded--	Saline Subirrigated (pe16-20)	4,500	3,750	2,500
Lesho, occasionally flooded-----	Saline Subirrigated (pe16-20)	6,500	5,500	4,000
Lc: Las Animas, occasionally flooded--	Saline Subirrigated (pe16-20)	5,000	4,500	3,250
Ld: Las Animas-----	Saline Subirrigated (pe16-20)	5,000	4,500	3,250
Le: Las, occasionally flooded-----	Saline Lowland (pe16-20)	6,000	5,000	4,000
Lh: Lesho, occasionally flooded-----	Saline Subirrigated (pe16-20)	9,000	8,000	7,000
Lk: Lesho-----	Saline Subirrigated (pe16-20)	9,000	8,000	7,000
Sweetwater-----	Saline Subirrigated (pe16-20)	5,000	4,250	3,500
LL: Lincoln, occasionally flooded----	Sandy Lowland (pe16-20)	3,000	2,300	1,800
Lo: Lofton-----	---	---	---	---
Lu: Lubbock-----	Loamy Upland (pe16-20)	3,500	2,500	1,500
Md: Penden-----	Limy Upland (pe20-26)	4,000	2,500	1,000
Me: Penden-----	Limy Upland (pe16-20)	4,000	2,500	1,000
Mf: Penden, Severely eroded-----	Limy Upland (pe16-20)	4,000	2,500	1,000
Mh: Penden, Severely Eroded-----	Limy Upland (pe16-20)	4,000	2,500	1,000
Mm: Manter-----	Sandy (pe16-20)	2,000	1,600	800
Mn: Manter-----	Sandy (pe16-20)	2,000	1,600	800
Mo: Manter-----	Sandy (pe16-20)	2,000	1,600	800
Mp: Campus-----	Limy Upland (pe16-20)	3,000	2,000	1,000
Canlon-----	Shallow Limy (pe16-20)	2,400	1,600	900
Mu:				

RANGELAND PRODUCTIVITY--Continued  
Gray County, Kansas

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
Manter-----	Sandy (pe16-20)	2,000	1,600	800
Ulysses-----	Loamy Upland (pe16-20)	2,400	1,800	1,000
Pa:				
Pratt-----	Sands (pe16-20)	4,500	3,500	2,500
Pg:				
Pratt-----	Sands (pe16-20)	4,500	3,500	2,500
Pt:				
Pratt-----	Sands (pe16-20)	4,500	3,500	2,500
Optima-----	Sands (pe17-20)	2,250	1,250	900
Ra:				
Ness-----	Lakebed (pe20-26)	2,000	1,500	500
Rm:				
Richfield-----	Loamy Upland (pe16-20)	2,400	1,800	800
Rn:				
Richfield-----	Loamy Upland (pe16-20)	2,400	1,800	800
Ro:				
Richfield-----	Loamy Upland (pe16-20)	2,400	1,800	800
Rs:				
Richfield-----	Loamy Upland (pe16-20)	2,400	1,800	800
Spearville-----	Clay Upland (pe16-20)	2,000	1,500	800
Sa:				
Satanta-----	Loamy Upland (pe16-20)	3,000	2,000	1,000
SAP:				
Sand Pit-----	---	---	---	---
Sb:				
Satanta-----	Loamy Upland (pe16-20)	3,000	2,000	1,000
Sp:				
Spearville-----	Clay Upland (pe16-20)	2,000	1,500	800
Sr:				
Spearville-----	Clay Upland (pe16-20)	2,000	1,500	800
Sw:				
Sweetwater-----	Saline Subirrigated (pe16-20)	5,000	4,250	3,500
Tf:				
Optima-----	Sands (pe17-20)	2,250	1,250	900
Ua:				
Ulysses-----	Loamy Upland (pe16-20)	2,400	1,800	1,000
Ub:				
Ulysses-----	Loamy Upland (pe16-20)	2,400	1,800	1,000
UBB:				
Uly-----	Loamy Upland (pe20-26)	3,700	3,200	2,700
Uc:				
Ulysses-----	Loamy Upland (pe16-20)	2,400	1,800	1,000
UCC:				
Uly-----	Loamy Upland (pe20-26)	3,700	3,200	2,700
Um:				
Ulysses-----	Loamy Upland (pe16-20)	2,400	1,800	1,000
Colby-----	Limy Upland (pe16-20)	2,400	1,800	1,000
W:				
Water-----	---	---	---	---

BUILDING SITE DEVELOPMENT  
Gray County, Kansas

### Building Site Development

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. The following tables show the degree and kind of soil limitations that affect dwellings with and without basements, small commercial buildings, local roads and streets, shallow excavations, and lawns and landscaping.

The ratings in the tables are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Local roads and streets have an all-weather surface and carry automobile and light truck traffic all year. They have a subgrade of cut or fill soil material; a base of gravel, crushed rock, or soil material stabilized by lime or cement; and a surface of flexible material (asphalt), rigid material (concrete), or gravel with a binder. The ratings are based on the soil properties that affect the ease of excavation and grading and the traffic-supporting capacity. The properties that affect the ease of excavation and grading are depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, depth to a water table, ponding, flooding, the amount of large stones, and slope. The properties that affect the traffic-supporting capacity are soil strength (as inferred from the AASHTO group index number), subsidence, linear extensibility (shrink-swell potential), the potential for frost action, depth to a water table, and ponding.

Shallow excavations are trenches or holes dug to a maximum depth of 5 or 6 feet for graves, utility lines, open ditches, or other purposes. The ratings are based on the soil properties that influence the ease of digging and the resistance to sloughing. Depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, the amount of large stones, and dense layers influence the ease of digging, filling, and compacting. Depth to the seasonal high water table, flooding, and ponding may restrict the period when excavations can be made. Slope influences the ease of using machinery. Soil texture, depth to the water table, and linear extensibility (shrink-swell potential) influence the resistance to sloughing.

Lawns and landscaping require soils on which turf and ornamental trees and shrubs can be established and maintained. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer.

BUILDING SITE DEVELOPMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
055BP: Bridgeport, rarely flooded-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
055LD: Las, occasionally flooded-----	55	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Shrink-swell	1.00 0.50
Las Animas, occasionally flooded-----	45	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
055TF: Valent-----	100	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
055TV: Valent-----	50	Not limited		Not limited		Somewhat limited Slope	0.86
Vona-----	50	Not limited		Not limited		Somewhat limited Slope	0.48
055VO: Vona-----	100	Not limited		Not limited		Not limited	
081LU: Lubbock-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
083UM: Uly-----	70	Not limited		Not limited		Somewhat limited Slope	0.12
Coly-----	30	Not limited		Not limited		Somewhat limited Slope	0.12
119HC: Harney, eroded-----	100	Not limited		Not limited		Not limited	
119RX: Roxbury-----	100	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 0.50
119UA: Uly-----	100	Not limited		Not limited		Not limited	
119UT: Uly-----	100	Not limited		Not limited		Somewhat limited Slope	0.12
An: Bridgeport, channeled-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
ARR: Arkansas River-----	100	Not rated		Not rated		Not rated	
Bc: Bridgeport-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Bd: Bridgeport-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Bo: Optima-----	70	Not limited		Not limited		Somewhat limited Slope	0.00
Da: Dale-----	100	Very limited Flooding	1.00	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding	1.00
Gr: Schamber-----	100	Not rated		Not rated		Not rated	
GRP: Pits-----	100	Not rated		Not rated		Not rated	
Ha: Harney-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
HB: Harney-----	100	Not limited		Not limited		Not limited	

BUILDING SITE DEVELOPMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
INL: Aquolls-----	100	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
La: Las Animas, occasionally flooded-----	50	Very limited  Flooding	 1.00	Very limited  Flooding Depth to saturated zone	 1.00 0.95	Very limited  Flooding	 1.00
Lesho, occasionally flooded-----	40	Very limited  Flooding	 1.00	Very limited  Flooding Depth to saturated zone	 1.00 0.95	Very limited  Flooding	 1.00
Lc: Las Animas, occasionally flooded-----	100	Very limited  Flooding Depth to saturated zone	 1.00 0.07	Very limited  Flooding Depth to saturated zone	 1.00 1.00	Very limited  Flooding Depth to saturated zone	 1.00 0.07
Ld: Las Animas-----	100	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
Le: Las, occasionally flooded-----	100	Very limited  Flooding Shrink-swell	 1.00 0.50	Very limited  Flooding Depth to saturated zone	 1.00 1.00	Very limited  Flooding Shrink-swell	 1.00 0.50
Lh: Lesho, occasionally flooded-----	100	Very limited  Flooding	 1.00	Very limited  Flooding Depth to saturated zone	 1.00 0.95	Very limited  Flooding	 1.00
Lk: Lesho-----	50	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.95	Very limited Flooding	1.00
Sweetwater-----	25	Very limited Flooding Depth to saturated zone	1.00 0.81	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.81
Ll: Lincoln, occasionally flooded-----	100	Very limited  Flooding	 1.00	Very limited  Flooding Depth to saturated zone	 1.00 0.03	Very limited  Flooding	 1.00
Lo: Lofton-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
Lu: Lubbock-----	100	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
Md: Penden-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
Me: Penden-----	100	Somewhat limited Shrink-swell Slope	0.50 0.37	Somewhat limited Shrink-swell Slope	0.50 0.37	Very limited Slope Shrink-swell	1.00 0.50
Mf: Penden, Severely eroded-----	100	Somewhat limited  Shrink-swell	 0.50	Somewhat limited  Shrink-swell	 0.50	Somewhat limited  Shrink-swell Slope	 0.50 0.00

BUILDING SITE DEVELOPMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Mh: Penden, Severely Eroded-----	100	Somewhat limited  Shrink-swell Slope	0.50 0.37	Somewhat limited  Shrink-swell Slope	0.50 0.37	Very limited  Slope Shrink-swell	1.00 0.50
Mm: Manter-----	100	Not limited		Not limited		Not limited	
Mn: Manter-----	100	Not limited		Not limited		Not limited	
Mo: Manter-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Mp: Campus-----	75	Very limited Depth to hard bedrock	1.00	Very limited Depth to hard bedrock	1.00	Very limited Depth to hard bedrock Slope	1.00 0.86
Canlon-----	25	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00
Mu: Manter-----	55	Not limited		Not limited		Somewhat limited Slope	0.00
Ulysses-----	45	Not limited		Not limited		Somewhat limited Slope	0.00
Pa: Pratt-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Pg: Pratt-----	100	Not limited		Not limited		Not limited	
Pt: Pratt-----	60	Not limited		Not limited		Somewhat limited Slope	0.48
Optima-----	40	Not limited		Not limited		Somewhat limited Slope	0.86
Ra: Ness-----	100	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00
Rm: Richfield-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Rn: Richfield-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Ro: Richfield-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Rs: Richfield-----	60	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Spearville-----	40	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Sa: Satanta-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
SAP: Sand Pit-----	100	Not rated		Not rated		Not rated	
Sb: Satanta-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Sp: Spearville-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Sr: Spearville-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Sw: Sweetwater-----	100	Very limited Flooding Depth to saturated zone	1.00 0.81	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.81



BUILDING SITE DEVELOPMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Tf: Optima-----	90	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
Ua: Ulysses-----	100	Not limited		Not limited		Not limited	
Ub: Ulysses-----	100	Not limited		Not limited		Not limited	
UBB: Uly-----	100	Not limited		Not limited		Not limited	
Uc: Ulysses-----	100	Not limited		Not limited		Somewhat limited Slope	0.12
UCC: Uly-----	100	Not limited		Not limited		Somewhat limited Slope	0.12
Um: Ulysses-----	55	Not limited		Not limited		Somewhat limited Slope	0.12
Colby-----	45	Not limited		Not limited		Somewhat limited Slope	0.12
W: Water-----	100	Not rated		Not rated		Not rated	

BUILDING SITE DEVELOPMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
055BP: Bridgeport, rarely flooded-----	100	Somewhat limited		Somewhat limited		Not limited	
		Frost action	0.50	Cutbanks cave	0.10		
		Flooding	0.40				
055LD: Las, occasionally flooded-----	55	Very limited		Very limited		Somewhat limited	
		Flooding	1.00	Cutbanks cave	1.00	Flooding	0.60
		Shrink-swell	0.50	Depth to saturated zone	1.00		
		Frost action	0.50	Flooding	0.60		
Las Animas, occasionally flooded-----	45	Very limited		Very limited		Somewhat limited	
		Flooding	1.00	Cutbanks cave	1.00	Flooding	0.60
		Frost action	0.50	Depth to saturated zone	1.00	Depth to saturated zone	0.03
		Depth to saturated zone	0.03	Flooding	0.60		
055TF: Valent-----	100	Very limited Slope	1.00	Very limited Cutbanks cave Slope	1.00 1.00	Very limited Slope Droughty	1.00 0.34
055TV: Valent-----	50	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.25
Vona-----	50	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
055VO: Vona-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
081LU: Lubbock-----	100	Very limited Shrink-swell	1.00	Somewhat limited Cutbanks cave	0.10	Not limited	
083UM: Uly-----	70	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Coly-----	30	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
119HC: Harney, eroded-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
119RX: Roxbury-----	100	Very limited Low strength Shrink-swell Frost action Flooding	1.00 0.50 0.50 0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
119UA: Uly-----	100	Very limited Low strength Frost action	1.00 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
119UT: Uly-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
An: Bridgeport, channeled-----	100	Very limited		Somewhat limited		Very limited	
		Flooding	1.00	Flooding	0.80	Flooding	1.00
		Frost action	0.50	Cutbanks cave	0.10		
ARR: Arkansas River-----	100	Not rated		Not rated		Not rated	
Bc: Bridgeport-----	100	Somewhat limited Frost action Flooding	0.50 0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
Bd: Bridgeport-----	100	Somewhat limited Frost action Flooding	0.50 0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
Bo: Optima-----	70	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.38

BUILDING SITE DEVELOPMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Da: Dale-----	100	Somewhat limited Frost action Flooding	0.50 0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
Gr: Schamber-----	100	Not rated		Not rated		Not rated	
GRP: Pits-----	100	Not rated		Not rated		Not rated	
Ha: Harney-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
HB: Harney-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
INL: Aquolls-----	100	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding Cutbanks cave	1.00 1.00 0.10	Very limited Depth to saturated zone Ponding	1.00 1.00
La: Las Animas, occasionally flooded-----	50	Very limited  Flooding Frost action	 1.00 0.50	Very limited  Cutbanks cave Depth to saturated zone Flooding	 1.00 0.95 0.60	Somewhat limited  Flooding Salinity	 0.60 0.13
Lesho, occasionally flooded-----	40	Very limited  Flooding Frost action	 1.00 0.50	Very limited  Cutbanks cave Depth to saturated zone Flooding	 1.00 0.95 0.60	Somewhat limited  Flooding Salinity	 0.60 0.13
LC: Las Animas, occasionally flooded-----	100	Very limited  Flooding Frost action Depth to saturated zone	 1.00 0.50 0.03	Very limited  Cutbanks cave Depth to saturated zone Flooding	 1.00 1.00 0.60	Somewhat limited  Flooding Depth to saturated zone	 0.60 0.03
Ld: Las Animas-----	100	Very limited Flooding Frost action Depth to saturated zone	1.00 0.50 0.03	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 1.00 0.60	Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Le: Las, occasionally flooded-----	100	Very limited  Flooding Shrink-swell Frost action	 1.00 0.50 0.50	Very limited  Cutbanks cave Depth to saturated zone Flooding	 1.00 1.00 0.60	Somewhat limited  Flooding	 0.60
Lh: Lesho, occasionally flooded-----	100	Very limited  Flooding Frost action	 1.00 0.50	Very limited  Cutbanks cave Depth to saturated zone Flooding	 1.00 0.95 0.60	Somewhat limited  Flooding	 0.60

BUILDING SITE DEVELOPMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Lk: Lesho-----	50	Very limited Flooding Frost action	1.00 0.50	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 0.95 0.60	Somewhat limited Flooding	0.60
Sweetwater-----	25	Very limited Flooding Depth to saturated zone	1.00 0.48	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 1.00 0.60	Somewhat limited Flooding Depth to saturated zone	0.60 0.48
Ll: Lincoln, occasionally flooded-----	100	Very limited  Flooding	 1.00	Very limited  Cutbanks cave Flooding Depth to saturated zone	 1.00 0.80 0.03	Very limited  Flooding	 1.00
Lo: Lofton-----	100	Very limited Shrink-swell	1.00	Somewhat limited Too clayey Cutbanks cave	0.12 0.10	Not limited	
Lu: Lubbock-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Md: Penden-----	100	Very limited Low strength Shrink-swell Frost action	1.00 0.50 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Me: Penden-----	100	Somewhat limited Shrink-swell Frost action Slope	0.50 0.50 0.37	Somewhat limited Slope Cutbanks cave	0.37 0.10	Somewhat limited Slope	0.37
Mf: Penden, Severely eroded-----	100	Somewhat limited  Shrink-swell Frost action	 0.50 0.50	Somewhat limited  Cutbanks cave	 0.10	Not limited	
Mh: Penden, Severely Eroded-----	100	Somewhat limited  Shrink-swell Frost action Slope	 0.50 0.50 0.37	Somewhat limited  Slope Cutbanks cave	 0.37 0.10	Somewhat limited  Slope	 0.37
Mm: Manter-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Mn: Manter-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Mo: Manter-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Mp: Campus-----	75	Very limited Depth to hard bedrock	1.00	Very limited Depth to hard bedrock Cutbanks cave	1.00 0.10	Very limited Depth to bedrock	1.00
Canlon-----	25	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Cutbanks cave Slope	1.00 0.10 0.04	Droughty Very limited Depth to bedrock Droughty Slope	0.08 1.00 1.00 0.04
Mu: Manter-----	55	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Ulysses-----	45	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Pa: Pratt-----	100	Not limited		Very limited Cutbanks cave	1.00	Not limited	
Pg: Pratt-----	100	Not limited		Very limited Cutbanks cave	1.00	Not limited	

BUILDING SITE DEVELOPMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Pt: Pratt-----	60	Not limited		Very limited Cutbanks cave	1.00	Not limited	
Optima-----	40	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.38
Ra: Ness-----	100	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Cutbanks cave Too clayey	1.00 1.00 1.00 0.50	Very limited Ponding Depth to saturated zone Too clayey	1.00 1.00 1.00
Rm: Richfield-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Rn: Richfield-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Ro: Richfield-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Rs: Richfield-----	60	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Spearville-----	40	Somewhat limited Shrink-swell	0.50	Somewhat limited Too clayey Cutbanks cave	0.18 0.10	Not limited	
Sa: Satanta-----	100	Somewhat limited Low strength Shrink-swell Frost action	0.78 0.50 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
SAP: Sand Pit-----	100	Not rated		Not rated		Not rated	
Sb: Satanta-----	100	Somewhat limited Low strength Shrink-swell Frost action	0.78 0.50 0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Sp: Spearville-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Too clayey Cutbanks cave	0.18 0.10	Not limited	
Sr: Spearville-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Too clayey Cutbanks cave	0.18 0.10	Not limited	
Sw: Sweetwater-----	100	Very limited Flooding  Depth to saturated zone	1.00  0.48	Very limited Depth to saturated zone Flooding Cutbanks cave	1.00  0.60 0.10	Somewhat limited Flooding  Depth to saturated zone	0.60  0.48
Tf: Optima-----	90	Somewhat limited Slope	0.63	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope Droughty	0.63 0.38
Ua: Ulysses-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Ub: Ulysses-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
UBB: Uly-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Uc: Ulysses-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
UCC: Uly-----	100	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	

BUILDING SITE DEVELOPMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Um:							
Ulysses-----	55	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Colby-----	45	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
W:							
Water-----	100	Not rated		Not rated		Not rated	

CONSTRUCTION MATERIALS  
Gray County, Kansas

### Construction Materials

The following tables give information about the soils as potential sources of gravel, sand, topsoil, reclamation material, and roadfill. Normal compaction, minor processing, and other standard construction practices are assumed.

The soils are rated good, fair, or poor as potential sources of topsoil, reclamation material, and roadfill. The features that limit the soils as sources of these materials are specified in the tables. The numerical ratings given after the specified features indicate the degree to which the features limit the soils as sources of topsoil, reclamation material, or roadfill. The lower the number, the greater the limitation.

The soils are rated as a probable or improbable source of sand and gravel. A rating of probable means that the source material is likely to be in or below the soil. The numerical ratings in these columns indicate the degree of probability. The number 0.00 indicates that the soil is an improbable source. A number between 0.00 and 1.00 indicates the degree to which the soil is a probable source of sand or gravel.

Sand and gravel are natural aggregates suitable for commercial use with a minimum of processing. They are used in many kinds of construction. Specifications for each use vary widely. In these tables, only the probability of finding material in suitable quantity is evaluated. The suitability of the material for specific purposes is not evaluated, nor are factors that affect excavation of the material. The properties used to evaluate the soil as a source of sand or gravel are gradation of grain sizes (as indicated by the Unified classification of the soil), the thickness of suitable material, and the content of rock fragments. If the lowest layer of the soil contains sand or gravel, the soil is rated as a probable source regardless of thickness. The assumption is that the sand or gravel layer below the depth of observation exceeds the minimum thickness.

Topsoil is used to cover an area so that vegetation can be established and maintained. The upper 40 inches of a soil is evaluated for use as topsoil. Also evaluated is the reclamation potential of the borrow area. The ratings are based on the soil properties that affect plant growth; the ease of excavating, loading, and spreading the material; and reclamation of the borrow area. Toxic substances, soil reaction, and the properties that are inferred from soil texture, such as available water capacity and fertility, affect plant growth. The ease of excavating, loading, and spreading is affected by rock fragments, slope, depth to a water table, soil texture, and thickness of suitable material. Reclamation of the borrow area is affected by slope, depth to a water table, rock fragments, depth to bedrock or a cemented pan, and toxic material.

The surface layer of most soils is generally preferred for topsoil because of its organic matter content. Organic matter greatly increases the absorption and retention of moisture and nutrients for plant growth.

Reclamation material is used in areas that have been drastically disturbed by surface mining or similar activities. When these areas are reclaimed, layers of soil material or unconsolidated geological material, or both, are replaced in a vertical sequence. The reconstructed soil favors plant growth. The ratings in the table do not apply to quarries and other mined areas that require an offsite source of reconstruction material. The ratings are based on the soil properties that affect erosion and stability of the surface and the productive potential of the reconstructed soil. These properties include the content of sodium, salts, and calcium carbonate; reaction; available water capacity; erodibility; texture; content of rock fragments; and content of organic matter and other features that affect fertility.

Roadfill is soil material that is excavated in one place and used in road embankments in another place. In this table, the soils are rated as a source of roadfill for low embankments, generally less than 6 feet high and less exacting in design than higher embankments.

The ratings are for the whole soil, from the surface to a depth of about 5 feet. It is assumed that soil layers will be mixed when the soil material is excavated and spread.

The ratings are based on the amount of suitable material and on soil properties that affect the ease of excavation and the performance of the material after it is in place. The thickness of the suitable material is a major consideration. The ease of excavation is affected by large stones, depth to a water table, and slope. How well the soil performs in place after it has been compacted and drained is determined by its strength (as inferred from the AASHTO classification of the soil) and linear extensibility (shrink-swell potential).

CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
055BP: Bridgeport, rarely flooded-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
055LD: Las, occasionally flooded-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.93
Las Animas, occasionally flooded-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.07
055TF: Valent-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Bottom layer	0.99
055TV: Valent-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.65 0.99
Vona-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.08 0.09
055VO: Vona-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.09
081LU: Lubbock-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
083UM: Uly-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Coly-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
119HC: Harney, eroded-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
119RX: Roxbury-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
119UA: Uly-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
119UT: Uly-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
An: Bridgeport, channeled-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00



CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
ARR: Arkansas River-----	100	Not rated		Not rated	
Bc: Bridgeport-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Bd: Bridgeport-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Bo: Optima-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.99 0.99
Da: Dale-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Gr: Schamber-----	100	Not rated		Not rated	
GRP: Pits-----	100	Not rated		Not rated	
Ha: Harney-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
HB: Harney-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
INL: Aquolls-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
La: Las Animas, occasionally flooded-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.96
Lesho, occasionally flooded-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.93
Lc: Las Animas, occasionally flooded-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.07
Ld: Las Animas-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.07
Le: Las, occasionally flooded-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.93

CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Lh: Lesho, occasionally flooded-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.96
Lk: Lesho-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.96
Sweetwater-----	25	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.87
Ll: Lincoln, occasionally flooded-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.09
Lo: Lofton-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Lu: Lubbock-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Md: Penden-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Me: Penden-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Mf: Penden, Severely eroded-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Mh: Penden, Severely Eroded-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Mm: Manter-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.08
Mn: Manter-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.08 0.09
Mo: Manter-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.08 0.09
Mp: Campus-----	75	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Canlon-----	25	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Mu: Manter-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.08 0.09
Ulysses-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Pa: Pratt-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.57
Pg: Pratt-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.57
Pt: Pratt-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.57
Optima-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.99 0.99
Ra: Ness-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rm: Richfield-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rn: Richfield-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ro: Richfield-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rs: Richfield-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Spearville-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Sa: Satanta-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
SAP: Sand Pit-----	100	Not rated		Not rated	
Sb: Satanta-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Sp: Spearville-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Sr: Spearville-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Sw: Sweetwater-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.08
Tf: Optima-----	90	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.99 0.99
Ua: Ulysses-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ub: Ulysses-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
UBB: Uly-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Uc: Ulysses-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
UCC: Uly-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Um: Ulysses-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Colby-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
W: Water-----	100	Not rated		Not rated	

CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
055BP: Bridgeport, rarely flooded-----	100	Fair Water erosion	0.90	Good		Good	
055LD: Las, occasionally flooded-----	55	Fair Low content of organic matter	0.02	Fair Depth to saturated zone	0.89	Fair Depth to saturated zone	0.89
Las Animas, occasionally flooded-----	45	Fair Low content of organic matter	0.12	Fair Depth to saturated zone	0.76	Fair Depth to saturated zone	0.76
055TF: Valent-----	100	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.08 0.25	Fair Slope	0.68	Poor Too sandy Slope	0.00 0.00
055TV: Valent-----	50	Poor Too sandy Wind erosion Low content of organic matter Droughty	0.00 0.00 0.08 0.77	Good		Poor Too sandy	0.00
Vona-----	50	Poor Wind erosion Low content of organic matter	0.00 0.50	Good		Good	
055VO: Vona-----	100	Poor Wind erosion Low content of organic matter	0.00 0.50	Good		Good	
081LU: Lubbock-----	100	Poor Too clayey Low content of organic matter	0.00 0.82	Fair Shrink-swell	0.67	Poor Too Clayey	0.00
083UM: Uly-----	70	Fair Low content of organic matter Water erosion	0.18 0.90	Good		Good	
Coly-----	30	Fair Low content of organic matter Water erosion	0.18 0.90	Good		Good	
119HC: Harney, eroded-----	100	Fair Low content of organic matter Water erosion	0.50 0.90	Good		Good	
119RX: Roxbury-----	100	Fair Water erosion Low content of organic matter	0.90 0.92	Poor Low strength Shrink-swell	0.00 0.96	Good	
119UA: Uly-----	100	Poor Too alkaline Low content of organic matter Water erosion	0.00 0.18 0.90	Poor Low strength	0.00	Good	

CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
119UT: Uly-----	100	Fair Low content of organic matter Water erosion	0.18 0.90	Good		Good	
An: Bridgeport, channeled-----	100	Fair Water erosion Low content of organic matter	0.90 0.92	Good		Good	
ARR: Arkansas River-----	100	Not rated		Not rated		Not rated	
Bc: Bridgeport-----	100	Fair Water erosion	0.90	Good		Good	
Bd: Bridgeport-----	100	Fair Water erosion	0.90	Good		Good	
Bo: Optima-----	70	Poor Too sandy Wind erosion Low content of organic matter	0.00 0.00 0.00	Good		Poor Too sandy	0.00
Da: Dale-----	100	Fair Low content of organic matter Water erosion	0.50 0.90	Fair Shrink-swell	0.98	Good	
Gr: Schamber-----	100	Not rated		Not rated		Not rated	
GRP: Pits-----	100	Not rated		Not rated		Not rated	
Ha: Harney-----	100	Fair Too clayey Low content of organic matter Water erosion	0.05 0.50 0.90	Good		Fair Too Clayey	0.03
HB: Harney-----	100	Fair Low content of organic matter Water erosion	0.50 0.90	Good		Good	
INL: Aguolls-----	100	Poor Low content of organic matter	0.00	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone	0.00
La: Las Animas, occasionally flooded-----	50	Poor Too alkaline Low content of organic matter Too sandy	0.00 0.50 0.68	Good		Fair Salinity Too sandy	0.50 0.68
Lesho, occasionally flooded-----	40	Poor Low content of organic matter Salinity Too clayey	0.00 0.88 0.95	Good		Fair Too Clayey Salinity	0.84 0.88

CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Lc: Las Animas, occasionally flooded-----	100	Poor Wind erosion  Low content of organic matter Droughty	0.00  0.12 0.99	Fair Depth to saturated zone	0.76	Fair Depth to saturated zone	0.76
Ld: Las Animas-----	100	Fair Low content of organic matter	0.12	Fair Depth to saturated zone	0.76	Fair Depth to saturated zone	0.76
Le: Las, occasionally flooded-----	100	Fair Low content of organic matter	0.02	Fair Depth to saturated zone	0.89	Fair Depth to saturated zone	0.89
Lh: Lesho, occasionally flooded-----	100	Fair Low content of organic matter Too clayey	0.08 0.95	Good		Fair Too Clayey	0.84
Lk: Lesho-----	50	Poor Low content of organic matter	0.00	Good		Good	
Sweetwater-----	25	Poor Too sandy  Low content of organic matter	0.00 0.18	Fair Depth to saturated zone	0.29	Poor Too sandy  Depth to saturated zone	0.00 0.29
Ll: Lincoln, occasionally flooded-----	100	Fair Low content of organic matter	0.08	Good		Good	
Lo: Lofton-----	100	Poor Too clayey Water erosion	0.00 0.90	Fair Shrink-swell	0.84	Poor Too Clayey	0.00
Lu: Lubbock-----	100	Poor Too clayey Low content of organic matter	0.00 0.32	Good		Poor Too Clayey	0.00
Md: Penden-----	100	Fair Low content of organic matter Carbonate content	0.18 0.80	Poor Low strength  Shrink-swell	0.00 0.87	Fair Carbonate content	0.80
Me: Penden-----	100	Fair Low content of organic matter Carbonate content	0.18 0.80	Fair Shrink-swell	0.87	Fair Slope  Carbonate content	0.63 0.80
Mf: Penden, Severely eroded-----	100	Fair Low content of organic matter Carbonate content	0.18 0.80	Fair Shrink-swell	0.87	Fair Carbonate content	0.80

CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Mh: Penden, Severely Eroded-----	100	Fair Low content of organic matter Carbonate content	0.18 0.80	Fair Shrink-swell	0.87	Fair Slope Carbonate content	0.63 0.80
Mm: Manter-----	100	Fair Low content of organic matter	0.02	Good		Fair Rock fragments	0.97
Mn: Manter-----	100	Fair Low content of organic matter	0.02	Good		Fair Rock fragments	0.97
Mo: Manter-----	100	Fair Low content of organic matter	0.02	Good		Fair Rock fragments	0.97
Mp: Campus-----	75	Poor Depth to bedrock Droughty Low content of organic matter Carbonate content	0.00 0.09 0.50 0.80	Poor Depth to bedrock	0.00	Poor Depth to bedrock Carbonate content	0.00 0.80
Canlon-----	25	Poor Droughty Depth to bedrock Low content of organic matter No water erosion limitation	0.00 0.00 0.88 0.99	Poor Depth to bedrock	0.00	Poor Depth to bedrock Slope Rock fragments	0.00 0.96 0.97
Mu: Manter-----	55	Fair Low content of organic matter	0.02	Good		Fair Rock fragments	0.97
Ulysses-----	45	Fair Low content of organic matter Water erosion	0.18 0.90	Good		Good	
Pa: Pratt-----	100	Poor Wind erosion Too sandy Low content of organic matter	0.00 0.00 0.02	Good		Poor Too sandy	0.00
Pg: Pratt-----	100	Poor Wind erosion Too sandy Low content of organic matter	0.00 0.00 0.08	Good		Poor Too sandy	0.00
Pt: Pratt-----	60	Poor Wind erosion Too sandy Low content of organic matter	0.00 0.00 0.02	Good		Poor Too sandy	0.00
Optima-----	40	Poor Too sandy Wind erosion Low content of organic matter	0.00 0.00 0.00	Good		Poor Too sandy	0.00



CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ra: Ness-----	100	Poor Too clayey Water erosion	0.00  0.90	Poor Depth to saturated zone Shrink-swell	0.00  0.56	Poor Too Clayey Depth to saturated zone	0.00  0.00
Rm: Richfield-----	100	Fair Low content of organic matter Water erosion	0.18  0.90	Fair Shrink-swell	0.87	Good	
Rn: Richfield-----	100	Fair Low content of organic matter Water erosion	0.18  0.90	Fair Shrink-swell	0.76	Good	
Ro: Richfield-----	100	Fair Low content of organic matter Water erosion	0.18  0.90	Fair Shrink-swell	0.76	Good	
Rs: Richfield-----	60	Fair Low content of organic matter Water erosion	0.18  0.90	Fair Shrink-swell	0.76	Good	
Spearville-----	40	Fair Low content of organic matter Too clayey Water erosion	0.12  0.88 0.90	Fair Shrink-swell	0.81	Fair Too Clayey	0.51
Sa: Satanta-----	100	Fair Low content of organic matter	0.92	Fair Low strength Shrink-swell	0.22 0.99	Good	
SAP: Sand Pit-----	100	Not rated		Not rated		Not rated	
Sb: Satanta-----	100	Fair Low content of organic matter	0.92	Fair Low strength Shrink-swell	0.22 0.99	Good	
Sp: Spearville-----	100	Fair Low content of organic matter Too clayey Water erosion	0.12  0.88 0.90	Fair Shrink-swell	0.81	Fair Too Clayey	0.51
Sr: Spearville-----	100	Fair Low content of organic matter Too clayey Water erosion	0.12  0.88 0.90	Fair Shrink-swell	0.81	Fair Too Clayey	0.51
Sw: Sweetwater-----	100	Fair Low content of organic matter	0.18	Fair Depth to saturated zone	0.29	Fair Depth to saturated zone	0.29
Tf: Optima-----	90	Poor Too sandy Wind erosion Low content of organic matter	0.00 0.00 0.00	Good		Poor Too sandy Slope	0.00 0.37

CONSTRUCTION MATERIALS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ua: Ulysses-----	100	Fair Low content of organic matter Water erosion	0.18 0.90	Good		Good	
Ub: Ulysses-----	100	Fair Low content of organic matter Water erosion	0.18 0.90	Good		Good	
UBB: Uly-----	100	Fair Water erosion	0.90	Good		Good	
Uc: Ulysses-----	100	Fair Low content of organic matter Water erosion	0.18 0.90	Good		Good	
UCC: Uly-----	100	Fair Low content of organic matter Water erosion	0.18 0.90	Good		Good	
Um: Ulysses-----	55	Fair Low content of organic matter Water erosion	0.18 0.90	Good		Good	
Colby-----	45	Fair Low content of organic matter Water erosion	0.18 0.90	Good		Good	
W: Water-----	100	Not rated		Not rated		Not rated	

RECREATIONAL INTERPRETATIONS  
Gray County, Kansas

### Recreation

The soils of the survey area are rated in the following tables according to limitations that affect their suitability for recreation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the recreational uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The ratings in the tables are based on restrictive soil features, such as wetness, slope, and texture of the surface layer. Susceptibility to flooding is considered. Not considered in the ratings, but important in evaluating a site, are the location and accessibility of the area, the size and shape of the area and its scenic quality, vegetation, access to water, potential water impoundment sites, and access to public sewer lines. The capacity of the soil to absorb septic tank effluent and the ability of the soil to support vegetation also are important. Soils that are subject to flooding are limited for recreational uses by the duration and intensity of flooding and the season when flooding occurs. In planning recreational facilities, onsite assessment of the height, duration, intensity, and frequency of flooding is essential.

The information in this table can be supplemented by other information in this survey, for example, interpretations for building site development, construction materials, sanitary facilities, and water management.

Camp areas require site preparation, such as shaping and leveling the tent and parking areas, stabilizing roads and intensively used areas, and installing sanitary facilities and utility lines. Camp areas are subject to heavy foot traffic and some vehicular traffic. The ratings are based on the soil properties that affect the ease of developing camp areas and the performance of the areas after development. Slope, stoniness, and depth to bedrock or a cemented pan are the main concerns affecting the development of camp areas.

The soil properties that affect the performance of the areas after development are those that influence trafficability and promote the growth of vegetation, especially in heavily used areas. For good trafficability, the surface of camp areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Picnic areas are subject to heavy foot traffic. Most vehicular traffic is confined to access roads and parking areas. The ratings are based on the soil properties that affect the ease of developing picnic areas and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of picnic areas. For good trafficability, the surface of picnic areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Playgrounds require soils that are nearly level, are free of stones, and can withstand intensive foot traffic. The ratings are based on the soil properties that affect the ease of developing playgrounds and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of playgrounds. For good trafficability, the surface of the playgrounds should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Paths and trails for hiking and horseback riding should require little or no slope modification through cutting and filling. The ratings are based on the soil properties that affect trafficability and erodibility. These properties are stoniness, depth to a water table, ponding, flooding, slope, and texture of the surface layer.

Golf fairways are subject to heavy foot traffic and some light vehicular traffic. Cutting or filling may be required. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer. The suitability of the soil for traps, tees, roughs, and greens is not considered in the ratings.

RECREATIONAL INTERPRETATIONS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
055BP: Bridgeport, rarely flooded-----	100	Very limited Flooding	1.00	Not limited		Not limited	
055LD: Las, occasionally flooded-----	55	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
Las Animas, occasionally flooded-----	45	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Flooding	0.60
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
055TF: Valent-----	100	Very limited Too sandy Slope	1.00 1.00	Very limited Too sandy Slope	1.00 1.00	Very limited Slope Too sandy	1.00 1.00
055TV: Valent-----	50	Somewhat limited Too sandy	0.95	Somewhat limited Too sandy	0.95	Very limited Slope Too sandy	1.00 0.95
Vona-----	50	Somewhat limited Too sandy	0.34	Somewhat limited Too sandy	0.34	Very limited Slope Too sandy	1.00 0.34
055VO: Vona-----	100	Somewhat limited Too sandy	0.34	Somewhat limited Too sandy	0.34	Somewhat limited Too sandy Slope	0.34 0.13
081LU: Lubbock-----	100	Not limited		Not limited		Not limited	
083UM: Uly-----	70	Not limited		Not limited		Somewhat limited Slope	0.87
Coly-----	30	Not limited		Not limited		Somewhat limited Slope	0.87
119HC: Harney, eroded-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
119RX: Roxbury-----	100	Very limited Flooding	1.00	Not limited		Not limited	
119UA: Uly-----	100	Not limited		Not limited		Not limited	
119UT: Uly-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
An: Bridgeport, channeled-----	100	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
ARR: Arkansas River-----	100	Not rated		Not rated		Not rated	
Bc: Bridgeport-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Bd: Bridgeport-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Slope	0.00
Bo: Optima-----	70	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy Slope	0.96 0.50
Da: Dale-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Gr: Schamber-----	100	Not rated		Not rated		Not rated	
GRP: Pits-----	100	Not rated		Not rated		Not rated	

RECREATIONAL INTERPRETATIONS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ha: Harney-----	100	Not limited		Not limited		Not limited	
HB: Harney-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
INL: Aguolls-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Restricted permeability	1.00
		Restricted permeability	1.00	Restricted permeability	1.00	Depth to saturated zone	1.00
		Ponding	1.00	Ponding	1.00	Ponding	1.00
La: Las Animas, occasionally flooded-----	50	Very limited		Somewhat limited		Somewhat limited	
		Flooding Salinity	1.00 0.13	Salinity	0.13	Flooding Salinity	0.60 0.13
Lesho, occasionally flooded-----	40	Very limited		Somewhat limited		Somewhat limited	
		Flooding Salinity	1.00 0.13	Salinity	0.13	Flooding Salinity	0.60 0.13
LC: Las Animas, occasionally flooded-----	100	Very limited		Somewhat limited		Somewhat limited	
		Flooding Too sandy	1.00 0.81	Too sandy Depth to saturated zone	0.81 0.03	Too sandy Flooding	0.81 0.60
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
Ld: Las Animas-----	100	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Flooding	0.60
		Depth to saturated zone	0.07			Depth to saturated zone	0.07
Le: Las, occasionally flooded-----	100	Very limited		Not limited		Somewhat limited	
		Flooding	1.00			Flooding	0.60
Lh: Lesho, occasionally flooded-----	100	Very limited		Not limited		Somewhat limited	
		Flooding	1.00			Flooding	0.60
Lk: Lesho-----	50	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
Sweetwater-----	25	Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.48	Somewhat limited Depth to saturated zone	0.81
		Depth to saturated zone	0.81	Restricted permeability	0.15	Flooding	0.60
		Restricted permeability	0.15			Restricted permeability	0.15
Ll: Lincoln, occasionally flooded-----	100	Very limited		Somewhat limited		Very limited	
		Flooding	1.00	Flooding	0.40	Flooding Slope	1.00 0.00
Lo: Lofton-----	100	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45
Lu: Lubbock-----	100	Not limited		Not limited		Not limited	
Md: Penden-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
Me: Penden-----	100	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope	1.00
Mf: Penden, Severely eroded-----	100	Not limited		Not limited		Somewhat limited Slope	0.50
Mh: Penden, Severely Eroded-----	100	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Slope	1.00

RECREATIONAL INTERPRETATIONS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Mm: Manter-----	100	Not limited		Not limited		Somewhat limited Gravel content	0.06
Mn: Manter-----	100	Not limited		Not limited		Somewhat limited Gravel content Slope	0.06 0.00
Mo: Manter-----	100	Not limited		Not limited		Somewhat limited Slope Gravel content	0.50 0.06
Mp: Campus-----	75	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Slope	1.00 1.00
Canlon-----	25	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Slope Gravel content	1.00 1.00 0.06
Mu: Manter-----	55	Not limited		Not limited		Somewhat limited Slope Gravel content	0.50 0.06
Ulysses-----	45	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Slope Dusty	0.50 0.50
Pa: Pratt-----	100	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy	0.37	Somewhat limited Slope Too sandy	0.50 0.37
Pg: Pratt-----	100	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy	0.37
Pt: Pratt-----	60	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy	0.37	Very limited Slope Too sandy	1.00 0.37
Optima-----	40	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy	0.96	Very limited Slope Too sandy	1.00 0.96
Ra: Ness-----	100	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
		Too clayey Restricted permeability	0.50 0.45	Too clayey Restricted permeability	0.50 0.45	Too clayey Restricted permeability	0.50 0.45
Rm: Richfield-----	100	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50
Rn: Richfield-----	100	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Dusty Slope	0.50 0.00
Ro: Richfield-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Rs: Richfield-----	60	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50
Spearville-----	40	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39
Sa: Satanta-----	100	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50
SAP: Sand Pit-----	100	Not rated		Not rated		Not rated	
Sb: Satanta-----	100	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Dusty Slope	0.50 0.00
Sp: Spearville-----	100	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39
Sr: Spearville-----	100	Somewhat limited		Somewhat limited		Somewhat limited	

RECREATIONAL INTERPRETATIONS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Sw: Sweetwater-----	100	Restricted permeability	0.39	Restricted permeability	0.39	Restricted permeability Slope	0.39 0.00
		Very limited Flooding	1.00	Somewhat limited Depth to saturated zone	0.48	Somewhat limited Depth to saturated zone	0.81
		Depth to saturated zone	0.81	Restricted permeability	0.15	Flooding	0.60
		Restricted permeability	0.15			Restricted permeability	0.15
Tf: Optima-----	90	Very limited Too sandy Slope	1.00 0.63	Very limited Too sandy Slope	1.00 0.63	Very limited Too sandy Slope	1.00 1.00
Ua: Ulysses-----	100	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50
Ub: Ulysses-----	100	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Dusty Slope	0.50 0.00
UBB: Uly-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Uc: Ulysses-----	100	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Slope Dusty	0.87 0.50
UCC: Uly-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
Um: Ulysses-----	55	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Slope Dusty	0.87 0.50
Colby-----	45	Somewhat limited Dusty	0.50	Somewhat limited Dusty	0.50	Somewhat limited Slope Dusty	0.87 0.50
W: Water-----	100	Not rated		Not rated		Not rated	

RECREATIONAL INTERPRETATIONS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
055BP: Bridgeport, rarely flooded-----	100	Not limited		Not limited	
055LD: Las, occasionally flooded-----	55	Not limited		Somewhat limited	
Las Animas, occasionally flooded-----	45	Not limited		Flooding Somewhat limited	0.60
				Flooding Depth to saturated zone	0.60 0.03
055TF: Valent-----	100	Very limited Too sandy Slope	1.00 0.32	Very limited Slope Droughty	1.00 0.34
055TV: Valent-----	50	Somewhat limited Too sandy	0.95	Somewhat limited Droughty	0.25
Vona-----	50	Somewhat limited Too sandy	0.34	Not limited	
055VO: Vona-----	100	Somewhat limited Too sandy	0.34	Not limited	
081LU: Lubbock-----	100	Not limited		Not limited	
083UM: Uly-----	70	Not limited		Not limited	
Coly-----	30	Not limited		Not limited	
119HC: Harney, eroded-----	100	Not limited		Not limited	
119RX: Roxbury-----	100	Not limited		Not limited	
119UA: Uly-----	100	Not limited		Not limited	
119UT: Uly-----	100	Not limited		Not limited	
An: Bridgeport, channeled-----	100	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
ARR: Arkansas River-----	100	Not rated		Not rated	
Bc: Bridgeport-----	100	Not limited		Not limited	
Bd: Bridgeport-----	100	Not limited		Not limited	
Bo: Optima-----	70	Somewhat limited Too sandy	0.96	Somewhat limited Droughty	0.38
Da: Dale-----	100	Not limited		Not limited	
Gr: Schamber-----	100	Not rated		Not rated	
GRP: Pits-----	100	Not rated		Not rated	
Ha: Harney-----	100	Not limited		Not limited	
HB: Harney-----	100	Not limited		Not limited	
INL: Aguolls-----	100	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
La: Las Animas, occasionally flooded-----	50	Not limited		Somewhat limited	



RECREATIONAL INTERPRETATIONS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Lesho, occasionally flooded-----	40	Not limited		Flooding Salinity Somewhat limited	0.60 0.13
Lc: Las Animas, occasionally flooded-----	100	Somewhat limited		Flooding Salinity Somewhat limited	0.60 0.13
		Too sandy	0.81	Flooding Depth to saturated zone	0.60 0.03
Ld: Las Animas-----	100	Not limited		Somewhat limited Flooding Depth to saturated zone	0.60 0.03
Le: Las, occasionally flooded-----	100	Not limited		Somewhat limited	
Lh: Lesho, occasionally flooded-----	100	Not limited		Flooding	0.60
Lk: Lesho-----	50	Not limited		Somewhat limited Flooding	0.60
Sweetwater-----	25	Somewhat limited Depth to saturated zone	0.11	Somewhat limited Flooding	0.60
				Depth to saturated zone	0.48
Ll: Lincoln, occasionally flooded-----	100	Somewhat limited		Very limited	
		Flooding	0.40	Flooding	1.00
Lo: Lofton-----	100	Not limited		Not limited	
Lu: Lubbock-----	100	Not limited		Not limited	
Md: Penden-----	100	Not limited		Not limited	
Me: Penden-----	100	Not limited		Somewhat limited Slope	0.37
Mf: Penden, Severely eroded-----	100	Not limited		Not limited	
Mh: Penden, Severely Eroded-----	100	Not limited		Somewhat limited Slope	0.37
Mm: Manter-----	100	Not limited		Not limited	
Mn: Manter-----	100	Not limited		Not limited	
Mo: Manter-----	100	Not limited		Not limited	
Mp: Campus-----	75	Not limited		Very limited Depth to bedrock Droughty	1.00 0.08
Canlon-----	25	Not limited		Very limited Depth to bedrock Droughty Slope	1.00 1.00 0.04
Mu: Manter-----	55	Not limited		Not limited	
Ulysses-----	45	Somewhat limited Dusty	0.50	Not limited	
Pa: Pratt-----	100	Somewhat limited Too sandy	0.37	Not limited	
Pg: Pratt-----	100	Somewhat limited Too sandy	0.37	Not limited	
Pt: Pratt-----	60	Somewhat limited		Not limited	

RECREATIONAL INTERPRETATIONS--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Optima-----	40	Too sandy	0.37		
		Somewhat limited		Somewhat limited	
		Too sandy	0.96	Droughty	0.38
Ra:					
Ness-----	100	Very limited		Very limited	
		Depth to	1.00	Ponding	1.00
		saturated zone			
		Ponding	1.00	Depth to	1.00
				saturated zone	
		Too clayey	0.50	Too clayey	1.00
Rm:					
Richfield-----	100	Somewhat limited		Not limited	
		Dusty	0.50		
Rn:					
Richfield-----	100	Somewhat limited		Not limited	
		Dusty	0.50		
Ro:					
Richfield-----	100	Not limited		Not limited	
Rs:					
Richfield-----	60	Somewhat limited		Not limited	
		Dusty	0.50		
Spearville-----	40	Not limited		Not limited	
Sa:					
Satanta-----	100	Somewhat limited		Not limited	
		Dusty	0.50		
SAP:					
Sand Pit-----	100	Not rated		Not rated	
Sb:					
Satanta-----	100	Somewhat limited		Not limited	
		Dusty	0.50		
Sp:					
Spearville-----	100	Not limited		Not limited	
Sr:					
Spearville-----	100	Not limited		Not limited	
Sw:					
Sweetwater-----	100	Somewhat limited		Somewhat limited	
		Depth to	0.11	Flooding	0.60
		saturated zone			
				Depth to	0.48
				saturated zone	
Tf:					
Optima-----	90	Very limited		Somewhat limited	
		Too sandy	1.00	Slope	0.63
				Droughty	0.38
Ua:					
Ulysses-----	100	Somewhat limited		Not limited	
		Dusty	0.50		
Ub:					
Ulysses-----	100	Somewhat limited		Not limited	
		Dusty	0.50		
UBB:					
Uly-----	100	Not limited		Not limited	
Uc:					
Ulysses-----	100	Somewhat limited		Not limited	
		Dusty	0.50		
UCC:					
Uly-----	100	Not limited		Not limited	
Um:					
Ulysses-----	55	Somewhat limited		Not limited	
		Dusty	0.50		
Colby-----	45	Somewhat limited		Not limited	
		Dusty	0.50		
W:					
Water-----	100	Not rated		Not rated	

WILDLIFE INTERPRETATIONS  
Gray County, Kansas

#### Use and Explanation of Wildlife Interpretations

Soils directly affect the kind and amount of vegetation that is available to wildlife as food and cover. They also affect the development of water impoundments. The kind and abundance of wildlife that populate an area depend largely on the amount and distribution of food, cover, water, and living space. If any one of these elements is missing, inadequate, or inaccessible, wildlife will be scarce or will not inhabit the area. If the soils have the potential, wildlife habitat can be created or improved by planting appropriate vegetation, properly managing the existing plant cover, and fostering the natural establishment of desirable plants.

In the Wildlife Interpretations table, the soils in the survey area are rated according to their potential for providing habitat for various kinds of wildlife. This information can be used in planning parks, wildlife refuges, nature study areas, and other developments for wildlife; in selecting soils that are suitable for establishing, improving, or maintaining specific elements of wildlife habitat; and in determining the intensity of management needed for each element of the habitat.

#### Suitability Ratings

The potential of the soil is rated good, fair, poor, or very poor.

Good - means that the element of wildlife habitat or the kind of habitat is easily created, improved, or maintained. Few or no limitations affect management, and satisfactory results can be expected if the soil is used for the designated purpose.

Fair - means that the element of wildlife habitat or kind of habitat can be created, improved, or maintained in most places. Moderately intensive management is required for satisfactory results.

Poor - means that limitations are severe for the designated element or kind of wildlife habitat. Habitat can be created, improved, or maintained in most places, but management is difficult and requires intensive effort.

Very Poor - means that limitations are very severe for the designated element or kind of wildlife habitat. Habitat is difficult to create, improve, or maintain in most places, and management is difficult and requires intensive effort.

#### Description of Wildlife Habitat Elements

Openland habitat consists of croplands, pastures, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. The kind of wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, killdeer, cottontail rabbit, red fox, and coyote.

Woodland habitat consists of hardwood or conifers, or a mixture of these and associated grasses, legumes and wild herbaceous plants. Examples of wildlife attracted to this habitat are wild turkey, thrushes, woodpeckers, owl, tree squirrels, raccoon, and deer.

Wetland habitat consists of water-tolerant plants in open, marshy or swampy, shallow water areas. Examples of wildlife attracted to this habitat are ducks, geese, herons, bitterns, rails, kingfishers, shorebirds, muskrat, mink, and beaver.

The elements of wildlife habitat are described in the following paragraphs.

Grain and seed crops are domestic grains and seed-producing herbaceous plants. Soil properties and features that affect the growth of grain and seed crops are depth of the root zone, texture of the surface layer, available water capacity, wetness, slope, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of grain and seed crops are corn, wheat, oats, and barley.

Grasses and legumes are domestic perennial grasses and herbaceous legumes. Soil properties and features that affect the growth of grasses and legumes are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, flooding, and slope. Soil temperature and soil moisture also are considerations. Examples of grasses and legumes are fescue, lovegrass, bromegrass, clover, and alfalfa.

Wild herbaceous plants are native or naturally established grasses and forbs, including weeds. Soil properties and features that affect the growth of these plants are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of wild herbaceous plants are bluestem, goldenrod, beggarweed, wheatgrass, and grama.

Hardwood trees and woody understory produce nuts or other fruit, buds, catkins, twigs, bark, and foliage. Soil properties and features that affect the growth of hardwood trees and shrubs are depth of the root zone, available water capacity, and wetness. Examples of these plants are oak, poplar, cherry, sweetgum, apple, hawthorn, dogwood, hickory, blackberry, and blueberry. Examples of fruit-producing shrubs that are suitable for planting on soils rated good are Russian-olive, autumn-olive, and crabapple.

Coniferous plants furnish browse and seeds. Soil properties and features that affect the growth of coniferous trees, shrubs, and ground cover are depth of the root zone, available water capacity, and wetness. Examples of coniferous plants are pine, spruce, fir, cedar, and juniper.

Shrubs are bushy woody plants that produce fruit, buds, twigs, bark, and foliage. Soil properties and features that affect the growth of shrubs are depth of the root zone, available water capacity, salinity, and soil moisture. Examples of shrubs are fragrant sumac, chokecherry, American plum, sand plum, and gorden currant.

Wetland plants are annual and perennial wild herbaceous plants that grow on moist or wet sites. Submerged or floating aquatic plants are excluded. Soil properties and features affecting wetland plants are texture of the surface layer, wetness, reaction, salinity, slope, and surface stoniness. Examples of wetland plants are smartweed, wild millet, saltgrass, cordgrass, rushes, sedges, and cattails.

Shallow water areas have an average depth of less than 5 feet. Some are naturally wet areas. Others are created by dams, levees, or other water-control structures. Soil properties and features affecting shallow water areas are depth to bedrock, wetness, surface stoniness, slope, and permeability. Examples of shallow water areas are marshes, waterfowl feeding areas, and ponds.

The habitat for various kinds of wildlife is described in the following paragraphs.

Habitat for openland wildlife consists of cropland, pasture, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. Wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, cottontail, red fox and coyote.

Habitat for woodland wildlife consists of areas of deciduous and/or coniferous plants and associated grasses, legumes, and wild herbaceous plants. Wildlife attracted to these areas include wild turkey, thrushes, woodpeckers, squirrels, gray fox, raccoon, and deer.

Habitat for wetland wildlife consists of open, marshy or swampy shallow water areas. Some of the wildlife attracted to such areas are ducks, geese, herons, shore birds, muskrat, mink, and beaver.

Habitat for rangeland wildlife consists of areas of shrubs and wild herbaceous plants. Wildlife attracted to rangeland include antelope, deer, cottontail rabbit, prairie chicken, meadowlark, quail, and pheasant.

WILDLIFE INTERPRETATIONS  
Gray County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
055BP: BRIDGEPORT-----	Good	Good	Good	---	---	Fair	Poor	Poor	Good	---	Poor	Fair
055LD: LAS-----	Fair	Fair	Fair	---	---	Fair	Fair	Fair	Fair	---	Fair	Fair
LAS ANIMAS-----	Fair	Good	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair	Good
055TF: VALENT-----	Poor	Fair	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
055TV: VALENT-----	Poor	Fair	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
VONA-----	Fair	Fair	Fair	---	---	Fair	Poor	Very poor	Fair	---	Very poor	Fair
055VO: VONA-----	Fair	Fair	Fair	---	---	Fair	Poor	Very poor	Fair	---	Very poor	Fair
081LU: LUBBOCK-----	Fair	Fair	Fair	---	---	Poor	Poor	Fair	Fair	---	Poor	Fair
083UM: ULY-----	Fair	Good	Good	Good	Fair	Fair	Very poor	Very poor	Fair	Good	Very poor	Good
COLY-----	Fair	Good	Good	Good	Fair	Fair	Very poor	Very poor	Fair	Good	Very poor	Fair
119HC: HARNEY-----	Good	Good	Good	Poor	Poor	Good	Poor	Fair	Good	---	Poor	Good
119RX: ROXBURY-----	Good	Good	Good	Fair	Fair	Fair	Poor	Fair	Good	Fair	Poor	Fair
119UA: ULY-----	Good	Good	Good	Good	Fair	Fair	Very poor	Very poor	Good	Good	Very poor	Good
119UT: ULY-----	Fair	Good	Good	Good	Fair	Fair	Very poor	Very poor	Fair	Good	Very poor	Good
An: BRIDGEPORT-----	Fair	Good	Fair	---	---	Poor	Poor	Poor	Fair	---	Poor	Poor
ARR: ARKANSAS RIVER--	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Poor	Very poor	Very poor	Poor	Very poor
Bc: BRIDGEPORT-----	Good	Good	Good	---	---	Fair	Poor	Poor	Good	---	Poor	Fair
Bd: BRIDGEPORT-----	Fair	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
Bo: OPTIMA-----	Poor	Fair	Fair	Poor	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor	Fair
Da: DALE-----	Good	Good	Fair	---	---	Good	Poor	Very poor	Good	---	Very poor	Fair
Gr: SCHAMBER-----	Very poor	Very poor	Poor	Poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Poor
GRP: PITS-----	Very poor	Very poor	Poor	Poor	Poor	Poor	Very poor	Fair	Very poor	Very poor	Poor	Poor
Ha: HARNEY-----	Good	Good	Good	Poor	Poor	Good	Poor	Fair	Good	---	Poor	Good
HB: HARNEY-----	Good	Good	Good	Poor	Poor	Good	Poor	Fair	Good	---	Poor	Good

WILDLIFE INTERPRETATIONS--Continued  
Gray County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
INL: AQUOLLS-----	---	---	---	---	---	---	---	---	---	---	---	---
La: LAS ANIMAS-----	Poor	Poor	Fair	Fair	Fair	Fair	Fair	Fair	Poor	Fair	Fair	Fair
LESHO-----	Fair	Fair	Good	---	---	Fair	Fair	Fair	Fair	---	Fair	Fair
Lc: LAS ANIMAS-----	Fair	Good	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair	Good
Ld: LAS ANIMAS-----	Fair	Good	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair	Good
Le: LAS-----	Fair	Fair	Fair	---	---	Fair	Fair	Fair	Fair	---	Fair	Fair
Lh: LESHO-----	Fair	Fair	Fair	---	---	Fair	Fair	Fair	Fair	---	Fair	Fair
Lk: LESHO-----	Fair	Fair	Fair	---	---	Fair	Fair	Fair	Fair	---	Fair	Fair
SWEETWATER-----	Poor	Fair	Good	---	Very poor	Fair	Good	Good	Fair	---	Good	Fair
Ll: LINCOLN-----	Fair	Fair	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Lo: LOFTON-----	---	---	---	---	---	---	---	---	---	---	---	---
Lu: LUBBOCK-----	Good	Good	Fair	---	---	Poor	Poor	Fair	Fair	---	Poor	Fair
Md: PENDEN-----	Fair	Good	Fair	---	---	Poor	Very poor	Poor	Fair	---	Very poor	Fair
Me: PENDEN-----	Poor	Fair	Fair	---	---	Poor	Very poor	Poor	Fair	---	Very poor	Fair
Mf: PENDEN-----	Poor	Fair	Fair	---	---	Poor	Very poor	Poor	Fair	---	Very poor	Fair
Mh: PENDEN-----	Poor	Fair	Fair	---	---	Poor	Very poor	Poor	Fair	---	Very poor	Fair
Mm: MANTER-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Mn: MANTER-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Mo: MANTER-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Mp: CAMPUS-----	Fair	Good	Good	---	---	Poor	Very poor	Very poor	Fair	---	Very poor	Fair
CANLON-----	Poor	Poor	Poor	---	---	Poor	Very poor	Very poor	Poor	---	Very poor	Poor
Mu: MANTER-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
ULYSSES-----	Fair	Good	Fair	---	---	Poor	Poor	Fair	Fair	---	Poor	Fair
Pa: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Pg: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair

WILDLIFE INTERPRETATIONS--Continued  
Gray County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
Pt: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
OPTIMA-----	Fair	Good	Fair	Poor	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor	Fair
Ra: NESS-----	Poor	Poor	Poor	---	---	Poor	Fair	Good	Poor	---	Good	Poor
Rm: RICHFIELD-----	Good	Good	Fair	---	---	Poor	Very poor	Very poor	Fair	---	Very poor	Fair
Rn: RICHFIELD-----	Good	Good	Fair	---	---	Poor	Very poor	Very poor	Fair	---	Very poor	Fair
Ro: RICHFIELD-----	Good	Good	Fair	---	---	Poor	Very poor	Very poor	Fair	---	Very poor	Fair
Rs: RICHFIELD-----	Good	Good	Fair	---	---	Poor	Very poor	Very poor	Fair	---	Very poor	Fair
SPEARVILLE-----	Good	Good	Fair	---	---	Poor	Poor	Good	Fair	---	Fair	Fair
Sa: SATANTA-----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
SAP: SAND PIT-----	---	---	---	---	---	---	---	---	---	---	---	---
Sb: SATANTA-----	Good	Good	Good	---	---	Good	Poor	Very poor	Good	---	Very poor	Good
Sp: SPEARVILLE-----	Good	Good	Fair	---	---	Poor	Poor	Good	Fair	---	Fair	Fair
Sr: SPEARVILLE-----	Fair	Good	Fair	---	---	Poor	Poor	Fair	Fair	---	Poor	Fair
Sw: SWEETWATER-----	Poor	Fair	Good	---	Very poor	Fair	Good	Good	Fair	---	Good	Fair
Tf: OPTIMA-----	Fair	Good	Fair	Poor	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor	Fair
Ua: ULYSSES-----	Good	Good	Fair	---	---	Poor	Poor	Fair	Fair	---	Poor	Fair
Ub: ULYSSES-----	Good	Good	Fair	---	---	Poor	Poor	Fair	Fair	---	Poor	Fair
UBB: ULY-----	Good	Good	Good	Good	Fair	Fair	Very poor	Very poor	Good	Good	Very poor	Good
Uc: ULYSSES-----	Fair	Good	Fair	---	---	Poor	Poor	Poor	Fair	---	Poor	Fair
UCC: ULY-----	Fair	Good	Good	Good	Fair	Fair	Very poor	Very poor	Fair	Good	Very poor	Good
Um: ULYSSES-----	Fair	Good	Fair	---	---	Poor	Poor	Poor	Fair	---	Poor	Fair
COLBY-----	Fair	Good	Fair	---	---	Poor	Poor	Poor	Fair	---	Poor	Poor
W: WATER-----	---	---	---	---	---	---	---	---	---	---	---	---

YIELDS PER ACRE OF PASTURE AND HAYLAND  
Gray County, Kansas

Use and Explanation of Pastureland and Hayland Interpretations

This subsection provides information concerning the suitability of soils for the production of pasture and hayland. This subsection may contain pasture and hayland suitability groupings, land capability and yield estimates, yield estimates for individual grasses or legumes, or other information pertaining to the production of forage.

Pasture and Hayland Suitability Groupings

Soils are placed in pasture and hayland groups according to their suitability for the production of forage. The soils in each group are enough alike to be suited to the same grasses or legumes, to have similar limitations and hazards, to require similar management, and to have similar productivity and other responses to management. Thus, the pasture and hayland suitability group is a convenient way of grouping the soils for their management. If used, these groupings are identified and described in other reports in the subsection.

Yield Estimates

The average yields per acre that can be expected of the principal pasture or hayland crops, under a high level of management, are presented in this subsection. In any given year, yields may be higher or lower than those indicated in the tables because of variations in rainfall or other climatic factors. The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations are also considered.

Under good management, proper grazing is essential for the production of high quality forage, stand survival, and erosion control. Proper grazing helps plants maintain sufficient and generally vigorous top growth during the growing season. Brush control is essential in many areas, and weed control generally is needed. Rotation grazing and renovation are also important management practices.

The Pasture and Hayland table show yield estimates in tons per acre and animal unit months for pasture and hayland groups. An animal unit month is the amount of forage required by one animal unit (AU) for 30 days. On animal unit (AU) is one (1000 pound) mature cow and a calf up to weaning age (usually six months of age) or their equivalent. The Natural Resources Conservation Service uses 900 pounds of air dry forage as the amount needed to meet this requirement. To maintain a healthy and vigorous plant community, the degree of use should never be greater than 50 percent. Therefore only 25 percent of the total biomass grown is considered consumed by the grazing animal. Animal Unit Months can be converted to air dry pounds per acre production by multiplying the AUM by 30 days, then by 30 pounds per day, and then by four. This figure is the amount of total forage production.

Planners of management systems for individual fields or farms should consider the detailed information given in the description of each soil in the Nontechnical Description section. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.



YIELDS PER ACRE OF PASTURE AND HAYLAND--Continued  
Gray County, Kansas

PAGE 2 OF 4

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)  
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
055BP: Bridgeport, rarely floode	2c	1	3.00	6.00
055LD: Las, occasionally flooded	3w	2w	---	---
Las Animas, occasionally flooded-----	4w	3w	---	---
055TF: Valent-----	7e	6e	---	---
055TV: Valent-----	6e	4e	---	3.50
Vona-----	6e	4e	---	4.00
055VO: Vona-----	4e	4e	---	4.00
081LU: Lubbock-----	3c	1	---	6.50
083UM: Uly-----	4e	---	---	---
Coly-----	3e	3e	---	4.50
119HC: Harney, eroded-----	3e	---	---	---
119RX: Roxbury-----	2c	1	3.00	7.00
119UA: Uly-----	2c	1	2.50	5.50
119UT: Uly-----	4e	---	---	---
An: Bridgeport, channeled----	5w	---	---	---
ARR: Arkansas River-----	8w	---	---	---
Bc: Bridgeport-----	2c	1	3.00	6.00
Bd: Bridgeport-----	3e	2e	---	5.50
Bo: Optima-----	6e	4e	---	---
Da: Dale-----	2c	---	3.00	---
Gr: Schamber-----	7s	---	---	---
GRP: Pits-----	8s	---	---	---
Ha: Harney-----	2c	1	---	6.50
HB: Harney-----	2e	2e	---	5.50
INL: Aquolls-----	5w	---	---	---
La: Las Animas, occasionally flooded-----	4w	3w	---	---
Lesho, occasionally flooded-----	4s	3s	---	---

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)  
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Lc: Las Animas, occasionally flooded-----	6s	---	---	---
Ld: Las Animas-----	3w	3w	---	---
Le: Las, occasionally flooded	4w	2w	---	---
Lh: Lesho, occasionally flooded-----	3w	---	---	---
Lk: Lesho-----	6s	---	---	---
Sweetwater-----	5w	---	---	---
Ll: Lincoln, occasionally flooded-----	7w	---	---	---
Lo: Lofton-----	---	---	---	---
Lu: Lubbock-----	2c	1	---	6.50
Md: Penden-----	3e	---	---	---
Me: Penden-----	6e	---	---	---
Mf: Penden, Severely eroded--	6e	---	---	---
Mh: Penden, Severely Eroded--	6e	---	---	---
Mm: Manter-----	2e	2s	---	6.00
Mn: Manter-----	2e	2e	---	5.50
Mo: Manter-----	3e	3e	---	4.50
Mp: Campus-----	4e	---	---	---
Canlon-----	6e	---	---	---
Mu: Manter-----	3e	3e	---	4.50
Ulysses-----	3e	3e	---	4.00
Pa: Pratt-----	3e	3e	---	5.50
Pg: Pratt-----	4e	3e	---	5.50
Pt: Pratt-----	4e	3e	---	5.50
Optima-----	6e	4e	---	---
Ra: Ness-----	6w	---	---	---
Rm: Richfield-----	2c	1	---	6.00
Rn: Richfield-----	2e	2e	---	5.00

YIELDS PER ACRE OF PASTURE AND HAYLAND--Continued  
Gray County, Kansas

PAGE 4 OF 4

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)  
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay	
	N	I	N	I
			Tons	Tons
Ro: Richfield-----	3e	3e	---	---
Rs: Richfield-----	2c	1	---	6.00
Spearville-----	2s	2s	---	6.00
Sa: Satanta-----	2c	1	---	6.00
SAP: Sand Pit-----	---	---	---	---
Sb: Satanta-----	2e	2e	---	5.00
Sp: Spearville-----	2s	2s	---	6.00
Sr: Spearville-----	3e	---	---	---
Sw: Sweetwater-----	5w	---	---	---
Tf: Optima-----	6e	4e	---	---
Ua: Ulysses-----	2c	1	---	6.00
Ub: Ulysses-----	2e	2e	---	5.00
UBB: Uly-----	2e	2e	2.40	5.00
Uc: Ulysses-----	3e	3e	---	4.00
UCC: Uly-----	3e	---	1.90	4.50
Um: Ulysses-----	4e	---	---	---
Colby-----	4e	3e	---	3.50
W: Water-----	---	---	---	---

CONSERVATION TREE AND SHRUB MANAGEMENT  
Gray County, Kansas

A Conservation Tree/Shrub Suitability Group (CTSG), formerly Windbreak Suitability Group, is a physiographic unit or area having similar climatic and edaphic characteristics that control the selection and height growth of trees and shrubs.

In this table, the Conservation Tree and Shrub Grouping is expressed as a group index number. The group index for Conservation Tree and Shrub groups (CTSG) are a guide for species best suited for different kinds of soil and for prediction height, growth, and effectiveness. The groupings can be used when selection woody plants for windbreaks, wildlife plantings riparian buffers, reforestation, other environmental plantings, recreation, landscaping, wetland restoration or enhancement and critical area plantings. CTSG's are developed to assure satisfactory species selection and adaptation to specific conditions of soil, climate and physiography. CTSG's are a guide for selection species best suited for different kinds of soil and prediction height growth and effectiveness.

All soil series mapped in the state have been placed in 10 groups of similar soil characteristics. Groups 1, 2, 3, 4, 6, and 9 are further divided into subgroups. In addition, all groups provide information by Major Land Resource Areas.

Each tree or shrub species has certain climatic and physiographic limits. Within these parameters a tree or shrub may be well or poorly suited because of soil characteristics. Each tree or shrub also has definable potentials of height growth depending on the factors just mentioned. Accurate definitions of potential heights are necessary for proper windbreak planning and design.

Windbreaks protect livestock, buildings, roads and yards from wind and snow. They also protect fruit trees and gardens, and they furnish habitat for wildlife. Several rows of low-growing and high-growing broadleaf and coniferous trees and shrubs provide the most protection.

Field windbreaks are narrow plantings made at right angles to the prevailing wind and at specific intervals across the field. The interval depends on the erodibility of the soil. Field windbreaks protect cropland and crops from wind, help to keep snow on the fields, and provide food and cover for wildlife.

Environmental plantings help to beautify and screen houses and other buildings and to abate noise. The plants, mostly evergreen shrubs and trees, are closely spaced. To ensure plant survival, a healthy planting stock of suitable species should be planted properly on a well prepared site and maintained in good condition.

Windbreaks are often planted on land that did not grow trees originally. Knowledge of how trees perform on such land can be gained only by observing and recording their performance where trees have been planted and survived. The problem is compounded by the fact that many favorite windbreak species are not indigenous to the areas in which they are planted.

The Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups shows the adapted species listing for each group index number. Showing the height that locally grown trees and shrubs are expected to reach in 20 years on various soils. The estimates are based on measurements and observation of established plantings that have been given adequate care. This information should be used to determine the placement of a windbreak, the area protected and the arrangement of species.

A number of attributes are included in the CTSG species tables for each group number found in this section of the Field Office Technical Guide. These attributes were rated subjectively and assigned a relative value to further assist those unfamiliar with individual species characteristics or desirability for the intended use. Definitions and explanations can be found. Additional information on planning windbreaks and screens and planting and caring for trees and shrubs can be obtained from the local office of the Natural Resources Conservation Service or of the Cooperative Extension Service or from a commercial nursery. See part 537 of the National Forestry Manual for additional information.

In the Tree and Shrub Management table interpretive ratings are given for various aspects of forest and conservation tree and shrub management. Some rating class terms indicate the degree to which the soils are suited to a specified forest management practice. Well suited indicates that the soil has features that are favorable for the specified practice and has no limitations. Good performance can be expected, and little or no maintenance is needed. Moderately well suited indicates that the soil has features that are moderately favorable for the specified practice. One or more soil properties are less than desirable and fair performance can be expected. Some maintenance is needed. Poorly suited indicates that the soil has one or more properties that are unfavorable for the specified practice. Overcoming the unfavorable properties requires special design, extra maintenance, and costly alteration. Unsuitable indicates that the expected performance of the soil is unacceptable for the specified practice or that extreme measures are needed to overcome the undesirable soil properties.

The paragraphs that follow indicate the soil properties considered in rating the soils for forest and conservation tree and shrub management practices. More detailed information about the criteria used in the ratings is available in the "National Forestry Manual," which is available in local offices of the Natural Resources Conservation Service or on the Internet. Also, in the Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups.

Ratings in the columns suitability for hand planting and suitability for mechanical planting are based on slope, depth to a restrictive layer, content of sand, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, moderately well suited, poorly suited, or unsuited to these methods of planting. It is assumed that necessary site preparation is completed before seedlings are planted.

Ratings in the column suitability for mechanical site preparation (surface) are based on slope, depth to a restrictive layer, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsuited to this management activity. The part of the soil from the surface to a depth of about 1-foot is considered in the ratings.

Ratings in the column suitability for mechanical site preparation (deep) are based on slope, depth to a restrictive layer, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsuited to this management activity. The part of the soil from the surface to a depth of about 3 feet is considered in the ratings.

Ratings in the column potential for seedling mortality are based on flooding, ponding, depth to a water table, content of lime, reaction, salinity, available water capacity, soil moisture regime, soil temperature regime, aspect, and slope. The soils are described as having a low, moderate, or high potential for seedling mortality. See the National Forestry Manual, Subpart B for criteria used in rating management concerns. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

CONSERVATION TREE AND SHRUB MANAGEMENT  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
055BP: Bridgeport, rarely flooded-----	1K	Well suited	Well suited	Well suited	Well suited	Low
055LD: Las, occasionally flooded-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate
Las Animas, occasionally flooded-----	2K	Well suited	Well suited	Well suited	Well suited	Soil reaction Moderate
055TF: Valent-----		Moderately suited Sandiness	Poorly suited Slope Sandiness	Poorly suited Slope	Poorly suited Slope	Soil reaction Low
055TV: Valent-----		Well suited	Moderately suited Slope	Well suited	Well suited	Low
Vona-----	5	Well suited	Moderately suited Slope	Well suited	Well suited	Low
055VO: Vona-----	5	Well suited	Well suited	Well suited	Well suited	Low
081LU: Lubbock-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
083UM: Uly-----	3	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Coly-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Soil reaction
119HC: Harney, eroded-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
119RX: Roxbury-----	1	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
119UA: Uly-----	3	Well suited	Well suited	Well suited	Well suited	Low
119UT: Uly-----	3	Well suited	Moderately suited Slope	Well suited	Well suited	Low
An: Bridgeport, channeled-----	1K	Well suited	Well suited	Well suited	Well suited	Low
ARR: Arkansas River-----		Not rated	Not rated	Not rated	Not rated	Not rated
Bc: Bridgeport-----	1K	Well suited	Well suited	Well suited	Well suited	Low
Bd: Bridgeport-----	1K	Well suited	Well suited	Well suited	Well suited	Low
Bo: Optima-----	7	Well suited	Well suited	Well suited	Well suited	Low
Da: Dale-----	1	Well suited	Well suited	Well suited	Well suited	Low
Gr: Schamber-----	10	Not rated	Not rated	Not rated	Not rated	Not rated
GRP: Pits-----		Not rated	Not rated	Not rated	Not rated	Not rated
Ha: Harney-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
HB: Harney-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
INL: Aquolls-----		Well suited	Well suited	Well suited	Well suited	High Wetness Soil reaction
La: Las Animas, occasionally flooded-----	2K	Well suited	Well suited	Well suited	Well suited	Moderate
Lesho, occasionally flooded-----	1K	Well suited	Well suited	Well suited	Well suited	Soil reaction Salinity Moderate
Lc: Las Animas, occasionally flooded-----	2K	Well suited	Well suited	Well suited	Well suited	Soil reaction Salinity Moderate
Ld: Las Animas-----	2K	Well suited	Well suited	Well suited	Well suited	Moderate
Le: Las, occasionally flooded-----	1K	Well suited	Well suited	Well suited	Well suited	Soil reaction
Lh: Lesho, occasionally flooded-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate
Lk: Lesho-----	1K	Well suited	Well suited	Well suited	Well suited	Soil reaction
Sweetwater-----	2K	Well suited	Well suited	Well suited	Unsuited Wetness	Moderate Soil reaction Moderate Soil reaction
Ll: Lincoln, occasionally flooded-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate
Lo: Lofton-----		Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Soil reaction Low
Lu: Lubbock-----	3	Well suited	Well suited	Well suited	Well suited	Low
Md: Penden-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Moderate
Me: Penden-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Soil reaction Moderate
Mf: Penden, Severely eroded-----	8	Well suited	Well suited	Well suited	Well suited	Soil reaction Moderate
Mh: Penden, Severely Eroded-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Lime Soil reaction Moderate
Mn: Manter-----	5	Well suited	Well suited	Well suited	Well suited	Lime Soil reaction Low
Mn: Manter-----	5	Well suited	Well suited	Well suited	Well suited	Low
Mo: Manter-----	5	Well suited	Well suited	Well suited	Well suited	Low
Mp: Campus-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Moderate
Canlon-----	10	Unsuited Restrictive layer Rock fragments	Unsuited Restrictive layer Rock fragments Slope	Unsuited Restrictive layer Rock fragments	Unsuited Restrictive layer	Lime Soil reaction Moderate Soil reaction

CONSERVATION TREE AND SHRUB MANAGEMENT  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Mu:						
Manter-----	5	Well suited	Well suited	Well suited	Well suited	Low
Ulysses-----	3	Well suited	Well suited	Well suited	Well suited	Low
Pa:						
Pratt-----	7	Well suited	Well suited	Well suited	Well suited	Low
Pg:						
Pratt-----	7	Well suited	Well suited	Well suited	Well suited	Low
Pt:						
Pratt-----	7	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Optima-----	7	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Ra:						
Ness-----	10	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	High Wetness
Rm:						
Richfield-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Rn:						
Richfield-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Ro:						
Richfield-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Rs:						
Richfield-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Spearville-----	4C	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
Sa:						
Satanta-----	3	Well suited	Well suited	Well suited	Well suited	Low
SAP:						
Sand Pit-----		Not rated	Not rated	Not rated	Not rated	Not rated
Sb:						
Satanta-----	3	Well suited	Well suited	Well suited	Well suited	Low
Sp:						
Spearville-----	4C	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
Sr:						
Spearville-----	4C	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
Sw:						
Sweetwater-----	2K	Well suited	Well suited	Well suited	Unsuited Wetness	Moderate Soil reaction
Tf:						
Optima-----	7	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Ua:						
Ulysses-----	3	Well suited	Well suited	Well suited	Well suited	Low
Ub:						
Ulysses-----	3	Well suited	Well suited	Well suited	Well suited	Low
UBB:						
Uly-----	3	Well suited	Well suited	Well suited	Well suited	Low
Uc:						
Ulysses-----	3	Well suited	Moderately suited Slope	Well suited	Well suited	Low
UCC:						
Uly-----	3	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Um:						
Ulysses-----	3	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Colby-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Soil reaction
W:						
Water-----		Not rated	Not rated	Not rated	Not rated	Not rated

ENGINEERING INDEX PROPERTIES  
Gray County, Kansas

Engineering Index Properties table gives the engineering classifications and the range of index properties for the layers of each soil in the survey area. Depth to the upper and lower boundaries of each layer is indicated. Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter. Loam, for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, gravelly. Textural terms are defined in the Glossary.

Classification of the soils is determined according to the Unified soil classification system (ASTM, 1998) and the system adopted by the American Association of State Highway and Transportation Officials (AASHTO, 1998). The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection. If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional refinement, the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest. The AASHTO classification for soils tested, with group index numbers in parentheses, is given in Engineering Index Properties table.

Rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in the field to weight percentage. Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field.

Liquid limit and plasticity index (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination. The estimates of particle-size distribution, liquid limit, and plasticity index are generally rounded to the nearest 5 percent. Thus, if the ranges of gradation and Atterberg limits extend a marginal amount (1 or 2 percentage points) across classification boundaries, the classification in the marginal zone is generally omitted in the table.



ENGINEERING INDEX PROPERTIES--Continued  
Gray County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
055BP: Bridgeport, rarely flooded	0-11	Clay loam	CL	A-6	0	0	100	100	95-100	75-95	30-40	11-20
	11-60	Loam	CL	A-4, A-6	0	0	100	100	90-100	65-100	25-40	8-20
055LD: Las, occasionally flooded-----	0-8	Loam	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	50-85	20-40	5-20
	8-31	Clay loam	CL, CL-ML	A-4, A-6, A-7	0	0	100	95-100	95-100	50-85	22-45	5-25
	31-60	Coarse sand	SM, SP-SM	A-1, A-2, A-3	0	0	100	95-100	30-70	5-35	---	NP
Las Animas, occasionally flooded-----	0-14	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-4	0	0	100	95-100	70-90	40-60	20-25	NP-5
	14-30	Stratified very fine sandy loam to loamy fine sand	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	95-100	90-100	55-90	25-55	20-25	NP-5
	30-60	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	75-100	5-25	---	NP
055TF: Valent-----	0-4	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	60-70	5-25	---	NP
	4-50	Fine sand	SM	A-2	0	0	100	95-100	75-90	10-30	---	NP
055TV: Valent-----	0-4	Loamy fine sand	SM, SP-SM	A-2	0	0	100	100	70-95	10-30	---	NP
	4-60	Fine sand	SM	A-2	0	0	100	95-100	75-90	10-30	---	NP
Vona-----	0-8	Loamy fine sand	SM	A-2	0	0	100	90-100	60-90	15-30	---	NP
	8-22	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	90-100	60-90	30-45	20-30	NP-10
	22-60	Fine sandy loam	SM	A-2	0	0	100	90-100	50-85	15-30	15-25	NP-5
055VO: Vona-----	0-8	Loamy fine sand	SM	A-2	0	0	100	90-100	60-90	15-30	---	NP
	8-22	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	90-100	60-90	30-45	20-30	NP-10
	22-60	Sandy loam	SM	A-2	0	0	100	90-100	50-85	15-30	15-25	NP-5
081LU: Lubbock-----	0-9	Silty clay loam	CL	A-6	0	0	100	100	95-100	65-100	30-40	10-20
	9-42	Silty clay loam	CH, CL	A-7	0	0	100	100	95-100	85-100	40-60	20-35
	42-72	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	85-100	30-40	7-17
083UM: Uly-----	0-6	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	100	95-100	20-40	2-20
	6-19	Silty clay loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-40	3-15
	19-60	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-40	3-15
Coly-----	0-6	Silt loam	CL, CL-ML, ML	A-4, A-6, A-7	0	0	100	100	85-100	85-100	20-45	2-20
	6-60	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	85-100	85-100	20-40	2-15
119HC: Harney, eroded-	0-6	Silty clay loam	CL	A-6, A-7-6	0	0	100	100	95-100	85-100	30-45	15-22
	6-20	Silty clay loam	CH, CL	A-7-6	0	0	100	100	95-100	85-100	40-60	15-35
	20-60	Silt loam	CL	A-6, A-7-6	0	0	100	100	95-100	85-100	30-45	10-20
119RX: Roxbury-----	0-20	Silt loam	CL	A-4, A-6	0	0	100	100	90-100	70-90	30-35	10-15
	20-60	Silty clay loam	CL	A-4, A-6, A-7-6	0	0	100	100	95-100	85-100	30-45	10-20
119UA: Uly-----	0-6	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
	6-12	Silt loam	CL	A-6, A-7	0	0	100	100	90-100	85-100	25-43	11-20
	12-18	Silt loam	CL	A-6, A-7	0	0	100	100	90-100	85-100	25-43	11-20
	18-32	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
	32-62	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
119UT: Uly-----	0-6	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	100	95-100	20-40	2-20
	6-14	Silty clay loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-40	3-15
	14-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	100	95-100	25-40	3-15
An: Bridgeport, channeled----	0-12	Loam	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	65-90	20-35	4-19
	12-60	Loam	CL	A-4, A-6	0	0	100	100	90-100	65-100	25-40	8-20
ARR: Arkansas River-	0-6	Sand	SP, SP-SM, SW, SW-SM	A-1, A-3	---	0-5	80-100	75-100	30-60	0-10	---	NP
	6-60	Stratified coarse sand to sandy loam	SM, SP, SP-SM	A-1, A-2, A-3	---	0-5	80-100	75-100	40-70	0-20	---	NP
Bc: Bridgeport-----	0-10	Silty clay loam	CL	A-6	0	0	100	100	95-100	75-95	30-40	11-20
	10-65	Silty clay loam	CL	A-4, A-6	0	0	100	100	90-100	65-100	25-40	8-20
Bd: Bridgeport-----	0-10	Silty clay loam	CL	A-6	0	0	100	100	95-100	75-95	30-40	11-20
	10-65	Silty clay loam	CL	A-4, A-6	0	0	100	100	90-100	65-100	25-40	8-20
Bo: Optima-----	0-8	Loamy fine sand	ML, SM, SP, SP-SM	A-2, A-3, A-4	0	0	100	100	75-100	4-55	13-21	NP-4
	8-17	Fine sand	ML, SM, SP-SM	A-2, A-4	0	0	100	100	75-100	10-55	13-21	NP-4
Da: Dale-----	17-80	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	50-80	4-35	13-16	NP
	0-25	Silt loam	CL, CL-ML	A-4, A-6	0	0	95-100	95-100	90-100	65-98	25-35	5-15
	25-60	Silt loam	CL	A-4, A-6, A-7	0	0	95-100	95-100	90-100	65-98	30-43	8-20
Gr: Schamber-----	0-6	Gravelly sandy loam	GM, GW-GM, SM, SW-SM	A-1, A-2	---	0-5	55-90	50-75	40-60	10-35	15-25	NP-5
	6-60	Very gravelly loamy sand	GW, GW-GM, SW, SW-SM	A-1	---	0-15	30-80	25-50	5-20	0-10	15-25	NP-5

ENGINEERING INDEX PROPERTIES--Continued  
Gray County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
GRP: Pits-----	0-60	Gravelly sand	GP-GM, SM, SP, SP-SM	A-1, A-2, A-3	---	0-5	45-100	40-100	0-80	0-40	0-14	NP
Ha: Harney-----	0-9 9-25 25-65	Silt loam Silty clay loam Silty clay loam	CL, CL-ML CH, CL CL	A-4, A-6 A-7-6 A-6, A-7-6	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	85-100 85-100 85-100	25-40 40-60 30-45	5-20 15-35 10-20
HB: Harney-----	0-6 6-20 20-60	Silt loam Silty clay loam Silt loam	CL, CL-ML CH, CL CL	A-4, A-6 A-7-6 A-6, A-7-6	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	85-100 85-100 85-100	25-40 40-60 30-45	5-20 15-35 10-20
INL: Aquolls-----	0-72	Variable			---	---	---	---	---	---	---	---
La: Las Animas, occasionally flooded-----	0-11 11-60	Sandy loam Sand	CL-ML, ML, SC-SM, SM CL-ML, ML, SC-SM, SM	A-4 A-2, A-4	0 0	0 0	100 95-100	100 90-100	70-85 55-90	40-55 25-55	20-25 20-25	NP-5 NP-5
Lesho, occasionally flooded-----	0-14 14-25 25-60	Clay loam Clay loam Coarse sand	CL CL SM, SP-SM	A-6, A-7-6 A-2, A-3	0 0 0	0 0 0	100 100 100	100 95-100	85-100 55-75	65-95 5-35	30-45 ---	11-22 NP
Lc: Las Animas, occasionally flooded-----	0-11 11-30 30-60	Loamy sand Stratified very fine sandy loam to loamy fine sand Fine sand	SM CL-ML, ML, SC-SM, SM SM, SP-SM	A-2, A-4 A-2, A-4 A-2, A-3	0 0 0	0 0 0	95-100 95-100 100	90-100 90-100 95-100	60-75 55-90 75-100	20-40 25-55 5-25	---	NP NP-5 NP
Ld: Las Animas-----	0-11 11-30 30-60	Sandy loam Stratified very fine sandy loam to loamy fine sand Fine sand	CL-ML, ML, SC-SM, SM CL-ML, ML, SC-SM, SM SM, SP-SM	A-4 A-2, A-4 A-2, A-3	0 0 0	0 0 0	100 95-100 100	95-100 90-100 95-100	70-90 55-90 75-100	40-60 25-55 5-25	20-25 20-25 ---	NP-5 NP-5 NP
Le: Las, occasionally flooded-----	0-8 8-31 31-60	Clay loam Clay loam Coarse sand	CL CL, CL-ML SM, SP-SM	A-6, A-7 A-4, A-6, A-7 A-1, A-2, A-3	0 0 0	0 0 0	100 100 100	100 95-100 95-100	90-100 95-100 30-70	50-85 5-35	30-45 22-45 ---	10-25 5-25 NP
Lh: Lesho, occasionally flooded-----	0-14 14-25 25-60	Clay loam Clay loam Coarse sand	CL CL SM, SP-SM	A-6, A-7-6 A-4, A-6, A- 7-6 A-1, A-2, A- 3, A-4	0 0 0	0 0 0	100 100 100	100 85-100 95-100	95-100 65-95 30-85	65-85 25-45 5-45	35-45 25-45 ---	15-22 7-22 NP
Lk: Lesho-----	0-14 14-25 25-60	Loam Clay loam Coarse sand	CL CL SM, SP-SM	A-4, A-6 A-4, A-6, A- 7-6 A-2, A-3, A- 4, A-1	0 0 0	0 0 0	100 100 100	100 100 95-100	85-100 85-100	60-85 65-95	25-40 25-45	7-20 7-22
Sweetwater-----	0-17 17-60	Clay loam Coarse sand	CL, CL-ML, SC SM	A-4, A-6 A-2	0 0	0 0	100 95-100	95-100 90-100	80-95 50-80	40-70 15-35	25-40 15-22	7-20 NP-2
Ll: Lincoln, occasionally flooded-----	0-8 8-60	Fine sandy loam Stratified fine sand to clay loam	CL-ML, ML, SC-SM, SM SM, SP-SM	A-4 A-2, A-3	0 0	0 0	100 100	98-100 98-100	94-100 82-100	36-60 5-35	15-24 ---	NP-7 NP
Lo: Lofton-----	0-6 6-36 36-60	Silty clay loam Silty clay Silty clay loam	CL CH, CL CL, ML	A-6, A-7 A-7 A-4, A-6	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	95-100 95-100 80-100	35-50 40-65 25-40	15-25 20-45 NP-15
Lu: Lubbock-----	0-15 15-36 36-60	Loam Silty clay loam Clay loam	CL, CL-ML CH, CL CL, ML	A-4, A-6 A-7 A-4, A-6	0 0 0	0 0 0	100 100 100	100 100 100	95-100 95-100 95-100	65-100 85-100 85-100	25-40 40-60 30-45	5-20 20-35 7-17
Md: Penden-----	0-14 14-32 32-60	Clay loam Clay loam Clay loam	CL CL CL	A-6, A-7-6 A-6, A-7-6 A-6, A-7-6	0 0 0	0 0 0	100 100 100	100 100 100	85-100 85-100 75-100	65-95 60-90 55-75	30-45 30-45 30-45	11-25 11-25 11-25
Me: Penden-----	0-15 15-34 34-60	Clay loam Clay loam Clay loam	CL CL CL	A-6, A-7-6 A-6, A-7-6 A-6, A-7-6	0 0 0	0 0 0	100 100 100	100 100 100	85-100 85-100 75-100	65-95 60-90 55-75	30-45 30-45 30-45	11-25 11-25 11-25

ENGINEERING INDEX PROPERTIES--Continued  
Gray County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct						
Mf: Penden, Severel eroded-----	In											
	0-4	Clay loam	CL	A-6, A-7-6	0	0	100	100	85-100	65-95	30-45	11-25
	4-22	Clay loam	CL	A-6, A-7-6	0	0	100	100	85-100	60-90	30-45	11-25
	22-30	Clay loam	CL	A-6, A-7-6	0	0	100	100	75-100	55-75	30-45	11-25
Mh: Penden, Severel Eroded-----	30-80	Clay loam			0	0	100	100	75-100	55-75	30-45	11-25
	0-4	Clay loam	CL	A-6, A-7-6	0	0	100	100	85-100	65-95	30-45	11-25
	4-22	Clay loam	CL	A-6, A-7-6	0	0	100	100	85-100	60-90	30-45	11-25
	22-30	Clay loam	CL	A-6, A-7-6	0	0	100	100	75-100	55-75	30-45	11-25
Mm: Manter-----	30-80	Clay loam			0	0	100	100	75-100	55-75	30-45	11-25
	0-18	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	95-100	75-100	45-85	25-55	20-30	NP-10
	18-30	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	95-100	75-100	50-85	30-55	15-25	NP-10
	30-62	Loam	SM	A-1, A-2, A-4	0	0	95-100	75-100	40-85	15-50	---	NP
Mn: Manter-----	0-18	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	95-100	75-100	45-85	25-55	20-30	NP-10
	18-30	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	95-100	75-100	50-85	30-55	15-25	NP-10
	30-62	Sandy loam	SM	A-1, A-2, A-4	0	0	95-100	75-100	40-85	15-50	---	NP
	0-18	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	95-100	75-100	45-85	25-55	20-30	NP-10
Mo: Manter-----	18-30	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	95-100	75-100	50-85	30-55	15-25	NP-10
	30-62	Sandy loam	SM	A-1, A-2, A-4	0	0	95-100	75-100	40-85	15-50	---	NP
	0-9	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	80-100	55-90	20-40	3-20
	9-19	Loam	CL, ML	A-4, A-6, A-7	0	0	100	100	75-95	50-80	33-45	8-20
Canlon-----	>19	Unweathered bedrock			---	---	---	---	---	---	---	---
	0-8	Silt loam	CL, CL-ML	A-4, A-6	0	0	90-100	75-100	65-100	50-90	20-40	4-20
	>8	Unweathered bedrock			---	---	---	---	---	---	---	---
	0-18	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	95-100	75-100	45-85	25-55	20-30	NP-10
Ulysses-----	18-30	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	95-100	75-100	50-85	30-55	15-25	NP-10
	30-62	Sandy loam	SM	A-1, A-2, A-4	0	0	95-100	75-100	40-85	15-50	---	NP
	0-5	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
	5-14	Silty clay loam	CL	A-6, A-7	0	0	100	100	90-100	85-100	25-43	11-20
Pa: Pratt-----	14-60	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
	0-11	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	---	NP
	11-26	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	26-60	Loamy fine sand	SP-SM, SM	A-2, A-3	0	0	100	95-100	80-100	5-35	---	NP
Pg: Pratt-----	0-11	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	---	NP
	11-40	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	40-60	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-35	---	NP
Pt: Pratt-----	0-11	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	---	NP
	11-26	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	26-60	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-35	---	NP
	Optima-----	0-8	Loamy fine sand	ML, SM, SP, SP-SM	A-2, A-3, A-4	0	0	100	100	75-100	4-55	13-21
Ra: Ness-----	8-17	Fine sand	ML, SM, SP-SM	A-2, A-4	0	0	100	100	75-100	10-55	13-21	NP-4
	17-80	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	50-80	4-35	13-16	NP
	0-31	Clay	CH	A-7-6	0	0	100	100	95-100	90-100	50-70	30-45
	31-60	Silt loam	CH, CL, MH, ML	A-4, A-6, A-7-6	0	0	100	100	95-100	90-100	30-55	8-30
Rm: Richfield-----	0-5	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	70-100	20-35	2-15
	5-17	Silty clay loam	CH, CL	A-7-6	0	0	100	100	95-100	90-100	40-60	20-35
	17-60	Silty clay loam	CL, CL-ML	A-4, A-6, A-7-6	0	0	100	100	95-100	75-100	25-45	5-20
	0-6	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	70-100	20-35	2-15
Rn: Richfield-----	6-19	Silty clay loam	CH, CL	A-7-6	0	0	100	100	95-100	90-100	40-60	20-35
	19-60	Silt loam	CL, CL-ML	A-4, A-6, A-7-6	0	0	100	100	95-100	75-100	25-45	5-20
	0-6	Silty clay loam	CL	A-6	0	0	100	100	90-100	75-100	30-40	10-20
	6-19	Silty clay loam	CH, CL	A-7-6	0	0	100	100	95-100	90-100	40-60	20-35
Ro: Richfield-----	19-60	Silt loam	CL, CL-ML	A-4, A-6, A-7-6	0	0	100	100	95-100	75-100	25-45	5-20

ENGINEERING INDEX PROPERTIES--Continued  
Gray County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
Rs:												
Richfield-----	0-6	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	70-100	20-35	2-15
	6-19	Silty clay loam	CH, CL	A-7-6	0	0	100	100	95-100	90-100	40-60	20-35
	19-60	Silt loam	CL, CL-ML	A-4, A-6, A-7-6	0	0	100	100	95-100	75-100	25-45	5-20
Spearville-----	0-7	Silty clay loam	CL	A-6	0	0	100	100	95-100	90-100	25-40	11-20
	7-20	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-100	50-70	25-45
	20-26	Silty clay loam	CH, CL	A-6, A-7-6	0	0	100	100	95-100	90-100	35-55	15-30
	26-60	Silt loam	CL	A-6, A-7-6	0	0	100	100	95-100	90-100	30-45	11-20
Sa:												
Satanta-----	0-10	Loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	80-100	60-75	20-35	5-15
	10-32	Clay loam	CL, SC	A-6, A-7	0	0	100	95-100	75-100	40-80	30-45	11-20
	32-50	Silt loam	CL, SC	A-4, A-6	0	0	100	95-100	65-100	40-80	20-35	5-15
SAP:												
Sand Pit-----	---	---	---	---	---	---	---	---	---	---	---	---
Sb:												
Satanta-----	0-10	Loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	80-100	60-75	20-35	5-15
	10-32	Clay loam	CL, SC	A-6, A-7	0	0	100	95-100	75-100	40-80	30-45	11-20
	32-50	Silt loam	CL, SC	A-4, A-6	0	0	100	95-100	65-100	40-80	20-35	5-15
Sp:												
Spearville-----	0-7	Silty clay loam	CL	A-6	0	0	100	100	95-100	90-100	25-40	11-20
	7-20	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-100	50-70	25-45
	20-26	Silty clay loam	CH, CL	A-6, A-7-6	0	0	100	100	95-100	90-100	35-55	15-30
	26-60	Silt loam	CL	A-6, A-7-6	0	0	100	100	95-100	90-100	30-45	11-20
Sr:												
Spearville-----	0-7	Silty clay loam	CL	A-6	0	0	100	100	95-100	90-100	25-40	11-20
	7-20	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-100	50-70	25-45
	20-26	Silty clay loam	CH, CL	A-6, A-7-6	0	0	100	100	95-100	90-100	35-55	15-30
	26-60	Silt loam	CL	A-6, A-7-6	0	0	100	100	95-100	90-100	30-45	11-20
Sw:												
Sweetwater-----	0-17	Clay loam	CL, CL-ML, SC	A-4, A-6	0	0	100	95-100	80-95	40-70	25-40	7-20
	17-60	Very fine sandy loam	SM	A-2	0	0	95-100	90-100	50-80	15-35	15-22	NP-2
Tf:												
Optima-----	0-8	Fine sand	ML, SM, SP, SP-SM	A-2, A-3, A-4	0	0	100	100	75-100	4-55	13-21	NP-4
	8-17	Fine sand	ML, SM, SP-SM	A-2, A-4	0	0	100	100	75-100	10-55	13-21	NP-4
	17-80	Fine sand	SM, SP, SP-SM	A-2, A-3	0	0	100	100	50-80	4-35	13-16	NP
Ua:												
Ulysses-----	0-5	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
	5-14	Silty clay loam	CL	A-6, A-7	0	0	100	100	90-100	85-100	25-43	11-20
	14-60	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
Ub:												
Ulysses-----	0-5	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
	5-14	Silty clay loam	CL	A-6, A-7	0	0	100	100	90-100	85-100	25-43	11-20
	14-60	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
UBB:												
Uly-----	0-8	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	100	95-100	20-40	2-20
	8-55	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-40	3-15
	55-60	Silt loam	CL, ML, CL-ML	A-6, A-4	0	0	100	100	100	95-100	25-40	3-15
Uc:												
Ulysses-----	0-6	Silt loam	CL, ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
	6-16	Silty clay loam	CL	A-6, A-7	0	0	100	100	90-100	85-100	25-43	11-20
	16-60	Silt loam	CL, ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
UCC:												
Uly-----	0-6	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	100	95-100	20-40	2-20
	6-14	Silty clay loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-40	3-15
	14-60	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-40	3-15
Um:												
Ulysses-----	0-5	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
	5-14	Silty clay loam	CL	A-7, A-6	0	0	100	100	90-100	85-100	25-43	11-20
	14-60	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
Colby-----	0-5	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
	5-60	Silt loam	CL, ML, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	25-40	3-15
W:												
Water-----	---	---	---	---	---	---	---	---	---	---	---	---

Physical Properties table shows estimates of some physical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Particle size is the effective diameter of a soil particle as measured by sedimentation, sieving, or micrometric methods. Particle sizes are expressed as classes with specific effective diameter class limits. The broad classes are sand, silt, and clay, ranging from the larger to the smaller.

Sand as a soil separate consists of mineral soil particles that are 0.05 millimeter to 2 millimeters in diameter. In this table, the estimated sand content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Silt as a soil separate consists of mineral soil particles that are 0.002 to 0.05 millimeter in diameter. In this table, the estimated silt content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Clay as a soil separate consists of mineral soil particles that are less than 0.002 millimeter in diameter. In this table, the estimated clay content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of sand, silt, and clay affects the physical behavior of a soil. Particle size is important for engineering and agronomic interpretations, for determination of soil hydrologic qualities, and for soil classification.

The amount and kind of clay affect the fertility and physical condition of the soil and the ability of the soil to adsorb cations and to retain moisture. They influence shrink-swell potential, permeability, plasticity, the ease of soil dispersion, and other soil properties. The amount and kind of clay in a soil also affect tillage and earth moving operations.

Moist bulk density is the weight of soil (oven-dry) per unit volume. Volume is measured when the soil is at field moisture capacity, that is, the moisture content at 1/3- or 1/10-bar (33kPa or 10kPa) moisture tension. Weight is determined after the soil is dried at 105 degrees C. In the table, the estimated moist bulk density of each soil horizon is expressed in grams per cubic centimeter of soil material that is less than 2 millimeters in diameter. Bulk density data are used to compute shrink-swell potential, available water capacity, total pore space, and other soil properties. The moist bulk density of a soil indicates the pore space available for water and roots. Depending on soil texture, a bulk density of more than 1.4 can restrict water storage and root penetration. Moist bulk density is influenced by texture, kind of clay, content of organic matter, and soil structure.

Permeability ( $K_{sat}$ ) refers to the ability of a soil to transmit water or air. The term "permeability," as used in soil surveys, indicates saturated hydraulic conductivity ( $K_{sat}$ ). The estimates in the table indicate the rate of water movement, in inches per hour, when the soil is saturated. They are based on soil characteristics observed in the field, particularly structure, porosity, and texture. Permeability is considered in the design of soil drainage systems and septic tank absorption fields.

Available water capacity refers to the quantity of water that the soil is capable of storing for use by plants. The capacity for water storage is given in inches of water per inch of soil for each soil layer. The capacity varies, depending on soil properties that affect retention of water. The most important properties are the content of organic matter, soil texture, bulk density, and soil structure. Available water capacity is an important factor in the choice of plants or crops to be grown and in the design and management of irrigation systems. Available water capacity is not an estimate of the quantity of water actually available to plants at any given time.

Linear extensibility refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. It is an expression of the volume change between the water content of the clod at 1/3- or 1/10-bar tension (33kPa or 10kPa tension) and oven dryness. The volume change is reported in the table as percent change for the whole soil. Volume change is influenced by the amount and type of clay minerals in the soil.

Linear extensibility is used to determine the shrink-swell potential of soils. The shrink-swell potential is low if the soil has a linear extensibility of less than 3 percent; moderate if 3 to 6 percent; high if 6 to 9 percent; and very high if more than 9 percent. If the linear extensibility is more than 3, shrinking and swelling can cause damage to buildings, roads, and other structures and to plant roots. Special design commonly is needed.

Organic matter is the plant and animal residue in the soil at various stages of decomposition. In Physical Properties table, the estimated content of organic matter is expressed as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter. The content of organic matter in a soil can be maintained by returning crop residue to the soil. Organic matter has a positive effect on available water capacity, water infiltration, soil organism activity, and tilth. It is a source of nitrogen and other nutrients for crops and soil organisms.

Erosion factors are shown in the Physical Properties table as the K factor ( $K_w$  and  $K_f$ ) and the T factor. Erosion factor K indicates the susceptibility of a soil to sheet and rill erosion by water. Factor K is one of six factors used in the Universal Soil Loss Equation (USLE) and the Revised Universal Soil Loss Equation (RUSLE) to predict the average annual rate of soil loss by sheet and rill erosion in tons per acre per year. The estimates are based primarily on percentage of silt, sand, and organic matter and on soil structure and permeability. Values of K range from 0.02 to 0.69. Other factors being equal, the higher the value, the more susceptible the soil is to sheet and rill erosion by water.

Erosion factor  $K_w$  indicates the erodibility of the whole soil. The estimates are modified by the presence of rock fragments.

Erosion factor  $K_f$  indicates the erodibility of the fine-earth fraction, or the material less than 2 millimeters in size.

Erosion factor T is an estimate of the maximum average annual rate of soil erosion by wind or water that can occur without affecting crop productivity over a sustained period. The rate is in tons per acre per year.

Wind erodibility groups are made up of soils that have similar properties affecting their susceptibility to

wind erosion in cultivated areas. The soils assigned to group 1 are the most susceptible to wind erosion, and those assigned to group 8 are the least susceptible. The groups are as follows:

1. Coarse sands, sands, fine sands, and very fine sands.
2. Loamy coarse sands, loamy sands, loamy fine sands, loamy very fine sands, ash material, and sapric soil material.
3. Coarse sandy loams, sandy loams, fine sandy loams, and very fine sandy loams.
- 4L. Calcareous loams, silt loams, clay loams, and silty clay loams.
4. Clays, silty clays, noncalcareous clay loams, and silty clay loams that are more than 35 percent clay.
5. Noncalcareous loams and silt loams that are less than 20 percent clay and sandy clay loams, sandy clays, and hemic soil material.
6. Noncalcareous loams and silt loams that are more than 20 percent clay and noncalcareous clay loams that are less than 35 percent clay.
7. Silts, noncalcareous silty clay loams that are less than 35 percent clay, and fibric soil material.
8. Soils that are not subject to wind erosion because of coarse fragments on the surface or because of surface wetness.

Wind erodibility index is a numerical value indicating the susceptibility of soil to wind erosion, or the tons per acre per year that can be expected to be lost to wind erosion. There is a close correlation between wind erosion and the texture of the surface layer, the size and durability of surface clods, rock fragments, organic matter, and a calcareous reaction. Soil moisture and frozen soil layers also influence wind erosion.

#### Explanation of Wind Erodibility Groups

Soil erodibility by wind is directly related to the percentage of dry non-erodible surface soil aggregates larger than 0.84 mm in diameter. From this percentage, the wind erodibility index (I-factor) is determined. The I-factor is an expression of the stability of these soil aggregates against breakdown by tillage and abrasion from wind erosion. Soils are placed in Wind Erodibility Groups (WEG) having similar percentages of dry soil aggregates larger than 0.84 mm as shown in the following table.

WEG	Properties of Soil Surface Layer	Dry Soil Aggregates >0.84mm Percent	Wind Erodibility Index T/Ac/Yr (I)
1	Very fine sand, fine sand, sand, or coarse sand	1 2 3 5 7	310 1/ 250 220 180 160
2	Loamy very fine sand, loamy fine sand, loamy sand, loamy coarse sand, organic soil materials.	10	134
3	Very fine sandy loam, fine sandy loam, sandy loam, or coarse sandy loam.	25	86
4	Clay, silty clay, non-calcareous clay loam, or silty clay loam with >35 percent clay content.	25	86
4L	Calcareous 2/ loam, silt loam, clay loam, or silty clay loam.	25	86
5	Non-calcareous loam and silt loam with <20 percent clay content, or sandy clay loam, sandy clay, and hemic 3/ organic soil materials.	40	56
6	Non-calcareous loam and silt loam with >20 percent clay content, or non-calcareous clay loam with <35 percent clay content.	45	48
7	Silt, non-calcareous silty clay loam with >35 percent clay content and fibric 3/ organic soil material.	50	38
8	Soils not suitable for cultivation due to coarse fragments or wetness; wind erosion is not a problem.	--	0

1/ The "I" values for WEG 1 vary from 160 for coarse sands to 310 for very fine sands. Use an "I" of 220 as an average figure. For coarser sand that has gravel, use a lower figure. For a soil that has no gravel and very fine sand, use a higher figure. (Modification for coarse fragments is preparation.)

2/ Calcareous is a strongly or violently effervescent reaction to cold dilute (1N) HCL.

3/ See Soil Taxonomy for definition.

PHYSICAL PROPERTIES OF THE SOILS--Continued  
Gray County, Kansas: Maintenance needed

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
055BP: Bridgeport, rarely flooded-----	0-11	27	44	27-32	1.35-1.45	0.60-2.00	0.17-0.19	3.0-5.9	1.0-3.0	.28	.28	5	4L	86
055LD: Las, occasionally flooded-----	11-60	33	43	18-30	1.35-1.45	0.60-2.00	0.17-0.22	0.0-2.9	0.5-2.0	.43	.43			
	0-8	42	37	15-27	1.35-1.45	0.60-2.00	0.20-0.22	3.0-5.9	0.5-1.0	.32	.32	4	4L	86
	8-31	35	38	18-35	1.35-1.45	0.20-0.60	0.12-0.19	3.0-5.9	0.1-0.5	.32	.32			
	31-60	91	6	1-5	1.50-1.60	6.00-20.00	0.02-0.04	0.0-2.9	0.0-0.5	.10	.10			
Las Animas, occasionally flooded-----	0-14	67	20	8-18	1.45-1.55	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.24	.24	5	3	86
	14-30	63	24	8-18	1.50-1.60	2.00-6.00	0.09-0.19	0.0-2.9	0.0-1.0	.28	.28			
	30-60	97	1	0-5	1.50-1.60	6.00-20.00	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
055TF: Valent-----	0-4	95	1	2-6	1.50-1.60	5.95-19.98	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	250
	4-50	94	1	2-8	1.55-1.65	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
055TV: Valent-----	0-4	87	7	3-10	1.50-1.60	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	4-60	94	1	2-8	1.55-1.65	5.95-19.98	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
Vona-----	0-8	78	16	3-8	1.55-1.65	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	4	2	134
	8-22	67	20	8-18	1.40-1.50	2.00-6.00	0.12-0.17	0.0-2.9	0.5-1.0	.24	.24			
	22-60	64	27	3-15	1.45-1.55	5.95-19.98	0.08-0.13	0.0-2.9	0.0-0.5	.20	.20			
055VO: Vona-----	0-8	78	16	3-8	1.55-1.65	5.95-19.98	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	4	2	134
	8-22	67	20	8-18	1.40-1.50	2.00-6.00	0.12-0.17	0.0-2.9	0.5-1.0	.24	.24			
	22-60	67	24	3-15	1.45-1.55	5.95-19.98	0.08-0.13	0.0-2.9	0.0-0.5	.20	.20			
081LU: Lubbock-----	0-9	20	49	27-35	1.40-1.50	0.60-2.00	0.21-0.23	0.0-2.9	1.0-3.0	.32	.32	5	7	38
	9-42	7	53	35-45	1.35-1.45	0.20-0.60	0.09-0.20	6.0-8.9	0.5-1.0	.32	.32			
	42-72	23	50	20-35	1.40-1.50	0.60-2.00	0.14-0.22	0.0-2.9	0.0-0.7	.32	.32			
083UM: Uly-----	0-6	11	67	17-27	1.20-1.30	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	6-19	7	68	20-30	1.20-1.30	0.60-2.00	0.18-0.22	0.0-2.9	0.5-1.0	.43	.43			
	19-60	10	68	18-27	1.10-1.20	0.60-2.00	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
Coly-----	0-6	11	68	18-24	1.35-1.45	0.60-2.00	0.22-0.24	0.0-2.9	0.5-1.0	.43	.43	5	4L	86
	6-60	11	68	18-24	1.35-1.45	0.60-2.00	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
119HC: Harney, eroded-----	0-6	20	48	28-35	1.30-1.40	0.60-2.00	0.21-0.23	3.0-5.9	0.7-2.0	.32	.32	5	7	38
	6-20	7	54	35-42	1.40-1.50	0.20-0.60	0.11-0.20	3.0-5.9	0.5-1.0	.37	.37			
	20-60	20	50	24-35	1.25-1.35	0.60-2.00	0.18-0.22	0.0-2.9	0.0-0.5	.43	.43			
119RX: Roxbury-----	0-20	10	68	18-27	1.35-1.45	0.60-2.00	0.22-0.24	0.0-2.9	1.0-3.0	.32	.32	5	4L	86
	20-60	7	66	18-35	1.40-1.50	0.60-2.00	0.17-0.22	3.0-5.9	0.5-1.0	.43	.43			
119UA: Uly-----	0-6	21-44	70	10-27	1.15-1.25	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	6-12	15-30	64	21-32	1.25-1.35	0.60-2.00	0.18-0.22	3.0-5.9	0.5-1.5	.43	.43			
	12-18	10-15	64	21-32	1.25-1.35	0.60-2.00	0.18-0.22	3.0-5.9	0.5-1.0	.43	.43			
	18-32	15-27	68	18-27	1.25-1.35	0.60-2.00	0.17-0.22	0.0-2.9	0.2-0.5	.43	.43			
	32-62	15-45	68	18-27	1.25-1.35	0.60-2.00	0.17-0.22	0.0-2.9	0.2-0.5	.43	.43			
119UT: Uly-----	0-6	11	67	17-27	1.20-1.30	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	6-14	7	68	20-30	1.20-1.30	0.60-2.00	0.21-0.24	0.0-2.9	0.5-1.0	.43	.43			
	14-60	10	68	18-27	1.10-1.20	0.60-2.00	0.17-0.22	0.0-2.9	0.2-0.5	.43	.43			
An: Bridgeport, channeled---	0-12	37	42	14-27	1.30-1.40	0.60-2.00	0.20-0.22	0.0-2.9	1.0-3.0	.28	.28	5	4L	86
	12-60	33	43	18-30	1.40-1.50	0.60-2.00	0.17-0.22	0.0-2.9	0.5-1.0	.43	.43			
ARR: Arkansas River-----	0-6	98	2	0-1	---	6.00-20.00	0.03-0.04	0.0-2.9	0.0-0.1	---	---	-	---	0
	6-60			0-5	---	6.00-20.00	0.04-0.06	0.0-2.9	---	---	---			
Bc: Bridgeport---	0-10	7	64	27-32	1.35-1.45	0.60-2.00	0.21-0.23	3.0-5.9	1.0-3.0	.32	.32	5	4L	86
	10-65	8	68	18-30	1.40-1.50	0.60-2.00	0.17-0.22	0.0-2.9	0.5-2.0	.43	.43			
Bd: Bridgeport---	0-10	7	64	27-32	1.35-1.45	0.60-2.00	0.21-0.23	3.0-5.9	1.0-3.0	.32	.32	5	4L	86
	10-65	8	68	18-30	1.40-1.50	0.60-2.00	0.17-0.22	0.0-2.9	0.5-2.0	.43	.43			
Bo: Optima-----	0-8	87	7	2-10	1.70-1.80	6.00-20.00	0.08-0.10	0.0-2.9	0.0-1.0	.17	.17	5	2	134
	8-17	93	1	2-10	1.70-1.80	6.00-20.00	0.08-0.10	0.0-2.9	0.0-0.5	.17	.17			
	17-80	94	1	2-8	1.70-1.80	6.00-20.00	0.05-0.08	---	0.0-0.0	.15	.15			
Da: Dale-----	0-25	11	68	15-26	1.40-1.50	0.60-2.00	0.22-0.24	0.0-2.9	1.0-3.0	.32	.32	5	5	56
	25-60	9	64	18-35	1.50-1.60	0.60-2.00	0.17-0.22	3.0-5.9	0.5-1.0	.43	.43			
Gr: Schamber-----	0-6	64	14	18-25	1.50-1.60	6.00-20.00	0.03-0.06	0.0-2.9	0.5-1.0	.17	.28	5	8	0
	6-60	85	9	2-10	1.55-1.65	6.00-20.00	0.03-0.06	0.0-2.9	0.0-0.5	.10	.43			
GRP: Pits-----	0-60	95	1	0-8	1.70-2.00	6.00-20.00	0.02-0.09	0.0-2.9	0.0-0.5	.10	.17	2	8	0

PHYSICAL PROPERTIES OF THE SOILS--Continued  
Gray County, Kansas: Maintenance needed

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Ha: Harney-----	0-9	24	51	22-27	1.30-1.40	0.60-2.00	0.22-0.24	0.0-2.9	1.0-3.0	.32	.32	5	6	48
	9-25	7	54	35-42	1.40-1.50	0.20-0.60	0.12-0.23	3.0-5.9	0.5-1.0	.37	.37			
	25-65	18	52	24-35	1.25-1.35	0.60-2.00	0.18-0.22	0.0-2.9	0.0-0.5	.43	.43			
HB: Harney-----	0-6	24	51	22-27	1.30-1.40	0.60-2.00	0.22-0.24	0.0-2.9	1.0-3.0	.32	.32	5	6	48
	6-20	7	54	35-42	1.40-1.50	0.20-0.60	0.11-0.20	3.0-5.9	0.5-1.5	.37	.37			
	20-60	20	50	24-35	1.25-1.35	0.60-2.00	0.18-0.22	0.0-2.9	0.0-0.5	.43	.43			
INL: Aguolls-----	0-72			---	---	---	---	---	---	---	---	-	---	0
La: Las Animas, occasionally flooded-----	0-11	67	20	8-18	1.50-1.60	2.00-6.00	0.13-0.15	0.0-2.9	0.5-1.0	.24	.24	5	3	86
	11-60	91		8-18	1.55-1.65	2.00-6.00	0.09-0.19	0.0-2.9	0.0-1.0	.24	.24			
	0-14	35	33	28-35	1.30-1.40	0.20-0.60	0.17-0.19	3.0-5.9	1.0-2.0	.28	.28	4	4L	86
	14-25	34	37	22-35	1.35-1.45	0.20-0.60	0.15-0.20	3.0-5.9	---	.32	.32			
	25-60	91	6	1-5	1.45-1.55	6.00-20.00	0.02-0.07	0.0-2.9	---	.15	.15			
Lc: Las Animas, occasionally flooded-----	0-11	84	9	4-10	1.50-1.60	6.00-20.00	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	11-30	63	24	8-18	1.50-1.60	2.00-6.00	0.09-0.19	0.0-2.9	0.0-1.0	.24	.24			
	30-60	97	1	0-5	1.50-1.60	6.00-20.00	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Ld: Las Animas---	0-11	67	20	8-18	1.45-1.55	2.00-6.00	0.13-0.15	0.0-2.9	0.5-1.0	.24	.24	5	3	86
	11-30	63	24	8-18	1.50-1.60	2.00-6.00	0.09-0.19	0.0-2.9	0.0-1.0	.24	.24			
	30-60	97	1	0-5	1.50-1.60	6.00-20.00	0.05-0.07	0.0-2.9	0.0-0.5	.15	.15			
Le: Las, occasionally flooded-----	0-8	35	34	27-35	1.35-1.45	0.20-0.60	0.17-0.19	3.0-5.9	0.5-1.0	.32	.32	4	4L	86
	8-31	35	38	18-35	1.40-1.50	0.20-0.60	0.12-0.19	3.0-5.9	0.1-0.5	.32	.32			
	31-60	91	6	1-5	1.50-1.60	6.00-20.00	0.02-0.04	0.0-2.9	0.0-0.5	.10	.10			
Lh: Lesho, occasionally flooded-----	0-14	35	33	28-35	1.30-1.40	0.20-0.60	0.17-0.19	3.0-5.9	1.0-2.0	.28	.28	5	4L	86
	14-25	35	38	18-35	1.35-1.45	0.20-0.60	0.15-0.19	3.0-5.9	0.0-0.5	.32	.32			
	25-60	92	4	1-8	1.45-1.55	2.00-20.00	0.02-0.07	0.0-2.9	0.0-0.5	.15	.15			
Lk: Lesho-----	0-14	40	38	18-27	1.30-1.40	0.60-2.00	0.20-0.22	0.0-2.9	1.0-2.0	.28	.28	5	4L	86
	14-25	35	38	18-35	1.35-1.45	0.20-0.60	0.15-0.19	3.0-5.9	---	.32	.32			
	25-60	92	4	1-8	1.45-1.55	2.00-20.00	0.02-0.07	0.0-2.9	---	.15	.15			
	0-17	35	38	18-35	1.40-1.50	0.20-0.60	0.17-0.19	0.0-2.9	0.5-1.0	.28	.28	3	4L	86
	17-60	90	2	3-15	1.50-1.60	5.95-19.98	0.05-0.11	0.0-2.9	0.0-0.5	.17	.17			
Ll: Lincoln, occasionally flooded-----	0-8	66	20	10-18	1.50-1.60	6.00-20.00	0.16-0.18	0.0-2.9	0.5-1.0	.24	.24	5	3	86
	8-60	43	47	5-15	1.40-1.50	6.00-20.00	0.05-0.19	0.0-2.9	0.0-0.5	.17	.17			
Lo: Lofton-----	0-6	19	48	27-40	1.10-1.30	0.20-0.60	0.19-0.21	3.0-5.9	1.0-3.0	.37	.37	5	4	86
	6-36	7	48	35-50	1.10-1.30	0.00-0.06	0.14-0.18	6.0-8.9	0.5-4.5	.28	.28			
	36-60	18	52	20-35	1.10-1.30	0.60-2.00	0.18-0.20	0.0-2.9	0.2-0.8	.43	.43			
Lu: Lubbock-----	0-15	39	37	20-27	1.45-1.55	0.60-2.00	0.20-0.22	0.0-2.9	1.0-2.0	.28	.28	5	6	48
	15-36	7	53	35-45	1.35-1.45	0.20-0.60	0.09-0.20	3.0-5.9	0.5-1.0	.32	.32			
	36-60	35	38	20-35	1.40-1.50	0.60-2.00	0.18-0.22	0.0-2.9	0.0-0.7	.32	.32			
Md: Penden-----	0-14	35	33	28-35	1.35-1.45	0.60-2.00	0.17-0.19	3.0-5.9	1.0-2.0	.28	.28	5	4L	86
	14-32	34	37	24-35	1.40-1.50	0.60-2.00	0.15-0.20	3.0-5.9	0.5-1.0	.32	.32			
	32-60	34	37	24-35	1.30-1.50	0.60-2.00	0.14-0.19	3.0-5.9	0.1-0.5	.32	.32			
Me: Penden-----	0-15	35	33	28-35	1.35-1.45	0.60-2.00	0.17-0.19	3.0-5.9	0.5-1.0	.28	.28	5	4L	86
	15-34	34	37	24-35	1.40-1.50	0.60-2.00	0.15-0.20	3.0-5.9	0.0-1.0	.32	.32			
	34-60	34	37	24-35	1.40-1.50	0.60-2.00	0.14-0.19	3.0-5.9	0.1-0.5	.32	.32			
Mf: Penden, Severely eroded-----	0-4	35	33	28-35	1.35-1.45	0.60-2.00	0.17-0.19	3.0-5.9	0.5-1.0	.28	.28	4	4L	86
	4-22	34	37	24-35	1.40-1.50	0.60-2.00	0.15-0.20	3.0-5.9	0.0-1.0	.32	.32			
	22-30	34	37	24-35	1.40-1.50	0.60-2.00	0.14-0.19	3.0-5.9	0.2-0.5	.32	.32			
	30-80	34	37	24-35	1.40-1.50	0.60-2.00	0.14-0.19	3.0-5.9	0.2-0.5	.32	.32			
Mh: Penden, Severely Eroded-----	0-4	35	33	28-35	1.35-1.45	0.60-2.00	0.17-0.19	3.0-5.9	0.5-1.0	.28	.28	4	4L	86
	4-22	34	37	24-35	1.40-1.50	0.60-2.00	0.15-0.20	3.0-5.9	0.0-1.0	.32	.32			
	22-30	34	37	24-35	1.40-1.50	0.60-2.00	0.14-0.19	3.0-5.9	0.2-0.5	.32	.32			
	30-80	34	37	24-35	1.40-1.50	0.60-2.00	0.14-0.19	3.0-5.9	0.2-0.5	.32	.32			



PHYSICAL PROPERTIES OF THE SOILS--Continued  
Gray County, Kansas: Maintenance needed

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Mm: Manter-----	0-18	65	20	10-20	1.35-1.45	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.20	.20	5	3	86
	18-30	66	20	9-18	1.40-1.50	2.00-6.00	0.12-0.17	0.0-2.9	0.2-0.5	.24	.24			
	30-62	46	44	5-15	1.50-1.60	2.00-6.00	0.08-0.13	0.0-2.9	0.0-0.5	.20	.20			
Mn: Manter-----	0-18	65	20	10-20	1.35-1.45	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.20	.20	5	3	86
	18-30	66	20	9-18	1.40-1.50	2.00-6.00	0.12-0.17	0.0-2.9	0.2-0.5	.24	.24			
	30-62	67	23	5-15	1.50-1.60	2.00-6.00	0.08-0.13	0.0-2.9	0.0-0.5	.24	.24			
Mo: Manter-----	0-18	65	20	10-20	1.30-1.40	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.20	.20	5	3	86
	18-30	66	20	9-18	1.40-1.50	2.00-6.00	0.12-0.17	0.0-2.9	0.2-0.5	.24	.24			
	30-62	67	23	5-15	1.50-1.60	2.00-6.00	0.08-0.13	0.0-2.9	0.0-0.5	.24	.24			
Mp: Campus-----	0-9	42	37	15-27	1.25-1.35	0.60-2.00	0.20-0.22	0.0-2.9	0.5-2.0	.28	.28	2	4L	86
	9-19	38	36	18-35	1.30-1.40	0.60-2.00	0.15-0.19	0.0-2.9	0.5-1.0	.32	.32			
	>19			---	---	0.00-0.60	---	---	---	---	---			
Canlon-----	0-8	27	54	12-27	1.35-1.45	0.60-2.00	0.22-0.24	0.0-2.9	0.5-1.0	.37	.37	1	4L	86
	>8			---	---	0.00-0.60	---	---	---	---	---			
Mu: Manter-----	0-18	65	20	10-20	1.30-1.40	2.00-6.00	0.16-0.18	0.0-2.9	0.5-1.0	.20	.20	5	3	86
	18-30	66	20	9-18	1.40-1.50	2.00-6.00	0.12-0.17	0.0-2.9	0.2-0.5	.24	.24			
	30-62	67	23	5-15	1.50-1.60	2.00-6.00	0.08-0.13	0.0-2.9	0.0-0.5	.24	.24			
Ulysses-----	0-5	12	70	10-27	1.15-1.25	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	5-14	7	66	21-32	1.25-1.35	0.60-2.00	0.21-0.24	3.0-5.9	0.5-1.0	.43	.43			
	14-60	10	68	18-27	1.25-1.35	0.60-2.00	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
Pa: Pratt-----	0-11	79	16	2-8	1.45-1.55	6.00-20.00	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	11-26	86	7	4-11	1.45-1.55	6.00-20.00	0.09-0.17	0.0-2.9	0.1-0.5	.17	.17			
	26-60	79	16	1-8	1.50-1.60	6.00-20.00	0.05-0.10	0.0-2.9	0.0-0.3	.17	.17			
Pg: Pratt-----	0-11	79	16	2-8	1.45-1.55	6.00-20.00	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	11-40	86	7	4-11	1.45-1.55	6.00-20.00	0.09-0.17	0.0-2.9	0.1-0.5	.17	.17			
	40-60	79	16	1-8	1.50-1.60	6.00-20.00	0.05-0.10	0.0-2.9	0.0-0.3	.17	.17			
Pt: Pratt-----	0-11	79	16	2-8	1.45-1.55	6.00-20.00	0.10-0.12	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	11-26	86	7	4-11	1.45-1.55	6.00-20.00	0.09-0.17	0.0-2.9	0.1-0.5	.17	.17			
	26-60	79	16	1-8	1.50-1.60	6.00-20.00	0.05-0.10	0.0-2.9	0.0-0.3	.17	.17			
Optima-----	0-8	87	7	2-10	1.70-1.80	6.00-20.00	0.08-0.10	0.0-2.9	0.0-1.0	.17	.17	5	2	134
	8-17	93	1	2-10	1.70-1.80	6.00-20.00	0.08-0.10	0.0-2.9	0.0-0.5	.17	.17			
	17-80	94	1	2-8	1.70-1.80	6.00-20.00	0.05-0.08	---	0.0-0.0	.15	.15			
Ra: Ness-----	0-31	22	28	40-60	1.35-1.45	0.00-0.06	0.09-0.13	6.0-8.9	1.0-3.0	.28	.28	5	4	86
	31-60	20	50	20-40	1.35-1.45	0.06-2.00	0.18-0.22	3.0-5.9	0.1-1.0	.43	.43			
Rm: Richfield----	0-5	29	54	10-24	1.30-1.40	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	5-17	7	54	35-42	1.40-1.50	0.20-0.60	0.11-0.20	3.0-5.9	0.5-1.2	.37	.37			
	17-60	20	54	18-35	1.25-1.35	0.60-2.00	0.18-0.22	3.0-5.9	0.2-0.5	.43	.43			
Rn: Richfield----	0-6	29	54	10-24	1.30-1.40	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	6-19	7	54	35-42	1.40-1.50	0.20-0.60	0.11-0.20	6.0-8.9	0.5-1.2	.37	.37			
	19-60	24	50	18-35	1.25-1.35	0.60-2.00	0.14-0.22	3.0-5.9	0.2-0.5	.43	.43			
Ro: Richfield----	0-6	18	52	27-32	1.40-1.50	0.20-0.60	0.21-0.23	3.0-5.9	1.0-2.0	.32	.32	5	7	38
	6-19	7	54	35-42	1.40-1.50	0.20-0.60	0.11-0.20	6.0-8.9	0.5-1.2	.37	.37			
	19-60	24	50	18-35	1.25-1.35	0.60-2.00	0.14-0.22	3.0-5.9	0.2-0.5	.43	.43			
Rs: Richfield----	0-6	29	54	10-24	1.30-1.40	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	6-19	7	54	35-42	1.40-1.50	0.20-0.60	0.11-0.20	6.0-8.9	0.5-1.2	.37	.37			
	19-60	24	50	18-35	1.25-1.35	0.60-2.00	0.14-0.22	3.0-5.9	0.2-0.5	.43	.43			
Spearville---	0-7	20	53	20-35	1.20-1.30	0.20-0.60	0.21-0.23	3.0-5.9	1.0-3.0	.32	.32	5	7	38
	7-20	6	48	42-50	1.35-1.45	0.06-0.20	0.11-0.13	3.0-5.9	0.5-1.0	.32	.32			
	20-26	7	51	35-50	1.35-1.45	0.20-0.60	0.11-0.20	6.0-8.9	0.0-0.5	.37	.37			
	26-60	18	50	25-40	1.30-1.40	0.20-0.60	0.18-0.22	3.0-5.9	0.0-0.5	.43	.43			
Sa: Satanta-----	0-10	43	40	10-25	1.30-1.40	0.60-2.00	0.20-0.22	0.0-2.9	1.0-2.0	.28	.28	5	6	48
	10-32	35	38	18-35	1.35-1.45	0.60-2.00	0.15-0.19	3.0-5.9	0.7-1.0	.32	.32			
	32-50	27	54	10-28	1.40-1.50	0.60-2.00	0.14-0.19	0.0-2.9	0.3-0.7	.32	.32			
SAP: Sand Pit-----	---	---	---	---	---	---	---	---	---	---	---	-	---	---
Sb: Satanta-----	0-10	43	40	10-25	1.30-1.40	0.60-2.00	0.20-0.22	0.0-2.9	1.0-2.0	.28	.28	5	6	48
	10-32	35	38	18-35	1.35-1.45	0.60-2.00	0.15-0.19	3.0-5.9	0.7-1.0	.32	.32			
	32-50	27	54	10-28	1.40-1.50	0.60-2.00	0.14-0.19	0.0-2.9	0.3-0.7	.32	.32			
Sp: Spearville---	0-7	20	53	20-35	1.20-1.30	0.20-0.60	0.21-0.23	3.0-5.9	1.0-3.0	.32	.32	5	7	38
	7-20	6	48	42-50	1.35-1.45	0.06-0.20	0.11-0.13	3.0-5.9	0.5-1.0	.32	.32			
	20-26	7	51	35-50	1.35-1.45	0.20-0.60	0.11-0.20	6.0-8.9	0.0-0.5	.37	.37			
	26-60	18	50	25-40	1.30-1.40	0.20-0.60	0.18-0.22	3.0-5.9	0.0-0.5	.43	.43			
Sr: Spearville---	0-7	20	53	20-35	1.20-1.30	0.20-0.60	0.21-0.23	3.0-5.9	1.0-3.0	.32	.32	5	7	38
	7-20	6	48	42-50	1.35-1.45	0.06-0.20	0.11-0.13	3.0-5.9	0.5-1.0	.32	.32			
	20-26	7	51	35-50	1.35-1.45	0.20-0.60	0.11-0.20	6.0-8.9	0.0-0.5	.37	.37			
	26-60	18	50	25-40	1.30-1.40	0.20-0.60	0.18-0.22	3.0-5.9	0.0-0.5	.43	.43			

PHYSICAL PROPERTIES OF THE SOILS--Continued  
Gray County, Kansas: Maintenance needed

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Sw:														
Sweetwater---	0-17	35	38	18-35	1.40-1.50	0.20-0.60	0.17-0.19	0.0-2.9	0.5-1.0	.28	.28	3	4L	86
	17-60	65	26	3-15	1.50-1.60	5.95-19.98	0.05-0.11	0.0-2.9	0.0-0.5	.17	.17			
Tf:														
Optima-----	0-8	93	1	2-10	1.70-1.80	6.00-20.00	0.08-0.10	0.0-2.9	0.0-1.0	.17	.17	5	2	134
	8-17	93	1	2-10	1.70-1.80	6.00-20.00	0.08-0.10	0.0-2.9	0.0-0.5	.17	.17			
	17-80	94	1	2-8	1.70-1.80	6.00-20.00	0.05-0.08	---	0.0-0.0	.15	.15			
Ua:														
Ulysses-----	0-5	12	70	10-27	1.15-1.25	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	5-14	7	66	21-32	1.25-1.35	0.60-2.00	0.18-0.22	3.0-5.9	0.5-1.0	.43	.43			
	14-60	10	68	18-27	1.25-1.35	0.60-2.00	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
Ub:														
Ulysses-----	0-5	12	70	10-27	1.15-1.25	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	5-14	7	66	21-32	1.25-1.35	0.60-2.00	0.18-0.22	3.0-5.9	0.5-1.0	.43	.43			
	14-60	10	68	18-27	1.25-1.35	0.60-2.00	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
UBB:														
Uly-----	0-8	11	67	17-27	1.20-1.30	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	8-55	9	66	20-30	1.20-1.30	0.60-2.00	0.18-0.22	0.0-2.9	0.7-1.5	.43	.43			
	55-60	10	68	18-27	1.10-1.20	0.60-2.00	0.17-0.22	0.0-2.9	0.4-0.7	.43	.43			
Uc:														
Ulysses-----	0-6	12	70	10-27	1.15-1.25	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	6-16	7	66	21-32	1.25-1.35	0.60-2.00	0.18-0.22	3.0-5.9	0.5-1.0	.43	.43			
	16-60	10	68	18-27	1.25-1.35	0.60-2.00	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
UCC:														
Uly-----	0-6	11	67	17-27	1.20-1.30	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	6-14	7	68	20-30	1.20-1.30	0.60-2.00	0.21-0.24	0.0-2.9	0.5-1.0	.43	.43			
	14-60	10	68	18-27	1.10-1.20	0.60-2.00	0.17-0.22	0.0-2.9	0.2-0.5	.43	.43			
Um:														
Ulysses-----	0-5	12	70	10-27	1.15-1.25	0.60-2.00	0.22-0.24	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	5-14	7	66	21-32	1.25-1.35	0.60-2.00	0.18-0.22	3.0-5.9	0.5-1.0	.43	.43			
	14-60	10	68	18-27	1.25-1.35	0.60-2.00	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
Colby-----	0-5	11	68	15-27	1.20-1.30	0.60-2.00	0.22-0.24	0.0-2.9	0.5-1.0	.43	.43	5	4L	86
	5-60	10	68	18-27	1.30-1.40	0.60-2.00	0.17-0.22	0.0-2.9	0.0-0.5	.43	.43			
W:														
Water-----	---			---	---	---	---	---	---	---	---	-	---	---

CHEMICAL PROPERTIES OF THE SOILS  
Gray County, Kansas

The Chemical Properties table shows estimates of some characteristics and features that affect soil behavior. These estimates are given for the major layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils. Depth to the upper and lower boundaries of each layer is indicated.

Cation-exchange capacity is the total amount of extractable bases that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. Soils having a high cation-exchange capacity can retain cations. The ability to retain cations helps to prevent the pollution of ground water.

Soil reaction is a measure of acidity or alkalinity and is expressed as a range in pH values. The range in pH of each major horizon is based on many field tests. For many soils, values have been verified by laboratory analyses. Soil reaction is important in selecting crops and other plants, in evaluating soil amendments for fertility and stabilization, and in determining the risk of corrosion.

Calcium carbonate equivalent is the percent of carbonates, by weight, in the fraction of the soil less than 2 millimeters in size. The availability of plant nutrients is influenced by the amount of carbonates in the soil. Incorporating nitrogen fertilizer into calcareous soils helps to prevent nitrite accumulation and ammonium-N volatilization.

Gypsum is expressed as a percent, by weight, of hydrated calcium sulfates in the fraction of the soil less than 20 millimeters in size. Gypsum is partially soluble in water and can be dissolved and removed by water. Soils that have a high content of gypsum may collapse if the gypsum is removed by percolating water.

Salinity is a measure of soluble salts in the soil at saturation. It is expressed as the electrical conductivity of the saturation extract, in millimhos per centimeter at 25 degrees C. Estimates are based on field and laboratory measurements at representative sites of nonirrigated soils. The salinity of irrigated soils is affected by the quality of the irrigation water and by the frequency of water application. Hence, the salinity of soils in individual fields can differ greatly from the value given in the table. Salinity affects the suitability of a soil for crop production, the stability of soil if used as construction material, and the potential of the soil to corrode metal and concrete.

Sodium adsorption ratio (SAR) is a measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste. It is the ratio of the Na concentration divided by the square root of one-half of the Ca + Mg concentration. Soils that have SAR values of 13 or more may be characterized by an increased dispersion of organic matter and clay particles, reduced permeability and aeration, and a general degradation of soil structure.

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Gray County, Kansas

PAGE 2 of 5

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
055BP: Bridgeport, rarely flooded-	0-11	11-21	6.6-8.4	1-5	0	0.0-2.0	0
	11-60	10-38	7.4-8.4	5-10	0	0.0-2.0	0-2
055LD: Las, occasionally flooded-----	0-8	11-24	7.4-8.4	1-5	0	0.0-4.0	0
	8-31	7.0-21	7.4-8.4	5-15	0	0.0-4.0	0
	31-60	2.0-10	7.4-8.4	1-15	0	0.0-4.0	0-4
Las Animas, occasionally flooded-----	0-14	3.0-11	7.4-8.4	1-5	---	0.0-4.0	---
	14-30	3.0-11	7.4-8.4	1-5	---	0.0-2.0	---
	30-60	0.0-3.0	7.4-8.4	5-10	---	---	---
055TF: Valent-----	0-4	1.0-4.0	6.6-7.8	---	---	---	---
	4-50	0.0-5.0	6.6-7.8	1-5	0	0	0
055TV: Valent-----	0-4	1.0-7.0	6.6-7.8	0	0	0	0
	4-60	0.0-5.0	6.6-7.8	1-5	0	0	0
Vona-----	0-8	1.0-5.0	6.6-7.8	0	0	0.0-2.0	0
	8-22	3.0-11	6.6-8.4	1-5	0	0.0-4.0	0
	22-60	1.0-9.0	7.9-9.0	1-5	0	0.0-4.0	0
055VO: Vona-----	0-8	1.0-5.0	6.6-7.8	0	0	0.0-2.0	0
	8-22	3.0-11	6.6-8.4	1-5	0	0.0-4.0	0
	22-60	1.0-9.0	7.9-9.0	1-5	0	0.0-4.0	0
081LU: Lubbock-----	0-9	11-23	6.6-7.8	---	---	---	---
	9-42	14-27	6.6-8.4	1-5	---	---	---
	42-72	8.0-21	7.9-8.4	5-10	---	---	---
083UM: Uly-----	0-6	10-25	6.1-7.8	0	0	0	0
	6-19	10-25	6.1-8.4	1-5	0	0	0
	19-60	10-23	7.4-8.4	5-10	0	0	0
Coly-----	0-6	7.0-15	7.4-8.4	5-10	0	0	0
	6-60	7.0-14	7.4-8.4	10-15	0	0	0
119HC: Harney, eroded--	0-6	11-23	5.6-7.8	0	0	0	0
	6-20	14-25	6.1-8.4	1-5	0	0	0
	20-60	9.0-21	7.4-8.4	5-10	0	0	0
119RX: Roxbury-----	0-20	7.0-18	7.4-8.4	1-5	---	---	---
	20-60	7.0-21	7.4-8.4	1-5	---	---	---
119UA: Uly-----	0-6	4.0-18	6.6-7.8	0-5	---	---	---
	6-12	8.0-25	7.4-8.4	1-10	---	---	---
	12-18	8.0-19	7.4-8.4	1-15	---	---	---
	18-32	7.0-20	7.9-8.6	5-10	---	---	---
	32-62	7.0-16	7.9-8.6	1-10	---	---	---
119UT: Uly-----	0-6	7.0-18	6.1-7.8	0	0	0	0
	6-14	8.0-18	6.1-8.4	1-5	0	0	0
	14-60	7.0-16	7.4-8.4	5-10	0	0	0
An: Bridgeport, channeled-----	0-12	6.0-18	6.6-8.4	1-5	0	0	0
	12-60	7.0-18	7.4-8.4	5-10	0	0	0
ARR: Arkansas River--	0-6	---	---	---	---	---	---
	6-60	---	---	---	---	---	---
Bc: Bridgeport-----	0-10	11-21	6.6-8.4	1-5	0	0.0-2.0	0-1
	10-65	10-38	7.4-8.4	5-10	0	0.0-2.0	0-2
Bd: Bridgeport-----	0-10	11-21	6.6-8.4	1-5	0	0.0-2.0	0-1
	10-65	10-38	7.4-8.4	5-10	0	0.0-2.0	0-2

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Gray County, Kansas

PAGE 3 of 5

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Bo:							
Optima-----	0-8	2.0-8.0	6.6-7.3	0	0	0	0
	8-17	0.0-5.0	6.6-7.3	0	0	0	0
	17-80	0.0-5.0	6.6-7.8	0-5	0	0	0
Da:							
Dale-----	0-25	6.0-18	6.1-7.8	0	0	0	0
	25-60	7.0-21	7.4-8.4	1-5	0	0	0
Gr:							
Schamber-----	0-6	7.0-16	6.1-8.4	1-5	---	0.0-2.0	---
	6-60	0.0-6.0	7.4-8.4	5-10	---	0.0-2.0	---
GRP:							
Pits-----	0-60	0.0-5.0	6.6-8.4	0	0	0	0
Ha:							
Harney-----	0-9	9.0-18	5.6-7.8	0	0	0	0
	9-25	14-25	6.1-8.4	1-5	0	0	0
	25-65	9.0-21	7.4-8.4	5-10	0	0	0
HB:							
Harney-----	0-6	9.0-18	5.6-7.8	0	0	0	0
	6-20	14-25	6.1-8.4	1-5	0	0	0
	20-60	9.0-21	7.4-8.4	5-10	0	0	0
INL:							
Aquolls-----	0-72	---	---	---	---	---	---
La:							
Las Animas, occasionally flooded-----	0-11	3.0-11	7.4-9.0	1-5	---	2.0-8.0	---
	11-60	3.0-11	7.8-9.6	1-5	---	4.0-8.0	---
Lesho, occasionally flooded-----	0-14	11-22	7.4-9.0	1-5	---	2.0-8.0	---
	14-25	8.0-21	7.9-9.0	5-10	---	4.0-16.0	---
	25-60	0.0-3.0	7.4-9.0	5-10	---	2.0-8.0	---
Lc:							
Las Animas, occasionally flooded-----	0-11	1.0-7.0	7.4-8.4	1-5	---	0.0-4.0	---
	11-30	3.0-11	7.4-8.4	1-5	---	0.0-2.0	---
	30-60	0.0-3.0	7.4-8.4	5-10	---	---	---
Ld:							
Las Animas-----	0-11	3.0-11	7.4-8.4	1-5	---	0.0-4.0	---
	11-30	3.0-11	7.4-8.4	1-5	---	0.0-2.0	---
	30-60	0.0-3.0	7.4-8.4	5-10	---	---	---
Le:							
Las, occasionally flooded-----	0-8	11-22	7.4-8.4	1-5	0	0.0-4.0	0
	8-31	7.0-21	7.4-8.4	5-15	0	0.0-4.0	0
	31-60	2.0-10	7.4-8.4	1-15	0	0.0-4.0	0-4
Lh:							
Lesho, occasionally flooded-----	0-14	11-22	7.4-8.4	1-5	---	0.0-4.0	---
	14-25	7.0-21	7.4-8.4	1-5	---	0.0-4.0	---
	25-60	0.0-5.0	7.4-9.0	5-10	---	0.0-4.0	---
Lk:							
Lesho-----	0-14	7.0-18	7.4-8.4	1-5	---	0.0-4.0	---
	14-25	7.0-21	7.4-8.4	1-5	---	0.0-4.0	---
	25-60	0.0-5.0	7.4-9.0	5-10	---	0.0-4.0	---
Sweetwater-----	0-17	7.0-22	7.4-8.4	1-5	0	0	0
	17-60	1.0-9.0	7.9-8.4	5-10	0	0.0-4.0	0-5
Ll:							
Lincoln, occasionally flooded-----	0-8	4.0-11	7.4-8.4	---	---	---	---
	8-60	2.0-9.0	7.9-8.4	1-5	---	---	---
Lo:							
Lofton-----	0-6	11-27	6.6-7.3	---	---	0	---
	6-36	14-27	6.6-7.8	---	---	0	---
	36-60	8.0-19	7.4-8.4	1-10	---	0.0-2.0	---

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Gray County, Kansas

PAGE 4 of 5

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Lu:							
Lubbock-----	0-15	8.0-18	6.6-7.8	---	---	---	---
	15-36	14-27	6.6-8.4	1-5	---	---	---
	36-60	8.0-21	7.9-8.4	5-10	---	---	---
Md:							
Penden-----	0-14	11-22	7.4-8.4	5-10	---	0.0-2.0	0
	14-32	9.0-21	7.9-8.4	15-30	---	0.0-2.0	0
	32-60	9.0-21	7.9-8.4	15-30	---	0.0-2.0	0
Me:							
Penden-----	0-15	11-22	7.4-8.4	10-15	---	0.0-2.0	0
	15-34	9.0-21	7.9-8.4	15-30	---	0.0-2.0	0
	34-60	9.0-21	7.9-8.4	15-30	---	0.0-2.0	0
Mf:							
Penden, Severely eroded-----	0-4	11-22	7.4-8.4	10-15	---	---	---
	4-22	9.0-21	7.9-8.4	15-30	---	---	---
	22-30	9.0-21	7.9-8.4	15-30	---	---	---
	30-80	9.0-21	7.9-8.4	15-30	---	---	---
Mh:							
Penden, Severely Eroded-----	0-4	11-22	7.4-8.4	10-15	---	---	---
	4-22	9.0-21	7.9-8.4	15-30	---	---	---
	22-30	9.0-21	7.9-8.4	15-30	---	---	---
	30-80	9.0-21	7.9-8.4	15-30	---	---	---
Mm:							
Manter-----	0-18	4.0-13	6.6-7.8	0	0	0	0
	18-30	3.0-11	6.6-7.8	1-5	0	0	0
	30-62	2.0-9.0	7.9-8.4	1-5	0	0.0-2.0	0
Mn:							
Manter-----	0-18	4.0-13	6.6-7.8	0	0	0	0
	18-30	3.0-11	6.6-7.8	1-5	0	0	0
	30-62	2.0-9.0	7.9-8.4	1-5	0	0.0-2.0	0
Mo:							
Manter-----	0-18	4.0-13	6.6-7.8	0	0	0	0
	18-30	3.0-11	6.6-7.8	1-5	0	0	0
	30-62	2.0-9.0	7.9-8.4	1-5	0	0.0-2.0	0
Mp:							
Campus-----	0-9	6.0-17	7.4-8.4	15-30	---	---	---
	9-19	7.0-21	7.4-8.4	15-30	---	---	---
	>19	---	---	---	---	---	---
Canlon-----	0-8	5.0-17	7.4-8.4	5-10	0	0	0
	>8	---	---	---	---	---	---
Mu:							
Manter-----	0-18	4.0-13	6.6-7.8	0	0	0	0
	18-30	3.0-11	6.6-7.8	1-5	0	0	0
	30-62	2.0-9.0	7.9-8.4	1-5	0	0.0-2.0	0
Ulysses-----	0-5	4.0-18	6.6-7.8	---	---	---	---
	5-14	8.0-19	7.4-8.4	5-10	---	---	---
	14-60	7.0-16	7.9-8.4	10-15	---	---	---
Pa:							
Pratt-----	0-11	1.0-5.0	5.6-7.3	---	---	---	---
	11-26	1.0-7.0	5.6-7.3	---	---	---	---
	26-60	0.0-5.0	6.1-7.3	1-5	---	---	---
Pg:							
Pratt-----	0-11	1.0-5.0	5.6-7.3	---	---	---	---
	11-40	1.0-7.0	5.6-7.3	---	---	---	---
	40-60	0.0-5.0	6.1-7.3	1-5	---	---	---
Pt:							
Pratt-----	0-11	1.0-5.0	5.6-7.3	---	---	---	---
	11-26	1.0-7.0	5.6-7.3	---	---	---	---
	26-60	0.0-5.0	6.1-7.3	1-5	---	---	---
Optima-----	0-8	2.0-8.0	6.6-7.3	0	0	0	0
	8-17	0.0-5.0	6.6-7.3	0	0	0	0
	17-80	0.0-5.0	6.6-7.8	0-5	0	0	0
Ra:							
Ness-----	0-31	16-38	6.1-8.4	---	---	---	---
	31-60	8.0-24	7.4-8.4	1-5	---	---	---
Rm:							
Richfield-----	0-5	4.0-16	6.6-7.8	0	0	0	0
	5-17	14-25	6.6-8.4	1-5	0	0	0
	17-60	7.0-21	7.9-9.0	5-10	0	0	0

CHEMICAL PROPERTIES OF THE SOILS--Continued  
Gray County, Kansas

PAGE 5 of 5

Map symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm	
Rn:							
Richfield-----	0-6	4.0-16	6.6-7.8	0	0	0	0
	6-19	14-25	6.6-8.4	1-5	0	0	0
	19-60	7.0-21	7.9-9.0	5-10	0	0	0
Ro:							
Richfield-----	0-6	11-21	6.6-7.8	0	0	0	0
	6-19	14-25	6.6-8.4	1-5	0	0	0
	19-60	7.0-21	7.9-9.0	5-10	0	0	0
Rs:							
Richfield-----	0-6	4.0-16	6.6-7.8	0	0	0	0
	6-19	14-25	6.6-8.4	1-5	0	0	0
	19-60	7.0-21	7.9-9.0	5-10	0	0	0
Spearville-----	0-7	8.0-23	6.6-7.8	---	---	---	---
	7-20	16-30	6.6-8.4	1-5	---	---	---
	20-26	14-30	7.9-8.4	1-5	---	---	---
	26-60	10-24	7.9-8.4	5-10	---	---	---
Sa:							
Satanta-----	0-10	4.0-16	6.1-7.8	---	---	---	---
	10-32	7.0-21	6.6-8.4	1-5	---	---	---
	32-50	4.0-17	7.4-8.4	1-5	---	---	---
SAP:							
Sand Pit-----	---	---	---	---	---	---	---
Sb:							
Satanta-----	0-10	4.0-16	6.1-7.8	---	---	---	---
	10-32	7.0-21	6.6-8.4	1-5	---	---	---
	32-50	4.0-17	7.4-8.4	1-5	---	---	---
Sp:							
Spearville-----	0-7	8.0-23	6.6-7.8	---	---	---	---
	7-20	16-30	6.6-8.4	1-5	---	---	---
	20-26	14-30	7.9-8.4	1-5	---	---	---
	26-60	10-24	7.9-8.4	5-10	---	---	---
Sr:							
Spearville-----	0-7	8.0-23	6.6-7.8	---	---	---	---
	7-20	16-30	6.6-8.4	1-5	---	---	---
	20-26	14-30	7.9-8.4	1-5	---	---	---
	26-60	10-24	7.9-8.4	5-10	---	---	---
Sw:							
Sweetwater-----	0-17	7.0-22	7.4-8.4	1-5	0	0	0
	17-60	1.0-9.0	7.9-8.4	5-10	0	0.0-4.0	0-5
Tf:							
Optima-----	0-8	2.0-8.0	6.6-7.3	0	0	0	0
	8-17	0.0-5.0	6.6-7.3	0	0	0	0
	17-80	0.0-5.0	6.6-7.8	0-5	0	0	0
Ua:							
Ulysses-----	0-5	4.0-18	6.6-7.8	---	---	---	---
	5-14	8.0-19	7.4-8.4	5-10	---	---	---
	14-60	7.0-16	7.9-8.4	10-15	---	---	---
Ub:							
Ulysses-----	0-5	4.0-18	6.6-7.8	---	---	---	---
	5-14	8.0-19	7.4-8.4	5-10	---	---	---
	14-60	7.0-16	7.9-8.4	10-15	---	---	---
UBB:							
Uly-----	0-8	10-25	6.1-7.8	0	0	0	0
	8-55	10-25	6.1-8.4	1-5	0	0	0
	55-60	10-23	7.4-8.4	5-10	0	0	0
Uc:							
Ulysses-----	0-6	4.0-18	6.6-7.8	---	---	---	---
	6-16	8.0-19	7.4-8.4	5-10	---	---	---
	16-60	7.0-16	7.9-8.4	10-15	---	---	---
UCC:							
Uly-----	0-6	7.0-18	6.1-7.8	0	0	0	0
	6-14	8.0-18	6.1-8.4	1-5	0	0	0
	14-60	7.0-16	7.4-8.4	5-10	0	0	0
Um:							
Ulysses-----	0-5	4.0-18	6.6-7.8	---	---	---	---
	5-14	8.0-19	7.4-8.4	5-10	---	---	---
	14-60	7.0-16	7.9-8.4	10-15	---	---	---
Colby-----	0-5	6.0-17	7.4-8.4	5-10	---	---	---
	5-60	7.0-16	7.4-8.4	10-15	---	---	---
W:							
Water-----	---	---	---	---	---	---	---

# WATER FEATURES Gray County, Kansas

The Water Features table gives estimates of various water features. The estimates are used in land use planning that involves engineering considerations. Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The four hydrologic soil groups are:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

The months in the table indicate the portion of the year in which the feature is most likely to be a concern.

Water table refers to a saturated zone in the soil. The Water Features table indicates, by month, depth to the top (upper limit) and base (lower limit) of the saturated zone in most years. Estimates of the upper and lower limits are based mainly on observations of the water table at selected sites and on evidence of a saturated zone, namely grayish colors or mottles (redoximorphic features) in the soil. A saturated zone that lasts for less than a month is not considered a water table. Ponding is standing water in a closed depression. Unless a drainage system is installed, the water is removed only by percolation, transpiration, or evaporation. The Water Features table indicates surface water depth and the duration and frequency of ponding. Duration is expressed as very brief if less than 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, rare, occasional, and frequent. None means that ponding is not probable; rare that it is unlikely but possible under unusual weather conditions (the chance of ponding is nearly 0 percent to 5 percent in any year); occasional that it occurs, on the average, once or less in 2 years (the chance of ponding is 5 to 50 percent in any year); and frequent that it occurs, on the average, more than once in 2 years (the chance of ponding is more than 50 percent in any year).

Flooding, the temporary inundation of an area, is caused by overflowing streams, by runoff from adjacent slopes, or by tides. Water standing for short periods after rainfall or snowmelt is not considered flooding, and water standing in swamps and marshes is considered ponding rather than flooding.

Duration and frequency are estimated. Duration is expressed as extremely brief if 0.1 hour to 4 hours, very brief if 4 hours to 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, very rare, rare, occasional, frequent, and very frequent. None means that flooding is not probable; very rare that it is very unlikely but possible under extremely unusual weather conditions (the chance of flooding is less than 1 percent in any year); rare that it is unlikely but possible under unusual weather conditions (the chance of flooding is 1 to 5 percent in any year); occasional that it occurs infrequently under normal weather conditions (the chance of flooding is 5 to 50 percent in any year); frequent that it is likely to occur often under normal weather conditions (the chance of flooding is more than 50 percent in any year but is less than 50 percent in all months in any year); and very frequent that it is likely to occur very often under normal weather conditions (the chance of flooding is more than 50 percent in all months of any year).

The information is based on evidence in the soil profile, namely thin strata of gravel, sand, silt, or clay deposited by floodwater; irregular decrease in organic matter content with increasing depth; and little or no horizon development.

Also considered are local information about the extent and levels of flooding and the relation of each soil on the landscape to historic floods. Information on the extent of flooding based on soil data is less specific than that provided by detailed engineering surveys that delineate flood-prone areas at specific flood frequency levels.



(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
055BP: Bridgeport, rarely flooded	B	April	---	---	---	---	---	Very brief	Rare
		May	---	---	---	---	---	Very brief	Rare
		June	---	---	---	---	---	Very brief	Rare
		July	---	---	---	---	---	Very brief	Rare
		August	---	---	---	---	---	Very brief	Rare
		September	---	---	---	---	---	Very brief	Rare
055LD: Las, occasionally flooded-	C	March	2.0-3.0	>6.0	---	---	---	---	None
		April	2.0-3.0	>6.0	---	---	---	Very brief	Occasional
		May	2.0-3.0	>6.0	---	---	---	Very brief	Occasional
		June	2.0-3.0	>6.0	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
Las Animas, occasionally flooded-----	C	January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
055TF: Valent-----	A		---	---	---	---	---	---	---
055TV: Valent-----	A		---	---	---	---	---	---	---
Vona-----	B		---	---	---	---	---	---	---
055VO: Vona-----	B		---	---	---	---	---	---	---
081LU: Lubbock-----	B		---	---	---	---	---	---	---
083UM: Uly-----	B		---	---	---	---	---	---	---
Coly-----	B		---	---	---	---	---	---	---
119HC: Harney, eroded-----	B		---	---	---	---	---	---	---
119RX: Roxbury-----	B	March	---	---	---	---	---	Very brief	Rare
		April	---	---	---	---	---	Very brief	Rare
		May	---	---	---	---	---	Very brief	Rare
		June	---	---	---	---	---	Very brief	Rare
		July	---	---	---	---	---	Very brief	Rare
		August	---	---	---	---	---	Very brief	Rare
119UA: Uly-----	B		---	---	---	---	---	---	---
119UT: Uly-----	B		---	---	---	---	---	---	---
An: Bridgeport, channeled----	B	April	---	---	---	---	---	Very brief	Frequent
		May	---	---	---	---	---	Very brief	Frequent
		June	---	---	---	---	---	Very brief	Frequent
		July	---	---	---	---	---	Very brief	Frequent
		August	---	---	---	---	---	Very brief	Frequent
		September	---	---	---	---	---	Very brief	Frequent
ARR:									

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Arkansas River-----	D		Ft	Ft	Ft				
		January	0.0-2.0	>6.0	---	---	---	Very long	Frequent
		February	0.0-2.0	>6.0	---	---	---	Very long	Frequent
		March	0.0-2.0	>6.0	---	---	---	Very long	Frequent
		April	0.0-2.0	>6.0	---	---	---	Very long	Frequent
		May	0.0-2.0	>6.0	---	---	---	Very long	Frequent
		June	0.0-2.0	>6.0	---	---	---	Very long	Frequent
		July	0.0-2.0	>6.0	---	---	---	Very long	Frequent
		August	0.0-2.0	>6.0	---	---	---	---	None
		September	0.0-2.0	>6.0	---	---	---	---	None
		October	0.0-2.0	>6.0	---	---	---	Very long	Frequent
		November	0.0-2.0	>6.0	---	---	---	Very long	Frequent
		December	0.0-2.0	>6.0	---	---	---	Very long	Frequent
Bc: Bridgeport-----	B								
		April	---	---	---	---	---	Very brief	Rare
		May	---	---	---	---	---	Very brief	Rare
		June	---	---	---	---	---	Very brief	Rare
		July	---	---	---	---	---	Very brief	Rare
		August	---	---	---	---	---	Very brief	Rare
		September	---	---	---	---	---	Very brief	Rare
Bd: Bridgeport-----	B								
		April	---	---	---	---	---	Very brief	Rare
		May	---	---	---	---	---	Very brief	Rare
		June	---	---	---	---	---	Very brief	Rare
		July	---	---	---	---	---	Very brief	Rare
		August	---	---	---	---	---	Very brief	Rare
		September	---	---	---	---	---	Very brief	Rare
Bo: Optima-----	A								
			---	---	---	---	---	---	---
Da: Dale-----	B								
		January	---	---	---	---	---	---	Rare
		February	---	---	---	---	---	---	Rare
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	---	---	---	---	---	---	Rare
		December	---	---	---	---	---	---	Rare
Gr: Schamber-----	A								
			---	---	---	---	---	---	---
GRP: Pits-----	A								
			---	---	---	---	---	---	---
Ha: Harney-----	B								
			---	---	---	---	---	---	---
HB: Harney-----	B								
			---	---	---	---	---	---	---
INL: Aquolls-----	C								
		March	0.0	>6.0	0.0-0.8	Brief	Occasional	---	None
		April	0.0	>6.0	0.0-0.8	Brief	Occasional	---	None
		May	0.0	>6.0	0.0-0.8	Brief	Occasional	---	None
		June	0.0	>6.0	0.0-0.8	Brief	Occasional	---	None
La: Las Animas, occasionally flooded-----	C								
		January	2.0-4.0	>6.0	---	---	---	---	None
		February	2.0-4.0	>6.0	---	---	---	---	None
		March	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		April	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		May	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		November	2.0-4.0	>6.0	---	---	---	---	None
		December	2.0-4.0	>6.0	---	---	---	---	None

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro-logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Lesho, occasionally flooded-----	C		Ft	Ft	Ft				
		March	2.0-4.0	>6.0	---	---	---	---	None
		April	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		May	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		June	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
Lc: Las Animas, occasionally flooded-----	C								
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Ld: Las Animas-----	C								
		January	1.5-3.0	>6.0	---	---	---	---	None
		February	1.5-3.0	>6.0	---	---	---	---	None
		March	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		April	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	1.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		November	1.5-3.0	>6.0	---	---	---	---	None
		December	1.5-3.0	>6.0	---	---	---	---	None
Le: Las, occasionally flooded-----	C								
		March	2.0-3.0	>6.0	---	---	---	---	None
		April	2.0-3.0	>6.0	---	---	---	Very brief	Occasional
		May	2.0-3.0	>6.0	---	---	---	Very brief	Occasional
		June	2.0-3.0	>6.0	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
Lh: Lesho, occasionally flooded-----	C								
		March	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		April	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		May	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		June	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
Lk: Lesho-----	C								
		March	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		April	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		May	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
		June	2.0-4.0	>6.0	---	---	---	Very brief	Occasional
Sweetwater-----	D								
		January	0.5-3.0	>6.0	---	---	---	---	None
		February	0.5-3.0	>6.0	---	---	---	---	None
		March	0.5-3.0	>6.0	---	---	---	---	None
		April	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		July	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		August	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		September	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		October	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		November	0.5-3.0	>6.0	---	---	---	---	None
		December	0.5-3.0	>6.0	---	---	---	---	None
Ll:									

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Lincoln, occasionally flooded-----	A		Ft	Ft	Ft				
		January	5.0-6.0	>6.0	---	---	---	---	None
		February	5.0-6.0	>6.0	---	---	---	---	None
		March	5.0-6.0	>6.0	---	---	---	---	None
		April	5.0-6.0	>6.0	---	---	---	Brief	Frequent
		May	5.0-6.0	>6.0	---	---	---	Brief	Frequent
		June	---	---	---	---	---	Brief	Frequent
		July	---	---	---	---	---	Brief	Frequent
		August	---	---	---	---	---	Brief	Frequent
		September	---	---	---	---	---	Brief	Frequent
		October	---	---	---	---	---	Brief	Frequent
		November	5.0-6.0	>6.0	---	---	---	---	None
		December	5.0-6.0	>6.0	---	---	---	---	None
Lo: Lofton-----	D		---	---	---	---	---	---	---
Lu: Lubbock-----	B		---	---	---	---	---	---	---
Md: Penden-----	B		---	---	---	---	---	---	---
Me: Penden-----	B		---	---	---	---	---	---	---
Mf: Penden, Severely eroded---	B		---	---	---	---	---	---	---
Mh: Penden, Severely Eroded---	B		---	---	---	---	---	---	---
Mm: Manter-----	B		---	---	---	---	---	---	---
Mn: Manter-----	B		---	---	---	---	---	---	---
Mo: Manter-----	B		---	---	---	---	---	---	---
Mp: Campus-----	B		---	---	---	---	---	---	---
Canlon-----	D		---	---	---	---	---	---	---
Mu: Manter-----	B		---	---	---	---	---	---	---
Ulysses-----	B		---	---	---	---	---	---	---
Pa: Pratt-----	A		---	---	---	---	---	---	---
Pg: Pratt-----	A		---	---	---	---	---	---	---
Pt: Pratt-----	A		---	---	---	---	---	---	---
Optima-----	A		---	---	---	---	---	---	---
Ra: Ness-----	D		---	---	---	---	---	---	---
		January	---	---	0.0-	---	---	---	None
		February	---	---	0.0-	---	---	---	None
		March	0.0	>6.0	0.0-1.0	Long	Frequent	---	None
		April	0.0	>6.0	0.0-1.0	Long	Frequent	---	None
		May	0.0	>6.0	0.0-1.0	Long	Frequent	---	None
		June	0.0	>6.0	0.0-1.0	Long	Frequent	---	None
		July	---	---	0.0-	---	---	---	None
		August	---	---	0.0-	---	---	---	None
		September	---	---	0.0-	---	---	---	None
		October	---	---	0.0-	---	---	---	None
		November	---	---	0.0-	---	---	---	None
		December	---	---	0.0-	---	---	---	None
Rm: Richfield-----	B		---	---	---	---	---	---	---
Rn: Richfield-----	B		---	---	---	---	---	---	---
Ro: Richfield-----	B		---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft	---	---	---	---
Rs: Richfield-----	B		---	---	---	---	---	---	---
Spearville-----	C		---	---	---	---	---	---	---
Sa: Satanta-----	B		---	---	---	---	---	---	---
SAP: Sand Pit-----	---		---	---	---	---	---	---	---
Sb: Satanta-----	B		---	---	---	---	---	---	---
Sp: Spearville-----	C		---	---	---	---	---	---	---
Sr: Spearville-----	C		---	---	---	---	---	---	---
Sw: Sweetwater-----	D		---	---	---	---	---	---	---
		January	0.5-3.0	>6.0	---	---	---	---	None
		February	0.5-3.0	>6.0	---	---	---	---	None
		March	0.5-3.0	>6.0	---	---	---	---	None
		April	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		May	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		June	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		July	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		August	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		September	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		October	0.5-3.0	>6.0	---	---	---	Brief	Occasional
		November	0.5-3.0	>6.0	---	---	---	---	None
		December	0.5-3.0	>6.0	---	---	---	---	None
Tf: Optima-----	A		---	---	---	---	---	---	---
Ua: Ulysses-----	B		---	---	---	---	---	---	---
Ub: Ulysses-----	B		---	---	---	---	---	---	---
UBB: Uly-----	B		---	---	---	---	---	---	---
Uc: Ulysses-----	B		---	---	---	---	---	---	---
UCC: Uly-----	B		---	---	---	---	---	---	---
Um: Ulysses-----	B		---	---	---	---	---	---	---
Colby-----	B		---	---	---	---	---	---	---
W: Water-----	---		---	---	---	---	---	---	---

The following table gives estimates of various soil features. The estimates are used in land use planning that involves engineering considerations.

A restrictive layer is a nearly continuous layer that has one or more physical, chemical, or thermal properties that significantly impede the movement of water and air through the soil or that restrict roots or otherwise provide an unfavorable root environment. Examples are bedrock, cemented layers, dense layers, and frozen layers. The table indicates the hardness and thickness of the restrictive layer, both of which significantly affect the ease of excavation. Depth to top is the vertical distance from the soil surface to the upper boundary of the restrictive layer.

Potential for frost action is the likelihood of upward or lateral expansion of the soil caused by the formation of segregated ice lenses (frost heave) and the subsequent collapse of the soil and loss of strength on thawing. Frost action occurs when moisture moves into the freezing zone of the soil. Temperature, texture, density, permeability, content of organic matter, and depth to the water table are the most important factors considered in evaluating the potential for frost action. It is assumed that the soil is not insulated by vegetation or snow and is not artificially drained. Silty and highly structured, clayey soils that have a high water table in winter are the most susceptible to frost action. Well drained, very gravelly, or very sandy soils are the least susceptible. Frost heave and low soil strength during thawing cause damage to pavements and other rigid structures.

Risk of corrosion pertains to potential soil-induced electrochemical or chemical action that corrodes or weakens uncoated steel or concrete. The rate of corrosion of uncoated steel is related to such factors as soil moisture, particle-size distribution, acidity, and electrical conductivity of the soil. The rate of corrosion of concrete is based mainly on the sulfate and sodium content, texture, moisture content, and acidity of the soil. Special site examination and design may be needed if the combination of factors results in a severe hazard of corrosion. The steel or concrete in installations that intersect soil boundaries or soil layers is more susceptible to corrosion than the steel or concrete in installations that are entirely within one kind of soil or within one soil layer.

For uncoated steel, the risk of corrosion, expressed as low, moderate, or high, is based on soil drainage class, total acidity, electrical resistivity near field capacity, and electrical conductivity of the saturation extract.

For concrete, the risk of corrosion also is expressed as low, moderate, or high. It is based on soil texture, acidity, and amount of sulfates in the saturation extract.

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness		Uncoated Steel	Concrete
		In	In				
055BP: Bridgeport, rarely flooded-	---	---	---	---	Moderate	Low	Low
055LD: Las, occasionally flooded-----	---	---	---	---	Moderate	High	Moderate
Las Animas, occasionally flooded-----	---	---	---	---	Moderate	High	Low
055TF: Valent-----	---	---	---	---	Low	Moderate	Low
055TV: Valent-----	---	---	---	---	Low	Moderate	Low
Vona-----	---	---	---	---	Low	High	Low
055VO: Vona-----	---	---	---	---	Low	High	Low
081LU: Lubbock-----	---	---	---	---	Low	High	Low
083UM: Uly-----	---	---	---	---	Moderate	Moderate	Low
Coly-----	---	---	---	---	Moderate	Moderate	Low
119HC: Harney, eroded--	---	---	---	---	Low	High	Low
119RX: Roxbury-----	---	---	---	---	Moderate	Low	Low
119UA: Uly-----	---	---	---	---	Moderate	High	Low
119UT: Uly-----	---	---	---	---	Moderate	High	Low
An: Bridgeport, channeled-----	---	---	---	---	Moderate	Low	Low
ARR: Arkansas River--	---	---	---	---	---	---	---
Bc: Bridgeport-----	---	---	---	---	Moderate	Low	Low
Bd: Bridgeport-----	---	---	---	---	Moderate	Low	Low
Bo: Optima-----	---	---	---	---	Low	Low	Low
Da: Dale-----	---	---	---	---	Moderate	Moderate	Low
Gr: Schamber-----	---	---	---	---	Low	Moderate	Low
GRP: Pits-----	---	---	---	---	Low	Low	Low
Ha: Harney-----	---	---	---	---	Low	High	Low
HB: Harney-----	---	---	---	---	Low	High	Low
INL: Aquolls-----	---	---	---	---	Low	---	---
La: Las Animas, occasionally flooded-----	---	---	---	---	Moderate	High	Moderate
Lesho, occasionally flooded-----	---	---	---	---	Moderate	High	Low
Lc: Las Animas, occasionally flooded-----	---	---	---	---	Moderate	High	Low
Ld: Las Animas-----	---	---	---	---	Moderate	High	Low
Le: Las, occasionally flooded-----	---	---	---	---	Moderate	High	Moderate
Lh: Lesho, occasionally flooded-----	---	---	---	---	Moderate	High	Low
Lk: Lesho-----	---	---	---	---	Moderate	High	Low
Sweetwater-----	---	---	---	---	Low	High	Low
Ll: Lincoln, occasionally flooded-----	---	---	---	---	Low	Low	Low
Lo: Lofton-----	---	---	---	---	Low	High	Low
Lu: Lubbock-----	---	---	---	---	Low	High	Low
Md: Penden-----	---	---	---	---	Moderate	Moderate	Low

SOIL FEATURES--Continued  
Gray County, Kansas

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness		Uncoated Steel	Concrete
		In	In				
Me: Penden-----	---	---	---	---	Moderate	Moderate	Low
Mf: Penden, Severely eroded-----	---	---	---	---	Moderate	Moderate	Low
Mh: Penden, Severely Eroded-----	---	---	---	---	Moderate	Moderate	Low
Mm: Manter-----	---	---	---	---	Moderate	High	Low
Mn: Manter-----	---	---	---	---	Moderate	High	Low
Mo: Manter-----	---	---	---	---	Moderate	High	Low
Mp: Campus-----	19-40	Bedrock (lithic)	---	Strongly cemented Indurated	Low	Low	Low
Canlon-----	8-20	Bedrock (lithic)	---		Low	Low	Low
Mu: Manter-----	---	---	---	---	Moderate	High	Low
Ulysses-----	---	---	---	---	Moderate	Moderate	Low
Pa: Pratt-----	---	---	---	---	Low	Low	Moderate
Pg: Pratt-----	---	---	---	---	Low	Low	Moderate
Pt: Pratt-----	---	---	---	---	Low	Low	Moderate
Optima-----	---	---	---	---	Low	Low	Low
Ra: Ness-----	---	---	---	---	Low	High	Low
Rm: Richfield-----	---	---	---	---	Low	High	Low
Rn: Richfield-----	---	---	---	---	Low	High	Low
Ro: Richfield-----	---	---	---	---	Low	High	Low
Rs: Richfield-----	---	---	---	---	Low	High	Low
Spearville-----	---	---	---	---	Low	High	Low
Sa: Satanta-----	---	---	---	---	Moderate	Low	Low
SAP: Sand Pit-----	---	---	---	---	---	---	---
Sb: Satanta-----	---	---	---	---	Moderate	Low	Low
Sp: Spearville-----	---	---	---	---	Low	High	Low
Sr: Spearville-----	---	---	---	---	Low	High	Low
Sw: Sweetwater-----	---	---	---	---	Low	High	Low
Tf: Optima-----	---	---	---	---	Low	Low	Low
Ua: Ulysses-----	---	---	---	---	Moderate	Moderate	Low
Ub: Ulysses-----	---	---	---	---	Moderate	Moderate	Low
UBB: Uly-----	---	---	---	---	Moderate	High	Low
Uc: Ulysses-----	---	---	---	---	Moderate	Moderate	Low
UCC: Uly-----	---	---	---	---	Moderate	High	Low
Um: Ulysses-----	---	---	---	---	Moderate	Moderate	Low
Colby-----	---	---	---	---	Low	Low	Low
W: Water-----	---	---	---	---	Low	---	---



WATER MANAGEMENT  
Gray County, Kansas

The soils of the survey area are rated in the Water Management table according to limitations that affect their suitability for water management. Soils are rated for pond reservoir areas, drainage, irrigation, terraces and diversions, and grassed waterways. Restrictive features that affect each soil for the specified use is also provided in the table.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the specified use. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Moderately limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Limited indicates that the soil has one or more features that are significant limitations for the specified use. The limitations can be overcome, but generally require special design, soil reclamation, or installation procedures that may result in additional expense. Fair performance and moderate to high maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Limitation class terms, such as very limited or limited, etc., limitation ratings, and numerical ratings are shown for each soil feature listed. As many as three soil features may be listed for each soil component if applicable. The overall limitation rating for the soil component is based on the most severe limitation.

Pond reservoir areas hold water behind a dam or embankment. Soils best suited to this use have low seepage potential in the upper 60 inches. The seepage potential is determined by the permeability of the soil and the depth to fractured bedrock or other permeable material. Excessive slope can affect the storage capacity of the reservoir area.

Embankments, dikes, and levees are raised structures of soil material, generally less than 20 feet high, constructed to impound water or to protect land against overflow. In this table, the soils are rated as a source of material for embankment fill. The ratings apply to the soil material below the surface layer to a depth of about 5 feet. It is assumed that soil layers will be uniformly mixed and compacted during construction.

The ratings do not indicate the ability of the natural soil to support an embankment. Soil properties to a depth even greater than the height of the embankment can affect performance and safety of the embankment. Generally, deeper onsite investigation is needed to determine these properties.

Soil material in embankments must be resistant to seepage, piping, and erosion and have favorable compaction characteristics. Unfavorable features include less than 5 feet of suitable material and a high content of stones or boulders, organic matter, or salts or sodium. A high water table affects the amount of usable material. It also affects traffic ability.

Aquifer-fed excavated ponds are pits or dugouts that extend to a ground-water aquifer or to a depth below a permanent water table. Excluded are ponds that are fed only by surface runoff and embankment ponds that impound water 3 feet or more above the original surface. Excavated ponds are affected by depth to a permanent water table, permeability of the aquifer, and quality of the water as inferred from the salinity of the soil. Depth to bedrock and the content of large stones affect the ease of excavation.

Drainage is the removal of excess surface and subsurface water from the soil. How easily and effectively the soil is drained depends on the depth to bedrock, to a cemented pan, or to other layers that affect the rate of water movement; permeability; depth to a high water table or depth of standing water if the soil is subject to ponding; slope; susceptibility to flooding; subsidence of organic layers; and the potential for frost action. Excavating and grading and the stability of ditch banks are affected by depth to bedrock or to a cemented pan, large stones, slope, and the hazard of cutbanks caving. The productivity of the soil after drainage is adversely affected by extreme acidity or by toxic substances in the root zone, such as salts, sodium, and sulfur. Availability of drainage outlets is not considered in the ratings.

Irrigation is the controlled application of water to supplement rainfall and support plant growth. The design and management of an irrigation system are affected by depth to the water table, the need for drainage, flooding, available water capacity, intake rate, permeability, erosion hazard, and slope. The construction of a system is affected by large stones and depth to bedrock or to a cemented pan. The performance of a system is affected by the depth of the root zone, the amount of salts or sodium, and soil reaction.

Terraces and diversions are embankments or a combination of channels and ridges constructed across a slope to control erosion and conserve moisture by intercepting runoff. Slope, wetness, large stones, and depth to bedrock or to a cemented pan affect the construction of terraces and diversions. A restricted rooting depth, a very limited hazard of wind erosion or water erosion, an excessively coarse texture, and restricted permeability adversely affect maintenance.

Grassed waterways are natural or constructed channels, generally broad and shallow, which conduct surface water to outlets at a non-erosive velocity. Large stones, wetness, slope, and depth to bedrock or to a cemented pan affect the construction of grassed waterways. A hazard of wind erosion, low available water capacity, restricted rooting depth, toxic substances such as salts and sodium, and restricted permeability adversely affect the growth and maintenance of the grass after construction.

WATER MANAGEMENT--Continued  
Gray County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
055BP: Bridgeport, rarely flooded--	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
055LD: Las, occasionally flooded-----	Limitation: flooding cutbanks cave	Limitation: flooding wetness	Limitation: too sandy wetness	Favorable
Las Animas, occasionally flooded-----	Limitation: flooding cutbanks cave	Limitation: flooding wetness soil blowing	Limitation: too sandy wetness soil blowing	Favorable
055TF: Valent-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope too arid droughty
055TV: Valent-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: too arid droughty
Vona-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: too arid droughty
055VO: Vona-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: too arid droughty
081LU: Lubbock-----	Limitation: deep to water	Favorable	Favorable	Favorable
083UM: Uly-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
Coly-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily	Limitation: erodes easily
119HC: Harney, eroded--	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
119RX: Roxbury-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
119UA: Uly-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
119UT: Uly-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
An: Bridgeport, channeled-----	Limitation: deep to water	Limitation: flooding	Limitation: erodes easily	Limitation: erodes easily
ARR: Arkansas River--	Limitation: flooding cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness	Limitation: wetness droughty
Bc: Bridgeport-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
Bd: Bridgeport-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
Bo: Optima-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: too arid droughty
Da: Dale-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
Gr: Schamber-----	Limitation: deep to water	Limitation: slope droughty	Limitation: slope too sandy	Limitation: slope too arid droughty

WATER MANAGEMENT--Continued  
Gray County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
GRP: Pits-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy	Limitation: rooting depth slope droughty
Ha: Harney-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
HB: Harney-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
INL: Aquolls-----	---	---	---	---
La: Las Animas, occasionally flooded-----	Limitation:  flooding	Limitation:  flooding wetness soil blowing	Limitation:  wetness soil blowing	Limitation:  excess salt
Lesho, occasionally flooded-----	Limitation:  excess salt flooding cutbanks cave	Limitation:  excess salt flooding wetness	Limitation:  too sandy wetness	Limitation:  excess salt
Lc: Las Animas, occasionally flooded-----	Limitation:  flooding cutbanks cave	Limitation:  fast intake wetness droughty	Limitation:  too sandy wetness soil blowing	Limitation:  droughty
Ld: Las Animas-----	Limitation: flooding cutbanks cave	Limitation: wetness soil blowing droughty	Limitation: too sandy wetness soil blowing	Limitation: droughty
Le: Las, occasionally flooded-----	Limitation:  flooding cutbanks cave	Limitation:  flooding wetness	Limitation:  too sandy wetness	Favorable
Lh: Lesho, occasionally flooded-----	Limitation:  flooding cutbanks cave	Limitation:  flooding wetness	Limitation:  too sandy wetness	Favorable
Lk: Lesho-----	Limitation: flooding cutbanks cave	Limitation: flooding wetness	Limitation: too sandy wetness	Favorable
Sweetwater-----	Limitation: flooding cutbanks cave	Limitation: flooding wetness	Limitation: too sandy wetness	Limitation: wetness
Ll: Lincoln, occasionally flooded-----	Limitation:  deep to water	Limitation:  flooding soil blowing droughty	Limitation:  too sandy soil blowing	Limitation:  droughty
Lo: Lofton-----	---	---	---	---
Lu: Lubbock-----	Limitation: deep to water	Favorable	Favorable	Favorable
Md: Penden-----	Limitation: deep to water	Limitation: slope	Favorable	Favorable
Me: Penden-----	Limitation: deep to water	Limitation: slope	Limitation: slope	Limitation: slope
Mf: Penden, Severely eroded-----	Limitation:  deep to water	Limitation:  slope	Limitation:  slope	Limitation:  slope
Mh: Penden, Severely Eroded-----	Limitation:  deep to water	Limitation:  slope	Limitation:  slope	Limitation:  slope

WATER MANAGEMENT--Continued  
Gray County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Mm: Manter-----	Limitation: deep to water	Limitation: soil blowing	Limitation: too sandy soil blowing	Limitation: too arid
Mn: Manter-----	Limitation: deep to water	Limitation: soil blowing	Limitation: too sandy soil blowing	Limitation: too arid
Mo: Manter-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: too sandy soil blowing	Limitation: too arid
Mp: Campus-----	Limitation: deep to water	Limitation: slope depth to rock	Limitation: depth to rock	Limitation: depth to rock
Canlon-----	Limitation: deep to water	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
Mu: Manter-----	Limitation: deep to water	Limitation: slope soil blowing	Limitation: too sandy soil blowing	Limitation: too arid
Ulysses-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily too arid
Pa: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
Pg: Pratt-----	Limitation: deep to water	Limitation: fast intake soil blowing droughty	Limitation: too sandy soil blowing	Limitation: droughty
Pt: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
Optima-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: too arid droughty
Ra: Ness-----	Limitation: percs slowly	Limitation: percs slowly slow intake wetness	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily percs slowly wetness
Rm: Richfield-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily too arid
Rn: Richfield-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily too arid
Ro: Richfield-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily too arid
Rs: Richfield-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily too arid
Spearville-----	Limitation: deep to water	Limitation: erodes easily percs slowly	Limitation: erodes easily	Limitation: erodes easily percs slowly
Sa: Satanta-----	Limitation: deep to water	Favorable	Favorable	Limitation: too arid
SAP: Sand Pit-----	---	---	---	---
Sb: Satanta-----	Limitation: deep to water	Favorable	Favorable	Limitation: too arid
Sp: Spearville-----	Limitation: deep to water	Limitation: percs slowly	Limitation: erodes easily	Limitation: erodes easily percs slowly
Sr: Spearville-----	Limitation: deep to water	Limitation: percs slowly	Limitation: erodes easily	Limitation: erodes easily percs slowly

WATER MANAGEMENT--Continued  
Gray County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Sw: Sweetwater-----	Limitation: flooding cutbanks cave	Limitation: flooding wetness	Limitation: too sandy wetness	Limitation: wetness
Tf: Optima-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: too arid droughty
Ua: Ulysses-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily too arid
Ub: Ulysses-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily too arid
UBB: Uly-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
Uc: Ulysses-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily too arid
UCC: Uly-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
Um: Ulysses-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily too arid
Colby-----	Limitation: deep to water	Limitation: erodes easily slope	Limitation: erodes easily	Limitation: erodes easily too arid
W: Water-----	---	---	---	---

WATER MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
055BP: Bridgeport, rarely flooded-----	100	Somewhat limited  Seepage	 0.70	Somewhat limited  Piping	 0.67	Very limited  Deep to water	 1.00
055LD: Las, occasionally flooded-----	55	Very limited  Seepage	 1.00	Somewhat limited  Seepage Depth to saturated zone	 0.93 0.86	Very limited  Cutbanks cave Deep to water	 1.00 0.06
Las Animas, occasionally flooded-----	45	Very limited  Seepage	 1.00	Very limited  Seepage Depth to saturated zone	 1.00 0.95	Very limited  Cutbanks cave Deep to water	 1.00 0.02
055TF: Valent-----	100	Very limited Seepage Slope	1.00 0.10	Very limited Seepage	1.00	Very limited Deep to water	1.00
055TV: Valent-----	50	Very limited Seepage	1.00	Somewhat limited Seepage	1.00	Very limited Deep to water	1.00
Vona-----	50	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
055VO: Vona-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
081LU: Lubbock-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
083UM: Uly-----	70	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
Coly-----	30	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
119HC: Harney, eroded-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.17	Very limited Deep to water	1.00
119RX: Roxbury-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.60	Very limited Deep to water	1.00
119UA: Uly-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
119UT: Uly-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
An: Bridgeport, channeled-----	100	Somewhat limited  Seepage	 0.70	Somewhat limited  Piping	 0.70	Very limited  Deep to water	 1.00
ARR: Arkansas River-----	100	Not rated		Not rated		Not rated	
Bc: Bridgeport-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.68	Very limited Deep to water	1.00
Bd: Bridgeport-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.68	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Bo: Optima-----	70	Very limited Seepage	1.00	Somewhat limited Seepage	1.00	Very limited Deep to water	1.00
Da: Dale-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.85	Very limited Deep to water	1.00
Gr: Schamber-----	100	Not rated		Not rated		Not rated	
GRP: Pits-----	100	Not rated		Not rated		Not rated	
Ha: Harney-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.10	Very limited Deep to water	1.00
HB: Harney-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.17	Very limited Deep to water	1.00
INL: Aquolls-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Somewhat limited Cutbanks cave	0.10
La: Las Animas, occasionally flooded-----	50	Very limited		Somewhat limited		Very limited	
		Seepage	1.00	Seepage Depth to saturated zone	0.96 0.43	Cutbanks cave Deep to water	1.00 0.25
						Salty water	0.06
Lesho, occasionally flooded-----	40	Very limited		Somewhat limited		Very limited	
		Seepage	1.00	Seepage Depth to saturated zone Salinity	0.93 0.43 0.12	Cutbanks cave Salty water Deep to water	1.00 0.50 0.25
Lc: Las Animas, occasionally flooded-----	100	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage Depth to saturated zone	1.00 0.95	Cutbanks cave Deep to water	1.00 0.02
Ld: Las Animas-----	100	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Deep to water	1.00 0.02
Le: Las, occasionally flooded-----	100	Very limited		Somewhat limited		Very limited	
		Seepage	1.00	Seepage Depth to saturated zone	0.93 0.86	Cutbanks cave Deep to water	1.00 0.06
Lh: Lesho, occasionally flooded-----	100	Very limited		Somewhat limited		Very limited	
		Seepage	1.00	Seepage Depth to saturated zone	0.96 0.43	Cutbanks cave Deep to water	1.00 0.25
Lk: Lesho-----	50	Very limited Seepage	1.00	Somewhat limited Seepage	0.96	Very limited Cutbanks cave	1.00

WATER MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Sweetwater-----	25	Very limited Seepage	1.00	Depth to saturated zone  Very limited Depth to saturated zone Seepage	0.43  1.00 0.87	Deep to water  Very limited Cutbanks cave	0.25  1.00
Ll: Lincoln, occasionally flooded-----	100	Very limited  Seepage	  1.00	Somewhat limited  Seepage	  0.09	Very limited  Deep to water	  1.00
Lo: Lofton-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
Lu: Lubbock-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.04	Very limited Deep to water	1.00
Md: Penden-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.08	Very limited Deep to water	1.00
Me: Penden-----	100	Somewhat limited Seepage Slope	0.70 0.00	Somewhat limited Piping	0.08	Very limited Deep to water	1.00
Mf: Penden, Severely eroded-----	100	Somewhat limited  Seepage	  0.70	Somewhat limited  Piping	  0.08	Very limited  Deep to water	  1.00
Mh: Penden, Severely Eroded-----	100	Somewhat limited  Seepage Slope	  0.70 0.00	Somewhat limited  Piping	  0.08	Very limited  Deep to water	  1.00
Mm: Manter-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.08	Very limited Deep to water	1.00
Mn: Manter-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
Mo: Manter-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
Mp: Campus-----	75	Very limited Seepage Depth to bedrock	1.00 1.00	Very limited Thin layer Piping	1.00 0.68	Very limited Deep to water	1.00
Canlon-----	25	Very limited Seepage Depth to bedrock	1.00 1.00	Very limited Thin layer	1.00	Very limited Deep to water	1.00
Mu: Manter-----	55	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Deep to water	1.00
Ulysses-----	45	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
Pa: Pratt-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
Pg: Pratt-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
Pt: Pratt-----	60	Very limited		Somewhat limited		Very limited	



WATER MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Optima-----	40	Seepage	1.00	Seepage	0.57	Deep to water	1.00
Ra: Ness-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	1.00	Very limited Deep to water	1.00
Rm: Richfield-----	100	Somewhat limited Seepage	0.43	Very limited Ponding Depth to saturated zone	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.57 0.10
Rn: Richfield-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.57	Very limited Deep to water	1.00
Ro: Richfield-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.46	Very limited Deep to water	1.00
Rs: Richfield-----	60	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.46	Very limited Deep to water	1.00
Sa: Satanta-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.82	Very limited Deep to water	1.00
SAP: Sand Pit-----	100	Not rated		Not rated		Not rated	
Sb: Satanta-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.82	Very limited Deep to water	1.00
Sp: Spearville-----	100	Somewhat limited Seepage	0.05	Not limited		Very limited Deep to water	1.00
Sr: Spearville-----	100	Somewhat limited Seepage	0.05	Not limited		Very limited Deep to water	1.00
Sw: Sweetwater-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.08	Somewhat limited Cutbanks cave	0.10
Tf: Optima-----	90	Very limited Seepage Slope	1.00 0.00	Somewhat limited Seepage	1.00	Very limited Deep to water	1.00
Ua: Ulysses-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
Ub: Ulysses-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
UBB: Uly-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
Uc: Ulysses-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
UCC: Uly-----	100	Somewhat limited		Very limited		Very limited	

WATER MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Um: Ulysses-----	55	Seepage	0.70	Piping	1.00	Deep to water	1.00
		Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
	45	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
W: Water-----	100	Not rated		Not rated		Not rated	

SANITARY FACILITIES  
Gray County, Kansas

Sanitary Facilities

The following tables show the degree and kind of soil limitations that affect septic tank absorption fields, sewage lagoons, sanitary landfills, and daily cover for landfill. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Septic tank absorption fields are areas in which effluent from a septic tank is distributed into the soil through subsurface tiles or perforated pipe. Only that part of the soil between depths of 24 and 60 inches is evaluated. The ratings are based on the soil properties that affect absorption of the effluent, construction and maintenance of the system, and public health. Permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, and flooding affect absorption of the effluent. Stones and boulders, ice, and bedrock or a cemented pan interfere with installation. Subsidence interferes with installation and maintenance. Excessive slope may cause lateral seepage and surfacing of the effluent in downslope areas.

Some soils are underlain by loose sand and gravel or fractured bedrock at a depth of less than 4 feet below the distribution lines. In these soils the absorption field may not adequately filter the effluent, particularly when the system is new. As a result, the ground water may become contaminated.

Sewage lagoons are shallow ponds constructed to hold sewage while aerobic bacteria decompose the solid and liquid wastes. Lagoons should have a nearly level floor surrounded by cut slopes or embankments of compacted soil. Nearly impervious soil material for the lagoon floor and sides is required to minimize seepage and contamination of ground water. Considered in the ratings are slope, permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, flooding, large stones, and content of organic matter.

Soil permeability is a critical property affecting the suitability for sewage lagoons. Most porous soils eventually become sealed when they are used as sites for sewage lagoons. Until sealing occurs, however, the hazard of pollution is severe. Soils that have a permeability rate of more than 2 inches per hour are too porous for the proper functioning of sewage lagoons. In these soils, seepage of the effluent can result in contamination of the ground water. Ground-water contamination is also a hazard if fractured bedrock is within a depth of 40 inches, if the water table is high enough to raise the level of sewage in the lagoon, or if floodwater overtops the lagoon.

A high content of organic matter is detrimental to proper functioning of the lagoon because it inhibits aerobic activity. Slope, bedrock, and cemented pans can cause construction problems, and large stones can hinder compaction of the lagoon floor. If the lagoon is to be uniformly deep throughout, the slope must be gentle enough and the soil material must be thick enough over bedrock or a cemented pan to make land smoothing practical.

A trench sanitary landfill is an area where solid waste is placed in successive layers in an excavated trench. The waste is spread, compacted, and covered daily with a thin layer of soil excavated at the site. When the trench is full, a final cover of soil material at least 2 feet thick is placed over the landfill. The ratings in the table are based on the soil properties that affect the risk of pollution, the ease of excavation, trafficability, and revegetation. These properties include permeability, depth to bedrock or a cemented pan, depth to a water table, ponding, slope, flooding, texture, stones and boulders, highly organic layers, soil reaction, and content of salts and sodium. Unless otherwise stated, the ratings apply only to that part of the soil within a depth of about 6 feet. For deeper trenches, onsite investigation may be needed.

Hard, nonrippable bedrock, creviced bedrock, or highly permeable strata in or directly below the proposed trench bottom can affect the ease of excavation and the hazard of ground-water pollution. Slope affects construction of the trenches and the movement of surface water around the landfill. It also affects the construction and performance of roads in areas of the landfill.

Soil texture and consistence affect the ease with which the trench is dug and the ease with which the soil can be used as daily or final cover. They determine the workability of the soil when dry and when wet. Soils that are plastic and sticky when wet are difficult to excavate, grade, or compact and are difficult to place as a uniformly thick cover over a layer of refuse.

The soil material used as the final cover for a trench landfill should be suitable for plants. It should not have excess sodium or salts and should not be too acid. The surface layer generally has the best workability, the highest content of organic matter, and the best potential for plants. Material from the surface layer should be stockpiled for use as the final cover.

SANITARY FACILITIES  
Gray County, Kansas

In an area sanitary landfill, solid waste is placed in successive layers on the surface of the soil. The waste is spread, compacted, and covered daily with a thin layer of soil from a source away from the site. A final cover of soil material at least 2 feet thick is placed over the completed landfill. The ratings in the table are based on the soil properties that affect trafficability and the risk of pollution. These properties include flooding, permeability, depth to a water table, ponding, slope, and depth to bedrock or a cemented pan.

Flooding is a serious problem because it can result in pollution in areas downstream from the landfill. If permeability is too rapid or if fractured bedrock, a fractured cemented pan, or the water table is close to the surface, the leachate can contaminate the water supply. Slope is a consideration because of the extra grading required to maintain roads in the steeper areas of the landfill. Also, leachate may flow along the surface of the soils in the steeper areas and cause difficult seepage problems.

Daily cover for landfill is the soil material that is used to cover compacted solid waste in an area sanitary landfill. The soil material is obtained offsite, transported to the landfill, and spread over the waste. The ratings in the table also apply to the final cover for a landfill. They are based on the soil properties that affect workability, the ease of digging, and the ease of moving and spreading the material over the refuse daily during wet and dry periods. These properties include soil texture, depth to a water table, ponding, rock fragments, slope, depth to bedrock or a cemented pan, reaction, and content of salts, sodium, or lime.

Loamy or silty soils that are free of large stones and excess gravel are the best cover for a landfill. Clayey soils may be sticky and difficult to spread; sandy soils are subject to wind erosion.

Slope affects the ease of excavation and of moving the cover material. Also, it can influence runoff, erosion, and reclamation of the borrow area.

After soil material has been removed, the soil material remaining in the borrow area must be thick enough over bedrock, a cemented pan, or the water table to permit revegetation. The soil material used as the final cover for a landfill should be suitable for plants. It should not have excess sodium, salts, or lime and should not be too acid.

SANITARY FACILITIES--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
055BP: Bridgeport, rarely flooded-----	100	Somewhat limited		Somewhat limited	
		Restricted permeability	0.50	Seepage	0.50
		Flooding	0.40	Flooding	0.40
055LD: Las, occasionally flooded-----	55	Very limited		Very limited	
		Flooding	1.00	Flooding	1.00
		Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
		Restricted permeability	1.00		
Las Animas, occasionally flooded-----	45	Very limited		Very limited	
		Flooding	1.00	Flooding	1.00
		Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
055TF: Valent-----	100	Very limited		Very limited	
		Slope	1.00	Slope	1.00
		Filtering capacity	1.00	Seepage	1.00
055TV: Valent-----	50	Very limited		Very limited	
		Filtering capacity	1.00	Seepage	1.00
Vona-----	50	Very limited		Slope	1.00
		Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.91
055VO: Vona-----	100	Very limited		Very limited	
		Filtering capacity	1.00	Seepage	1.00
				Slope	0.09
081LU: Lubbock-----	100	Very limited		Somewhat limited	
		Restricted permeability	1.00	Seepage	0.50
083UM: Uly-----	70	Somewhat limited		Somewhat limited	
		Restricted permeability	0.50	Slope	0.67
				Seepage	0.50
Coly-----	30	Somewhat limited		Somewhat limited	
		Restricted permeability	0.50	Slope	0.67
				Seepage	0.50
119HC: Harney, eroded-----	100	Somewhat limited		Somewhat limited	
		Restricted permeability	0.50	Seepage	0.50
				Slope	0.00
119RX: Roxbury-----	100	Somewhat limited		Somewhat limited	
		Restricted permeability	0.50	Seepage	0.50
		Flooding	0.40	Flooding	0.40
119UA: Uly-----	100	Somewhat limited		Somewhat limited	
		Restricted permeability	0.50	Seepage	0.50
119UT: Uly-----	100	Somewhat limited		Somewhat limited	
		Restricted permeability	0.50	Slope	0.67
				Seepage	0.50
An: Bridgeport, channeled-----	100	Very limited		Very limited	
		Flooding	1.00	Flooding	1.00

SANITARY FACILITIES--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
ARR: Arkansas River-----	100	Restricted permeability	0.50	Seepage	0.50
		Not rated		Not rated	
Bc: Bridgeport-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
		Flooding	0.40	Flooding	0.40
Bd: Bridgeport-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
		Flooding	0.40	Flooding	0.40
Bo: Optima-----	70	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
				Slope	0.33
Da: Dale-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
		Flooding	0.40	Flooding	0.40
Gr: Schamber-----	100	Not rated		Not rated	
GRP: Pits-----	100	Not rated		Not rated	
Ha: Harney-----	100	Very limited Restricted permeability	1.00	Somewhat limited Seepage	0.50
HB: Harney-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
				Slope	0.00
INL: Aguolls-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Ponding	1.00	Ponding	1.00
La: Las Animas, occasionally flooded-----	50	Very limited		Very limited	
		Flooding	1.00	Flooding	1.00
		Depth to saturated zone	1.00	Seepage	1.00
				Depth to saturated zone	1.00
Lesho, occasionally flooded-----	40	Very limited		Very limited	
		Flooding	1.00	Flooding	1.00
		Filtering capacity	1.00	Seepage	1.00
		Restricted permeability	1.00	Depth to saturated zone	1.00
		Depth to saturated zone	1.00		
Lc: Las Animas, occasionally flooded-----	100	Very limited		Very limited	
		Flooding	1.00	Flooding	1.00
		Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
Ld: Las Animas-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00

SANITARY FACILITIES--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Le: Las, occasionally flooded-----	100	Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00
Lh: Lesho, occasionally flooded-----	100	Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
		Restricted permeability	1.00		
		Very limited		Very limited	
Lk: Lesho-----	50	Flooding	1.00	Flooding	1.00
		Restricted permeability	1.00	Seepage	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00		
Sweetwater-----	25	Very limited		Very limited	
		Flooding	1.00	Flooding	1.00
		Restricted permeability	1.00	Seepage	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00
Ll: Lincoln, occasionally flooded-----	100	Filtering capacity	1.00	Very limited	
		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00
		Filtering capacity	1.00	Seepage	1.00
Lo: Lofton-----	100	Depth to saturated zone	0.08	Slope	0.00
		Very limited			
Lu: Lubbock-----	100	Restricted permeability	1.00	Somewhat limited Seepage	0.50
		Very limited			
Md: Penden-----	100	Restricted permeability	1.00	Somewhat limited Seepage	0.50
		Somewhat limited			
Me: Penden-----	100	Restricted permeability	0.50	Somewhat limited Slope	0.67
		Slope	0.37	Seepage	0.50
Mf: Penden, Severely eroded-----	100	Somewhat limited		Very limited Slope	1.00
		Restricted permeability	0.50	Seepage	0.50
Mh: Penden, Severely Eroded-----	100	Somewhat limited		Slope	0.33
		Restricted permeability	0.50	Very limited	
Mm: Manter-----	100	Slope	0.37	Slope	1.00
		Not limited		Seepage	0.50
				Very limited	

SANITARY FACILITIES--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Mn: Manter-----	100	Not limited		Seepage	1.00
Mo: Manter-----	100	Not limited		Very limited Seepage Slope	1.00 0.00
Mp: Campus-----	75	Very limited Depth to bedrock	1.00	Very limited Seepage Slope	1.00 0.33
Canlon-----	25	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.50 1.00
Mu: Manter-----	55	Not limited		Very limited Seepage Slope	1.00 0.33
Ulysses-----	45	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.33
Pa: Pratt-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.33
Pg: Pratt-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
Pt: Pratt-----	60	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.91
Optima-----	40	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 1.00
Ra: Ness-----	100	Very limited Restricted permeability Ponding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Seepage	1.00 1.00 0.18
Rm: Richfield-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Rn: Richfield-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.00
Ro: Richfield-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.00
Rs: Richfield-----	60	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Spearville-----	40	Very limited Restricted permeability	1.00	Not limited	
Sa: Satanta-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
SAP: Sand Pit-----	100	Not rated		Not rated	



SANITARY FACILITIES--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Sb: Satanta-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
Sp: Spearville-----	100	Very limited Restricted permeability	1.00	Slope	0.00
Sr: Spearville-----	100	Very limited Restricted permeability	1.00	Not limited	
Sw: Sweetwater-----	100	Somewhat limited Restricted permeability	1.00	Somewhat limited Slope	0.00
Tf: Optima-----	90	Very limited Flooding	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Seepage	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00
Ua: Ulysses-----	100	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
Ub: Ulysses-----	100	Slope	0.63	Slope	1.00
UBB: Uly-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
		Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
		Somewhat limited Restricted permeability	0.50	Slope	0.00
Uc: Ulysses-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
UCC: Uly-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope	0.67
		Somewhat limited Restricted permeability	0.50	Seepage	0.50
		Somewhat limited Restricted permeability	0.50	Somewhat limited Slope	0.67
Um: Ulysses-----	55	Somewhat limited Restricted permeability	0.50	Seepage	0.50
Colby-----	45	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope	0.67
		Somewhat limited Restricted permeability	0.50	Seepage	0.50
		Somewhat limited Restricted permeability	0.50	Somewhat limited Slope	0.67
W: Water-----	100	Not rated		Not rated	

SANITARY FACILITIES--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
055BP: Bridgeport, rarely flooded-----	100	Somewhat limited		Somewhat limited		Not limited	
		Flooding	0.40	Flooding	0.40		
055LD: Las, occasionally flooded-----	55	Very limited		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
		Seepage	1.00	Seepage	1.00	Depth to saturated zone	0.47
		Too Sandy	1.00				
Las Animas, occasionally flooded-----	45	Very limited		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
		Seepage	1.00	Seepage	1.00	Depth to saturated zone	0.68
		Too Sandy	1.00				
055TF: Valent-----	100	Very limited		Very limited		Very limited	
		Too Sandy	1.00	Slope	1.00	Too Sandy	1.00
		Slope	1.00			Seepage	1.00
						Slope	1.00
055TV: Valent-----	50	Very limited		Not limited		Very limited	
		Too Sandy	1.00			Too Sandy	1.00
						Seepage	1.00
Vona-----	50	Not limited		Not limited		Very limited	1.00
						Seepage	1.00
055VO: Vona-----	100	Not limited		Not limited		Very limited	
						Seepage	1.00
081LU: Lubbock-----	100	Somewhat limited		Not limited		Very limited	
		Too clayey	0.50			Hard to compact	1.00
						Too clayey	0.50
083UM: Uly-----	70	Not limited		Not limited		Not limited	
Coly-----	30	Not limited		Not limited		Not limited	
119HC: Harney, eroded-----	100	Not limited		Not limited		Not limited	
119RX: Roxbury-----	100	Somewhat limited		Somewhat limited		Somewhat limited	
		Too clayey	0.50	Flooding	0.40	Too clayey	0.50
		Flooding	0.40				
119UA: Uly-----	100	Not limited		Not limited		Not limited	
119UT: Uly-----	100	Not limited		Not limited		Not limited	
An: Bridgeport, channeled-----	100	Very limited		Very limited		Not limited	
		Flooding	1.00	Flooding	1.00		
ARR: Arkansas River-----	100	Not rated		Not rated		Not rated	
Bc: Bridgeport-----	100	Somewhat limited		Somewhat limited		Somewhat limited	
		Too clayey	0.50	Flooding	0.40	Too clayey	0.50
		Flooding	0.40				
Bd: Bridgeport-----	100	Somewhat limited		Somewhat limited		Somewhat limited	
		Too clayey	0.50	Flooding	0.40	Too clayey	0.50
		Flooding	0.40				
Bo: Optima-----	70	Very limited		Not limited		Very limited	
		Too Sandy	1.00			Too Sandy	1.00
						Seepage	1.00
Da: Dale-----	100	Somewhat limited		Somewhat limited		Not limited	
		Flooding	0.40	Flooding	0.40		
Gr: Schamber-----	100	Not rated		Not rated		Not rated	

SANITARY FACILITIES--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
GRP: Pits-----	100	Not rated		Not rated		Not rated	
Ha: Harney-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
HB: Harney-----	100	Not limited		Not limited		Not limited	
INL: Aquolls-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Ponding	1.00	Ponding	1.00	Ponding	1.00
		Seepage	1.00				
La: Las Animas, occasionally flooded-----	50	Very limited		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to	1.00	Depth to	1.00	Seepage	0.50
		saturated zone	1.00	saturated zone	1.00	Depth to	0.09
		Seepage		Seepage		saturated zone	
Lesho, occasionally flooded-----	40	Too Sandy	1.00	Very limited		Very limited	
		Very limited					
		Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to	1.00	Depth to	1.00	Seepage	1.00
		saturated zone	1.00	saturated zone	1.00	Depth to	0.09
		Seepage		Seepage		saturated zone	
		Too Sandy	1.00				
Lc: Las Animas, occasionally flooded-----	100	Very limited		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to	1.00	Depth to	1.00	Seepage	1.00
		saturated zone	1.00	saturated zone	1.00	Depth to	0.68
		Seepage		Seepage		saturated zone	
		Too Sandy	1.00				
Ld: Las Animas-----	100	Very limited		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to	1.00	Depth to	1.00	Seepage	1.00
		saturated zone	1.00	saturated zone	1.00	Depth to	0.68
		Seepage		Seepage		saturated zone	
		Too Sandy	1.00				
Le: Las, occasionally flooded-----	100	Very limited		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to	1.00	Depth to	1.00	Seepage	1.00
		saturated zone	1.00	saturated zone	1.00	Depth to	0.47
		Seepage		Seepage		saturated zone	
		Too Sandy	1.00				
Lh: Lesho, occasionally flooded-----	100	Very limited		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to	1.00	Depth to	1.00	Seepage	1.00
		saturated zone	1.00	saturated zone	1.00	Depth to	0.09
		Seepage		Seepage		saturated zone	
		Too Sandy	1.00				
Lk: Lesho-----	50	Very limited		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to	1.00	Depth to	1.00	Seepage	1.00
		saturated zone	1.00	saturated zone	1.00	Depth to	0.09
		Seepage		Seepage		saturated zone	
		Too Sandy	1.00				
Sweetwater-----	25	Very limited		Very limited		Very limited	

SANITARY FACILITIES--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ll: Lincoln, occasionally flooded-----	100	Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
		Seepage	1.00	Seepage	1.00	Depth to saturated zone	0.96
		Too Sandy	1.00				
Lo: Lofton-----	100	Very limited		Very limited		Very limited	
		Flooding	1.00	Flooding	1.00	Too Sandy	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Seepage	1.00
		Seepage	1.00	Seepage	1.00		
Lu: Lubbock-----	100	Too Sandy	1.00				
		Very limited	1.00	Not limited		Very limited Too clayey Hard to compact	1.00 1.00
Md: Penden-----	100	Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
		Somewhat limited	0.50	Not limited		Somewhat limited Too clayey	0.50
Me: Penden-----	100	Too clayey	0.50	Somewhat limited	0.37	Somewhat limited Too clayey Slope	0.50 0.37
		Slope	0.37	Slope	0.37		
Mf: Penden, Severely eroded-----	100	Somewhat limited		Not limited		Somewhat limited	
		Too clayey	0.50			Too clayey	0.50
Mh: Penden, Severely Eroded-----	100	Somewhat limited		Somewhat limited		Somewhat limited	
		Too clayey	0.50	Slope	0.37	Too clayey Slope	0.50 0.37
Mm: Manter-----	100	Slope	0.37	Not limited		Somewhat limited Seepage	0.50
		Not limited		Not limited		Somewhat limited Seepage	0.50
Mn: Manter-----	100	Not limited		Not limited		Somewhat limited Seepage	0.50
		Not limited		Not limited		Somewhat limited Seepage	0.50
Mo: Manter-----	100	Not limited		Not limited		Somewhat limited Seepage	0.50
		Not limited		Not limited		Somewhat limited Seepage	0.50
Mp: Campus-----	75	Very limited		Very limited		Very limited	
		Depth to bedrock	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
Canlon-----	25	Very limited		Very limited		Very limited	
		Depth to bedrock	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
Mu: Manter-----	55	Slope	0.04	Slope	0.04	Slope	0.04
		Not limited		Not limited		Somewhat limited Seepage	0.50
Ulysses-----	45	Not limited		Not limited		Not limited	
		Not limited		Not limited		Not limited	
Pa: Pratt-----	100	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Seepage	1.00
Pg: Pratt-----	100	Too Sandy	1.00			Too Sandy	0.50
		Very limited		Very limited		Very limited	
Pt: Pratt-----	60	Seepage	1.00	Seepage	1.00	Seepage	1.00
		Too Sandy	1.00			Too Sandy	0.50
Optima-----	40	Very limited		Very limited		Very limited	
		Seepage	1.00	Seepage	1.00	Seepage	1.00
Ra: Ness-----	100	Too Sandy	1.00	Not limited		Too Sandy	0.50
		Very limited	1.00			Very limited Too Sandy Seepage	1.00 1.00
Rm: Richfield-----	100	Depth to saturated zone	1.00	Very limited Ponding	1.00	Very limited Ponding	1.00
		Ponding	1.00	Depth to saturated zone	1.00	Depth to saturated zone Hard to compact	1.00 1.00
Richfield-----	100	Not limited		Not limited		Not limited	
		Not limited		Not limited		Not limited	

SANITARY FACILITIES--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Rn: Richfield-----	100	Not limited		Not limited		Not limited	
Ro: Richfield-----	100	Not limited		Not limited		Not limited	
Rs: Richfield-----	60	Not limited		Not limited		Not limited	
Spearville-----	40	Not limited		Not limited		Not limited	
Sa: Satanta-----	100	Not limited		Not limited		Not limited	
SAP: Sand Pit-----	100	Not rated		Not rated		Not rated	
Sb: Satanta-----	100	Not limited		Not limited		Not limited	
Sp: Spearville-----	100	Not limited		Not limited		Not limited	
Sr: Spearville-----	100	Not limited		Not limited		Not limited	
Sw: Sweetwater-----	100	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 0.96
Tf: Optima-----	90	Very limited Too Sandy Slope	1.00 0.63	Somewhat limited Slope	0.63	Very limited Too Sandy Seepage Slope	1.00 1.00 0.63
Ua: Ulysses-----	100	Not limited		Not limited		Not limited	
Ub: Ulysses-----	100	Not limited		Not limited		Not limited	
UBB: Uly-----	100	Not limited		Not limited		Not limited	
Uc: Ulysses-----	100	Not limited		Not limited		Not limited	
UCC: Uly-----	100	Not limited		Not limited		Not limited	
Um: Ulysses-----	55	Not limited		Not limited		Not limited	
Colby-----	45	Not limited		Not limited		Not limited	
W: Water-----	100	Not rated		Not rated		Not rated	

AGRICULTURAL WASTE MANAGEMENT  
Gray County, Kansas

The nature of the soil is also important in the application of organic wastes and wastewater to land as fertilizers and irrigation; it is also important when the soil is used as a medium for treatment and disposal of these wastes. Favorable soil properties are required to prevent environmental damage.

The use of organic wastes and wastewater as production resources will result in energy conservation, prevent the waste of these important resources, and prevent problems associated with their disposal. Where disposal is the goal, and a maximum amount is disposed in a minimum area to hold costs to a minimum, risk of environmental damage is the principal constraint. Where the reuse goal is pursued, and a minimum amount is applied to a maximum area to obtain the greatest benefit, environmental damage is unlikely.

Interpretations developed for waste management may include ratings for (1) manure and food processing wastes; (2) municipal sewage sludge; (3) irrigation use of wastewater; or (4) treatment of wastewater by the slow rate process, overland flow process, or rapid infiltration process. If available, these should be located in this subsection.

Soil properties are important considerations in areas where soils are used as sites for the treatment and disposal of organic waste and wastewater. Selection of soils with properties that favor waste management can help to prevent environmental damage.

The Ag-Waste tables show the degree and kind of soil limitations affecting the treatment of agricultural waste, including municipal and food-processing wastewater and effluent from lagoons or storage ponds. Municipal wastewater is the waste stream from a municipality. It contains domestic waste and may contain industrial waste. It may have received primary or secondary treatment. It is rarely untreated sewage. Food-processing wastewater results from the preparation of fruits, vegetables, milk, cheese, and meats for public consumption. In places it is high in content of sodium and chloride. In the context of these tables, the effluent in lagoons and storage ponds is from facilities used to treat or store food-processing wastewater or domestic or animal waste. Domestic and food-processing wastewater is very dilute, and the effluent from the facilities that treat or store it commonly is very low in content of carbonaceous and nitrogenous material; the content of nitrogen commonly ranges from 10 to 30 milligrams per liter. The wastewater from animal waste treatment lagoons or storage ponds, however, has much higher concentrations of these materials, mainly because the manure has not been diluted as much as the domestic waste. The content of nitrogen in this wastewater generally ranges from 50 to 2,000 milligrams per liter. When wastewater is applied, checks should be made to ensure that nitrogen, phosphorus, heavy metals, and salts are not added in excessive amounts.

The ratings in the tables are for waste management systems that not only dispose of and treat organic waste or wastewater but also are beneficial to crops (application of manure and food-processing waste, application of sewage sludge, and disposal of wastewater by irrigation) and for waste management systems that are designed only for the purpose of wastewater disposal and treatment (overland flow of wastewater, rapid infiltration of wastewater, and slow rate treatment of wastewater).

The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect agricultural waste management. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are generally favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Application of manure and food-processing waste not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. Manure is the excrement of livestock and poultry, and food-processing waste is damaged fruit and vegetables and the peelings, stems, leaves, pits, and soil particles removed in food preparation. The manure and food-processing waste are either solid, slurry, or liquid. Their nitrogen content varies. A high content of nitrogen limits the application rate. Toxic or otherwise dangerous wastes, such as those mixed with the lye used in food processing, are not considered in the ratings.

The ratings are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the waste is applied, and the method by which the waste is applied. The properties that affect absorption include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, and available water capacity. The properties that affect plant growth and microbial activity include reaction, the sodium adsorption ratio, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of waste. Permanently frozen soils are unsuitable for waste treatment.

Application of sewage sludge not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. In the context of this table, sewage sludge is the residual product of the treatment of municipal sewage. The solid component consists mainly of cell mass, primarily bacteria cells that developed during secondary treatment and have incorporated soluble organics into their own bodies. The sludge has small amounts of sand, silt, and other solid debris. The content of nitrogen varies. Some sludge has constituents that are toxic to plants or hazardous to the food chain, such as heavy metals and exotic organic compounds, and should be analyzed chemically prior to use.

AGRICULTURAL WASTE MANAGEMENT  
Gray County, Kansas

The content of water in the sludge ranges from about 98 percent to less than 40 percent. The sludge is considered liquid if it is more than about 90 percent water, slurry if it is about 50 to 90 percent water, and solid if it is less than about 50 percent water.

The ratings in the table are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the sludge is applied, and the method by which the sludge is applied. The properties that affect absorption, plant growth, and microbial activity include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, available water capacity, reaction, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of sludge. Permanently frozen soils are unsuitable for waste treatment.

Disposal of wastewater by irrigation not only disposes of municipal wastewater and wastewater from food-processing plants, lagoons, and storage ponds but also can improve crop production by increasing the amount of water available to crops. The ratings in the table are based on the soil properties that affect the design, construction, management, and performance of the irrigation system. The properties that affect design and management include the sodium adsorption ratio, depth to a water table, ponding, available water capacity, permeability, slope, and flooding. The properties that affect construction include stones, cobbles, depth to bedrock or a cemented pan, depth to a water table, and ponding.

The properties that affect performance include depth to bedrock or a cemented pan, bulk density, the sodium adsorption ratio, salinity, reaction, and the cation-exchange capacity, which is used to estimate the capacity of a soil to adsorb heavy metals. Permanently frozen soils are not suitable for disposal of wastewater by irrigation.

See the National Soil Handbook, September 1992, Part 620, for criteria used in rating soils for sanitary facilities and waste management.

AGRICULTURAL WASTE MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
055BP: Bridgeport, rarely flooded-----	100	Not limited		Somewhat limited Flooding	0.40	Not limited	
055LD: Las, occasionally flooded-----	55	Very limited Filtering capacity 1.00 Depth to 0.86 saturated zone Flooding 0.60 Restricted 0.30 permeability		Very limited Filtering capacity 1.00 Flooding 1.00 Depth to 0.86 saturated zone Restricted 0.22 permeability		Very limited Filtering capacity 1.00 Depth to 0.86 saturated zone Flooding 0.60 Restricted 0.22 permeability	
Las Animas, occasionally flooded-----	45	Very limited Filtering capacity 1.00 Depth to 0.95 saturated zone Flooding 0.60		Very limited Filtering capacity 1.00 Flooding 1.00 Depth to 0.95 saturated zone		Very limited Filtering capacity 1.00 Depth to 0.95 saturated zone Flooding 0.60	
055TF: Valent-----	100	Very limited Slope 1.00 Filtering capacity 1.00 Droughty 0.75 Leaching limitation 0.45		Very limited Slope 1.00 Filtering capacity 1.00 Droughty 0.75		Very limited Too steep for surface application 1.00 Too steep for sprinkler application 1.00 Filtering capacity 1.00 Droughty 0.75	
055TV: Valent-----	50	Very limited Filtering capacity 1.00 Leaching limitation 0.45 Droughty 0.23		Very limited Filtering capacity 1.00 Droughty 0.23		Very limited Filtering capacity 1.00 Too steep for surface application 0.91 Droughty 0.23 Too steep for sprinkler application 0.02	
Vona-----	50	Very limited Filtering capacity 1.00		Very limited Filtering capacity 1.00		Very limited Filtering capacity 1.00 Too steep for surface application 0.66 Too steep for sprinkler application 0.00	
055VO: Vona-----	100	Very limited Filtering capacity 1.00		Very limited Filtering capacity 1.00		Very limited Filtering capacity 1.00 Too steep for surface application 0.00	
081LU: Lubbock-----	100	Somewhat limited Restricted permeability 0.30		Somewhat limited Restricted permeability 0.22		Somewhat limited Restricted permeability 0.22	
083UM: Uly-----	70	Not limited		Not limited		Somewhat limited Too steep for surface application 0.31	
Coly-----	30	Not limited		Not limited		Somewhat limited Too steep for surface application 0.31	
119HC: Harney, eroded-----	100	Somewhat limited		Somewhat limited		Somewhat limited	



AGRICULTURAL WASTE MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
119RX: Roxbury-----	100	Restricted permeability	0.30	Restricted permeability	0.22	Restricted permeability	0.22
119UA: Uly-----	100	Not limited		Somewhat limited Flooding	0.40	Not limited	
119UT: Uly-----	100	Not limited		Not limited		Not limited	
An: Bridgeport, channeled-----	100	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
ARR: Arkansas River-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Bc: Bridgeport-----	100	Not rated		Not rated		Not rated	
Bd: Bridgeport-----	100	Not limited		Somewhat limited Flooding	0.40	Not limited	
Bo: Optima-----	70	Not limited		Somewhat limited Flooding	0.40	Not limited	
Da: Dale-----	100	Very limited Filtering capacity Depth to dense layer	1.00 1.00	Very limited Filtering capacity	1.00	Very limited Filtering capacity Too steep for surface application	1.00 0.08
Gr: Schamber-----	100	Leaching limitation	0.45				
GRP: Pits-----	100	Not limited		Somewhat limited Flooding	0.40	Not limited	
Ha: Harney-----	100	Not rated		Not rated		Not rated	
HB: Harney-----	100	Not rated		Not rated		Not rated	
INL: Aquolls-----	100	Not rated		Not rated		Not rated	
La: Las Animas, occasionally flooded-----	50	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
		Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Low adsorption	1.00	Low adsorption	1.00	Low adsorption	1.00
		Ponding	1.00	Ponding	1.00	Ponding	1.00
		Somewhat limited		Very limited		Somewhat limited	
		Flooding	0.60	Flooding	1.00	Flooding	0.60
		Depth to saturated zone	0.43	Depth to saturated zone	0.43	Depth to saturated zone	0.43
		Salinity	0.06	Salinity	0.13	Salinity	0.13
		Filtering capacity	0.00	Filtering capacity	0.00	Filtering capacity	0.00

AGRICULTURAL WASTE MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food-processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Lesho, occasionally flooded-----	40	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Flooding	0.60	Flooding	1.00	Flooding	0.60
		Salinity	0.50	Depth to saturated zone	0.43	Depth to saturated zone	0.43
		Depth to saturated zone	0.43	Restricted permeability	0.22	Restricted permeability	0.22
		Restricted permeability	0.30	Salinity	0.13	Salinity	0.13
Lc: Las Animas, occasionally flooded-----	100	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Depth to saturated zone	0.95	Flooding	1.00	Depth to saturated zone	0.95
		Flooding	0.60	Depth to saturated zone	0.95	Flooding	0.60
		Droughty	0.01	Droughty	0.01	Droughty	0.01
Ld: Las Animas-----	100	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Depth to saturated zone	0.95	Flooding	1.00	Depth to saturated zone	0.95
		Flooding	0.60	Depth to saturated zone	0.95	Flooding	0.60
Le: Las, occasionally flooded-----	100	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Depth to saturated zone	0.86	Flooding	1.00	Depth to saturated zone	0.86
		Flooding	0.60	Depth to saturated zone	0.86	Flooding	0.60
		Restricted permeability	0.30	Restricted permeability	0.22	Restricted permeability	0.22
Lh: Lesho, occasionally flooded-----	100	Somewhat limited		Very limited		Somewhat limited	
		Flooding	0.60	Flooding	1.00	Flooding	0.60
		Depth to saturated zone	0.43	Depth to saturated zone	0.43	Depth to saturated zone	0.43
		Restricted permeability	0.30	Restricted permeability	0.22	Restricted permeability	0.22
		Filtering capacity	0.00	Filtering capacity	0.00	Filtering capacity	0.00
Lk: Lesho-----	50	Somewhat limited		Very limited		Somewhat limited	
		Flooding	0.60	Flooding	1.00	Flooding	0.60
		Depth to saturated zone	0.43	Depth to saturated zone	0.43	Depth to saturated zone	0.43
		Restricted permeability	0.30	Restricted permeability	0.22	Restricted permeability	0.22
		Filtering capacity	0.00	Filtering capacity	0.00	Filtering capacity	0.00
Sweetwater-----	25	Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Flooding	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00	Filtering capacity	1.00
		Flooding	0.60	Filtering capacity	1.00	Flooding	0.60
		Runoff limitation	0.40	Restricted permeability	0.22	Restricted permeability	0.22
		Restricted permeability	0.30				
Ll: Lincoln, occasionally flooded-----	100	Very limited		Very limited		Very limited	
		Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Flooding	1.00	Flooding	1.00	Flooding	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Lo: Lofton-----	100	Leaching limitation	0.45				
		Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Runoff limitation	0.40				
Lu: Lubbock-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
Md: Penden-----	100	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
Me: Penden-----	100	Somewhat limited Slope	0.37	Somewhat limited Slope	0.37	Very limited Too steep for surface application	1.00
						Too steep for sprinkler application	0.59
Mf: Penden, Severely eroded-----	100	Not limited		Not limited		Somewhat limited  Too steep for surface application	0.08
Mh: Penden, Severely Eroded-----	100	Somewhat limited  Slope	0.37	Somewhat limited  Slope	0.37	Very limited	
						Too steep for surface application	1.00
						Too steep for sprinkler application	0.59
Mm: Manter-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00
Mn: Manter-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00
Mo: Manter-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Too steep for surface application	0.08
						Filtering capacity	0.00
Mp: Campus-----	75	Very limited Depth to bedrock Droughty	1.00 0.91	Very limited Depth to bedrock Droughty	1.00 0.91	Very limited Depth to bedrock Droughty	1.00 0.91
						Too steep for surface application	0.91
						Too steep for sprinkler application	0.02
Canlon-----	25	Very limited Depth to bedrock Droughty Runoff limitation	1.00 1.00 0.40	Very limited Droughty Depth to bedrock Slope	1.00 1.00 0.04	Very limited Droughty	1.00
						Depth to bedrock	1.00
						Too steep for surface application	1.00
Mu: Manter-----	55	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Too steep for sprinkler application	0.22
						Somewhat limited Too steep for surface application	0.08

AGRICULTURAL WASTE MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ulysses-----	45	Not limited		Not limited		Filtering capacity Somewhat limited Too steep for surface application	0.00 0.08
Pa: Pratt-----	100	Very limited Filtering capacity Leaching limitation Droughty	1.00 0.45 0.00	Very limited Filtering capacity Droughty	1.00 0.00	Very limited Filtering capacity Too steep for surface application Droughty	1.00 0.08 0.00
Pg: Pratt-----	100	Very limited Filtering capacity Leaching limitation	1.00 0.45	Very limited Filtering capacity	1.00	Very limited Filtering capacity	1.00
Pt: Pratt-----	60	Very limited Filtering capacity Leaching limitation Droughty	1.00 0.45 0.00	Very limited Filtering capacity Droughty	1.00 0.00	Very limited Filtering capacity Too steep for surface application Droughty Too steep for sprinkler application	1.00 0.66 0.00 0.00
Optima-----	40	Very limited Filtering capacity Depth to dense layer Leaching limitation	1.00 1.00 0.45	Very limited Filtering capacity	1.00	Very limited Filtering capacity Too steep for surface application Too steep for sprinkler application	1.00 0.91 0.02
Ra: Ness-----	100	Very limited Restricted permeability Ponding Depth to saturated zone Runoff limitation	1.00 1.00 1.00 0.40	Very limited Restricted permeability Ponding Depth to saturated zone	1.00 1.00 1.00	Very limited Restricted permeability Ponding Depth to saturated zone	1.00 1.00 1.00
Rm: Richfield-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
Rn: Richfield-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
Ro: Richfield-----	100	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
Rs: Richfield-----	60	Somewhat limited Restricted permeability	0.30	Somewhat limited Restricted permeability	0.22	Somewhat limited Restricted permeability	0.22
Spearville-----	40	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
Sa: Satanta-----	100	Not limited		Not limited		Not limited	
SAP: Sand Pit-----	100	Not rated		Not rated		Not rated	
Sb: Satanta-----	100	Not limited		Not limited		Not limited	
Sp: Spearville-----	100	Very limited		Very limited		Very limited	

AGRICULTURAL WASTE MANAGEMENT--Continued  
Gray County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Sr: Spearville-----	100	Restricted permeability	1.00	Restricted permeability	1.00	Restricted permeability	1.00
		Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
Sw: Sweetwater-----	100	Very limited	1.00	Very limited	1.00	Very limited	1.00
		Depth to saturated zone	1.00	Flooding	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00	Depth to saturated zone	1.00	Filtering capacity	1.00
		Flooding	0.60	Filtering capacity	1.00	Flooding	0.60
		Runoff limitation	0.40	Restricted permeability	0.22	Restricted permeability	0.22
Tf: Optima-----	90	Restricted permeability	0.30				
		Very limited	1.00	Very limited	1.00	Very limited	1.00
		Filtering capacity	1.00	Filtering capacity	0.63	Filtering capacity	1.00
		Depth to dense layer	1.00	Slope		Too steep for surface application	1.00
Ua: Ulysses-----	100	Slope	0.63			Too steep for sprinkler application	0.77
		Leaching limitation	0.45				
		Not limited		Not limited		Not limited	
		Not limited		Not limited		Not limited	
		Not limited		Not limited		Not limited	
Uc: Ulysses-----	100	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
		Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
UCC: Uly-----	100	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
		Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
Um: Ulysses-----	55	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
		Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
Colby-----	45	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
		Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
W: Water-----	100	Not rated		Not rated		Not rated	

WIN-PST SPISP II  
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL\_KS Sort Order: MUSYM

Gray County, Kansas: KS069

MUSYM/SEQ#	COMPONENT/TEXTURE/MU%	HYD	KFACT	SURFACE DEPTH	% OM	SPISP II Ratings		
						Leaching (SLP)	Solution Runoff (SSRP)	Adsorbed Runoff (SARP)
055BP 1	BRIDGEPORT CL 100%	B	0.28	11"	2.0%	I	I	I
055LD 1	LAS L 55%	C	0.32	8"	0.8%	H (w)	H	H
055LD 2	LAS ANIMAS FSL 45%	C	0.24	14"	0.8%	H (w)	H	H
055TF 1	OPTIMA LFS 90%	A	0.17	8"	0.5%	H	L	I (s)
055TV 1	EVA LFS 50%	B	0.10	13"	0.9%	H	I	L
055TV 2	OPTIMA LFS 50%	A	0.17	8"	0.5%	H	L	L
055VO 1	EVA LFS 85%	B	0.10	13"	0.9%	H	I	L
081LU 1	LUBBOCK SICL 100%	B	0.32	9"	2.0%	I	I	I
083UM 1	ULY SIL 70%	B	0.32	6"	1.5%	H	I	I
083UM 2	COLY SIL 30%	B	0.43	6"	0.8%	H	I	I
119HC 1	HARNEY SICL 100%	B	0.32	6"	1.4%	H	I	I
119RX 1	ROXBURY SIL 100%	B	0.32	20"	2.0%	I	I	I
119UA 1	ULY SIL 100%	B	0.32	6"	1.5%	H	I	I
119UT 1	ULY SIL 100%	B	0.32	6"	1.5%	H	I	I
An 1	BRIDGEPORT L 100%	B	0.28	12"	2.0%	I	I	I
ARR 1	ARKANSAS RIVER S 100%	D	0.00	6"	0.1%	H (w)	H	L
Bc 1	BRIDGEPORT SICL 100%	B	0.32	10"	2.0%	I	I	I
Bd 1	BRIDGEPORT SICL 100%	B	0.32	10"	2.0%	I	I	I
Bo 1	OPTIMA LFS 70%	A	0.17	8"	0.5%	H	L	L
Da 1	DALE SIL 100%	B	0.32	25"	2.0%	L	I	I
Gr 1	SCHAMBER GR-SL 100%	A	0.17	6"	0.8%	H	L	I (s)
GRP 1	PITS GR-S 100%	A	0.10	60"	0.3%	H	L	I (s)
Ha 1	HARNEY SIL 100%	B	0.32	9"	2.0%	I	I	I
HB 1	HARNEY SIL 100%	B	0.32	6"	2.0%	I	I	I
INL 1	AQUOLLS VAR 100%	C	0.00	72"	0.0%	?	H	?
La 1	LAS ANIMAS SL 50%	C	0.24	11"	0.8%	H (w)	H	H
La 2	LESHO CL 40%	C	0.28	14"	1.5%	H (w)	H	H
Lc 1	LAS ANIMAS LS 100%	C	0.17	11"	0.8%	H (w)	H	I

WIN-PST SPISP II  
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL\_KS Sort Order: MUSYM

Gray County, Kansas: KS069

Ld 1	LAS ANIMAS SL 100%	C	0.24	11"	0.8% H (w)	H	H
Le 1	LAS CL 100%	C	0.32	8"	0.8% H (w)	H	H
Lh 1	LESHO CL 100%	C	0.28	14"	1.5% H (w)	H	H
Lk 1	LESHO L 50%	C	0.28	14"	1.5% H (w)	H	H
Lk 2	SWEETWATER CL 25%	D	0.28	17"	0.8% H (w)	H	H
Ll 1	LINCOLN FSL 100%	A	0.24	8"	0.8% H	L	L
Lo 1	LOFTON SICL 100%	D	0.37	6"	2.0% V	H	H
Lu 1	LUBBOCK L 100%	B	0.28	15"	1.5% I	I	I
Md 1	PENDEN CL 100%	B	0.28	14"	1.5% I	I	I
Me 1	PENDEN CL 100%	B	0.28	15"	0.8% I	I	I
Mf 1	PENDEN CL 100%	B	0.28	4"	0.8% H	I	I
Mh 1	PENDEN CL 100%	B	0.28	4"	0.8% H	I	I
Mm 1	MANTER FSL 100%	B	0.20	18"	0.8% H	I	I
Mn 1	MANTER FSL 100%	B	0.20	18"	0.8% H	I	I
Mo 1	MANTER FSL 100%	B	0.20	18"	0.8% H	I	I
Mp 1	CAMPUS L 75%	B	0.28	9"	1.3% I	I	I
Mp 2	CANLON SIL 25%	D	0.37	8"	0.8% V	H	H
Mu 1	MANTER FSL 55%	B	0.20	18"	0.8% H	I	I
Mu 2	ULYSSES SIL 45%	B	0.32	5"	1.5% H	I	I
Pa 1	PRATT LFS 100%	A	0.17	11"	0.8% H	L	L
Pg 1	PRATT LFS 100%	A	0.17	11"	0.8% H	L	L
Pt 1	PRATT LFS 60%	A	0.17	11"	0.8% H	L	L
Pt 2	OPTIMA LFS 40%	A	0.17	8"	0.5% H	L	L
Ra 1	NESS C 100%	D	0.28	31"	2.0% H (w)	H	H
Rm 1	RICHFIELD SIL 100%	B	0.32	5"	1.5% H	I	I
Rn 1	RICHFIELD SIL 100%	B	0.32	6"	1.5% H	I	I
Ro 1	RICHFIELD SICL 100%	B	0.32	6"	1.5% H	I	I
Rs 1	RICHFIELD SIL 60%	B	0.32	6"	1.5% H	I	I
Rs 2	SPEARVILLE SICL 40%	C	0.37	7"	2.0% L	H	H
Sa 1	SATANTA L 100%	B	0.28	10"	1.5% I	I	I

WIN-PST SPISP II  
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL\_KS Sort Order: MUSYM

Gray County, Kansas: KS069

SAP 1	SAND PIT 100%		0.00	0"	0.0% ?	?	?
Sb 1	SATANTA L 100%	B	0.28	10"	1.5% I	I	I
Sp 1	SPEARVILLE SICL 100%	C	0.32	7"	2.0% L	H	H
Sr 1	SPEARVILLE SICL 100%	C	0.32	7"	2.0% L	H	H
Sw 1	SWEETWATER CL 100%	D	0.28	17"	0.8% H (w)	H	H
Tf 1	OPTIMA LFS 90%	A	0.17	8"	0.5% H	L	I (s)
Ua 1	ULYSSES SIL 100%	B	0.32	5"	1.5% H	I	I
Ub 1	ULYSSES SIL 100%	B	0.32	5"	1.5% H	I	I
UBB 1	ULY SIL 100%	B	0.32	8"	1.5% I	I	I
Uc 1	ULYSSES SIL 100%	B	0.32	6"	1.5% H	I	I
UCC 1	ULY SIL 100%	B	0.32	6"	1.5% H	I	I
Um 1	ULYSSES SIL 55%	B	0.32	5"	1.5% H	I	I
Um 2	COLBY SIL 45%	B	0.43	5"	0.8% H	I	I
W 1	WATER 100%		0.00	0"	0.0% ?	?	?

(.\REPORTS\SOILS.TXT generated on 12/12/01 at 12:11:15)

H -- High  
I -- Intermediate  
L -- Low  
V -- Very Low

Conditions that affect ratings:

- m -- There are macropores in the surface horizon deeper than 24"
- w -- The high water table comes within 24" of the surface during the growing season
- s -- The field slope is greater than 15%

SPISP II S-Ratings:

- SLP -- Soil Leaching Potential
- SSRP -- Soil Solution Runoff Potential
- SARP -- Soil Adsorbed Runoff Potential



In this section, hydric soils are defined and described and the hydric soils in the survey area are listed. The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin and others, 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for each of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part (Federal Register, 1994). These soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 1995). These criteria are used to identify a phase of a soil series that normally is associated with wetlands. The criteria used are selected estimated soil properties that are described in "Soil Taxonomy" (USDA, 1999) and "Keys to Soil Taxonomy" (USDA, 1998) and in the "Soil Survey Manual" (USDA, 1993).

If soils are wet enough for a long enough period to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils in this survey area are specified in "Field Indicators of Hydric Soils in the United States" (Hurt and others, 1996).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

Map units in the Hydric Soil Interpretations table meet the definition of hydric soils and, in addition, have at least one of the hydric soil indicators. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; Hurt and others, 1996).

Map units that are made up of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units made up of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform.

These map units, in general, do not meet the definition of hydric soils because they do not have one of the hydric soil indicators. A portion of these map units, however, may include hydric soils. Onsite investigation is recommended to determine whether hydric soils occur and the location of the included hydric soils.

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Gray County, Kansas

PAGE 2 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
055BP: BRIDGEPORT CLAY LOAM, RARELY FLOODED	BRIDGEPORT	No	flood plain	---	---	---	---
055LD: LAS-LAS ANIMAS COMPLEX, OCCASIONALLY FLOODED	SWEETWATER	Yes	flood plain	2B3	YES	NO	NO
	LAS	No	flood plain	---	---	---	---
	LAS ANIMAS	No	flood plain	---	---	---	---
055TF: VALENT FINE SAND, 6 TO 20 PERCENT SLOPES	VALENT	No	dune, paleoterrace	---	---	---	---
055TV: VALENT-VONA LOAMY FINE SANDS, 3 TO 8 PERCENT SLOPES	VALENT	No	dune, paleoterrace	---	---	---	---
	VONA	No	dune, paleoterrace	---	---	---	---
	UNNAMED HYDRIC SOILS	Yes	depression	2B3	YES	NO	NO
055VO: VONA LOAMY FINE SAND, 1 TO 5 PERCENT SLOPES	VONA	No	dune, paleoterrace	---	---	---	---
	UNNAMED HYDRIC SOILS	Yes	depression	2B3	YES	NO	NO
081LU: LUBBOCK SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES	LUBBOCK	No	paleoterrace	---	---	---	---
083UM: ULY-COLY SILT LOAMS, 3 TO 6 PERCENT SLOPES, ERODED	ULY	No	plain	---	---	---	---
	COLY	No	plain	---	---	---	---
119HC: HARNEY SILTY CLAY LOAM, 1 TO 3 PERCENT SLOPES, ERODED	HARNEY	No	plain	---	---	---	---
119RX: ROXBURY SILT LOAM, RARELY FLOODED	ROXBURY	No	flood plain	---	---	---	---
119UA: ULY SILT LOAM, 0 TO 1 PERCENT SLOPES	ULY	No	plain	---	---	---	---
	NESS	Yes	playa	2B3,3	YES	NO	YES
119UT: ULY SILT LOAM, 3 TO 6 PERCENT SLOPES, ERODED	ULY	No	plain	---	---	---	---
An: BRIDGEPORT SILT LOAM, CHANNELED	BRIDGEPORT	No	flood plain	---	---	---	---
ARR: ARKANSAS RIVER	ARKANSAS RIVER	Unranked	---	---	---	---	---
Bc: BRIDGEPORT SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES	BRIDGEPORT	No	flood plain	---	---	---	---
Bd: BRIDGEPORT SILTY CLAY LOAM, 1 TO 3 PERCENT SLOPES	BRIDGEPORT	No	flood plain	---	---	---	---
Bo: OPTIMA FINE SAND, 0 TO 10 PERCENT SLOPES	OPTIMA	No	dune, paleoterrace	---	---	---	---
	EVA	No	dune, paleoterrace	---	---	---	---
Da: DALE SILT LOAM, RARELY FLOODED	DALE	No	flood plain	---	---	---	---
Gr: SCHAMBER GRAVELLY SANDY LOAM, 5 TO 25 PERCENT SLOPES	SCHAMBER	No	fan remnant, paleoterrace	---	---	---	---
GRP: GRAVEL PIT	PITS	Unranked	---	---	---	---	---
Ha: HARNEY SILT LOAM, 0 TO 1 PERCENT SLOPES	HARNEY	No	plain	---	---	---	---
	NESS	Yes	depression	3,2B3	YES	NO	YES
HB: HARNEY SILT LOAM, 1 TO 3 PERCENT SLOPES	HARNEY	No	plain	---	---	---	---

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Gray County, Kansas

PAGE 3 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
INL: AQUOLLS	AQUOLLS	Yes	depression, terrace	3,2B3	YES	NO	YES
La: LAS ANIMAS-LESHO COMPLEX, ALKALI, OCCASIONALLY FLOODED	LAS ANIMAS	No	flood plain	---	---	---	---
	LESHO SLICKSPOTS SWEETWATER	No No Yes	flood plain stream terrace flood plain	---	---	---	---
Lc: LAS ANIMAS SOILS, OCCASIONALLY FLOODED	LAS ANIMAS	No	flood plain	---	---	---	---
	SWEETWATER	Yes	flood plain	---	---	---	---
Ld: LAS ANIMAS SANDY LOAM, OCCASIONALLY FLOODED	LAS ANIMAS	No	flood plain	---	---	---	---
	SWEETWATER	Yes	flood plain	2B3	YES	NO	NO
Le: LAS CLAY LOAM, OCCASIONALLY FLOODED	LAS	No	flood plain	---	---	---	---
	SWEETWATER	Yes	flood plain	2B3	YES	NO	NO
Lh: LESHO CLAY LOAM, OCCASIONALLY FLOODED	LESHO	No	flood plain	---	---	---	---
	SWEETWATER	Yes	flood plain	2B3	YES	NO	NO
Lk: LESHO-SWEETWATER COMPLEX, OCCASIONALLY FLOODED	LESHO	No	flood plain	---	---	---	---
	SWEETWATER	Yes	flood plain	2B3	YES	NO	NO
	LAS ANIMAS OPTIMA	No No	flood plain dune, paleoterrace	---	---	---	---
Ll: LINCOLN SOILS, FREQUENTLY FLOODED	LINCOLN	No	flood plain	---	---	---	---
	SWEETWATER	Yes	flood plain	2B3	YES	NO	NO
Lo: LOFTON SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES	LOFTON	Yes	bench, playa	2B3,3	YES	NO	YES
Lu: LUBBOCK LOAM, 0 TO 1 PERCENT SLOPES	LUBBOCK	No	paleoterrace	---	---	---	---
Md: PENDEN CLAY LOAM, 3 TO 6 PERCENT SLOPES	PENDEN	No	plain	---	---	---	---
Me: PENDEN CLAY LOAM, 6 TO 15 PERCENT SLOPES	PENDEN	No	plain	---	---	---	---
Mf: PENDEN CLAY LOAM, 3 TO 6 PERCENT SLOPES, SEVERELY ERODED	PENDEN	No	plain	---	---	---	---
Mh: PENDEN COMPLEX, 6 TO 15 PERCENT SLOPES, SEVERELY ERODED	PENDEN	No	plain	---	---	---	---
Mm: MANTER FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES	MANTER	No	paleoterrace, sand sheet	---	---	---	---
	UNNAMED HYDRIC SOILS	Yes	depression	2B3	YES	NO	NO
Mn: MANTER FINE SANDY LOAM, 1 TO 3 PERCENT SLOPES	MANTER	No	paleoterrace, sand sheet	---	---	---	---
Mo: MANTER FINE SANDY LOAM, 1 TO 6 PERCENT SLOPES, ERODED	MANTER	No	paleoterrace, sand sheet	---	---	---	---
	UNNAMED HYDRIC SOILS	Yes	depression	2B3	YES	NO	NO
Mp: CAMPUS-CANLON COMPLEX, 6 TO 15 PERCENT SLOPES	CAMPUS	No	---	---	---	---	---
	CANLON	No	---	---	---	---	---

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Gray County, Kansas

PAGE 4 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Mu: MANter-ULYSSES COMPLEX, 1 TO 6 PERCENT SLOPES	MANter	No	paleoterrace, sand sheet	---	---	---	---
Pa: PRATT LOAMY FINE SAND, 1 TO 6 PERCENT SLOPES	ULYSSES	No	plain	---	---	---	---
	PRATT	No	dune, paleoterrace depression	---	---	---	---
	UNNAMED HYDRIC SOILS	Yes		2B3	YES	NO	NO
Pg: PRATT LOAMY FINE SAND, GRAVEL SUBSTRATUM, 0 TO 1 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
Pt: PRATT-OPTIMA LOAMY FINE SANDS, 3 TO 9 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
	OPTIMA	No	dune, paleoterrace	---	---	---	---
Ra: NESS CLAY	NESS	Yes	playa	2B3,3	YES	NO	YES
Rm: RICHFIELD SILT LOAM, 0 TO 1 PERCENT SLOPES	RICHFIELD	No	plain	---	---	---	---
	NESS	Yes	playa	2B3,3	YES	NO	YES
Rn: RICHFIELD SILT LOAM, 1 TO 3 PERCENT SLOPES	RICHFIELD	No	plain	---	---	---	---
	NESS	Yes	playa	2B3,3	YES	NO	YES
Ro: RICHFIELD SILTY CLAY LOAM, 1 TO 3 PERCENT SLOPES, ERODED	RICHFIELD	No	plain	---	---	---	---
Rs: RICHFIELD-SPEARVILLE COMPLEX, 0 TO 1 PERCENT SLOPES	RICHFIELD	No	plain	---	---	---	---
	SPEARVILLE NESS	No Yes	plain depression	--- 2B3,3	--- YES	--- NO	--- YES
Sa: SATANTA LOAM, 0 TO 1 PERCENT SLOPES	SATANTA	No	paleoterrace, sand sheet	---	---	---	---
	NESS	Yes	playa	2B3,3	YES	NO	YES
SAP: SAND PIT	SAND PIT	Unranked	---	---	---	---	---
Sb: SATANTA LOAM, 1 TO 3 PERCENT SLOPES	SATANTA	No	paleoterrace, sand sheet	---	---	---	---
Sp: SPEARVILLE SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES	SPEARVILLE	No	plain	---	---	---	---
	NESS	Yes	depression	2B3,3	YES	NO	YES
Sr: SPEARVILLE COMPLEX, 1 TO 3 PERCENT SLOPES, ERODED	SPEARVILLE	No	plain	---	---	---	---
Sw: SWEETWATER SOILS, OCCASIONALLY FLOODED	SWEETWATER	Yes	flood plain	2B3	YES	NO	NO
Tf: OPTIMA FINE SAND, 6 TO 20 PERCENT SLOPES	OPTIMA	No	dune, paleoterrace	---	---	---	---
	EVA	No	dune, paleoterrace	---	---	---	---
	UNNAMED HYDRIC SOILS	Yes	depression	2B3	YES	NO	NO
Ua: ULYSSES SILT LOAM, 0 TO 1 PERCENT SLOPES	ULYSSES	No	plain	---	---	---	---
	NESS	Yes	depression	2B3,3	YES	NO	YES
Ub: ULYSSES SILT LOAM, 1 TO 3 PERCENT SLOPES	ULYSSES	No	plain	---	---	---	---
UBB: ULY SILT LOAM, 1 TO 3 PERCENT SLOPES	ULY	No	plain	---	---	---	---
	NESS	Yes	playa	2B3,3	YES	NO	YES
Uc: ULYSSES SILT LOAM, 3 TO 6 PERCENT SLOPES	ULYSSES	No	plain	---	---	---	---

HYDRIC SOIL INTERPRETATIONS  
HYDRIC SOILS LIST  
Gray County, Kansas

PAGE 5 of 5

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
UCC: ULY SILT LOAM, 3 TO 6 PERCENT SLOPES	ULY	No	plain	---	---	---	---
Um: ULYSSES-COLBY SILT LOAMS, 3 TO 6 PERCENT SLOPES, ERODED	ULYSSES	No	plain	---	---	---	---
	COLBY	No	hillslope	---	---	---	---
W: WATER	WATER	Yes	---	4,3	NO	YES	YES

FOOTNOTE: There may be small areas of included soils or miscellaneous areas that are significant to use and management of the soil; yet are too small to delineate on the soil map at the map's original scale. These may be designated as spot symbols and are defined in the published Soil Survey Report or the USDA-NRCS Technical Guide, Part II. Areas mapped as water or any map unit that contains one of the following conventional symbols is considered a hydric soil map unit: marshes or swamps; wet spots; depressions; streams, lakes and ponds.

1. All Histosols except Folists, or
2. Soils in Aquic suborders, great groups, or subgroups, Albolls suborder, Aquisalids, Pachic subgroups, or Cumulic subgroups that are:
  - a. Somewhat poorly drained with a water table equal to 0.0 foot (ft) from the surface during the growing season, or
  - b. poorly drained or very poorly drained and have either:
    - (1) water table equal to 0.0 ft during the growing season if textures are coarse sand, sand, or fine sand in all layers within 20 inches (in), or for other soils
    - (2) water table at less than or equal to 0.5 ft from the surface during the growing season if permeability is equal to or greater than 6.0 in/hour (h) in all layers within 20 in, or
    - (3) water table at less than or equal to 1.0 ft from the surface during the growing season if permeability is less than 6.0 in/h in any layer within 20 in, or
3. Soils that are frequently ponded for long duration or very long duration during the growing season, or
4. Soils that are frequently flooded for long duration or very long duration during the growing season.