

Prime farmland is one of several kinds of important farmland defined by the U.S. Department of Agriculture. It is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil qualities, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. It is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

A recent trend in land use in some parts of the survey area has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

The map units in the survey area that are considered prime farmland are listed in the following table. This list does not constitute a recommendation for a particular land use. On some soils included in the list, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures. The extent of each listed map unit is shown in the "Acres and Proportionate Extent of Soils" table. The location is shown on the detailed soil maps. The soil qualities that affect use and management are described in other tables in this document."

Map symbol	Mapunit name	Farmland Classification
085WB	Wymore silty clay loam, 1 to 3 percent slopes	All areas are prime farmland
601GT	Grundy silty clay loam, 1 to 3 percent slopes	All areas are prime farmland
601SH	Shelby loam, 4 to 8 percent slopes	All areas are prime farmland
Bp	Belvue silt loam, escarpment, 2 to 12 percent slopes	All areas are prime farmland
Bx	Bourbonais-bismarckgrove complex, 0 to 2 percent slopes, rarely flooded	All areas are prime farmland
Eg	Eudora silt loam, 0 to 2 percent slopes, rarely flooded	All areas are prime farmland
Gb	Grundy silty clay loam, 0 to 2 percent slopes	All areas are prime farmland
Gc	Grundy silty clay loam, 2 to 5 percent slopes	All areas are prime farmland
Gy	Gymer silt loam, 3 to 7 percent slopes	All areas are prime farmland
Kb	Kennebec silt loam, 0 to 2 percent slopes, occasionally flooded	All areas are prime farmland
Mb	Martin silty clay loam, 1 to 3 percent slopes	All areas are prime farmland
MR	Morrill clay loam, 3 to 7 percent slopes	All areas are prime farmland
Mv	Morrill loam, 3 to 7 percent slopes	All areas are prime farmland
Pb	Pawnee clay loam, 1 to 3 percent slopes	All areas are prime farmland
Re	Reading silt loam, 0 to 2 percent slopes, very rarely flooded, moderately wet	All areas are prime farmland
Rs	Rossville silt loam, 0 to 2 percent slopes, very rarely flooded	All areas are prime farmland
Hc	Haig silty clay loam, 0 to 2 percent slopes	Prime farmland if drained
Wc	Wabash silty clay loam, 0 to 1 percent slopes, occasionally flooded	Prime farmland if drained

