

Table of Contents Section II – Soil and Site Information

	Issue Date	Date of Last Review	Responsible Staff
<i>Use and Explanation of Soil Interpretations</i>			
<i>Explanation of Key Phrases Used in Soil Interpretations</i>			
Soils Legends	1/02	1/02	SOI
<i>* Acreage and Proportionate Extent of the Soils</i>			
Soil Descriptions - Nontechnical	1/02	1/93	SOI
<i>Use and Explanation of Nontechnical Descriptions</i>			
<i>*Nontechnical Soils Description Report</i>			
Soil Descriptions - Technical	1/02	1/02	SOI
<i>*Map Unit Description Report</i>			
Cropland Interpretations - Technical	1/02	1/02	SOI
<i>*Prime Farmland Report</i>			
<i>* Kansas Soil Rating for Plant Growth Index</i>			
<i>*Soil Properties for Conservation Planning</i>			
Rangeland, Grazed Forestland, Native Pastureland Interpretations	1/02	1/02	SOI
<i>*Rangeland Productivity Report</i>			
<i>*Range Site Descriptions</i>			
Forestland Interpretations	1/93	1/93	SOI
<i>Use and Explanation of Forestland Interpretations</i>			
<i>*Woodland Management and Productivity</i>			
Nonagricultural Interpretations	1/02	1/02	SOI
<i>*Building Site Development Report</i>			
<i>*Construction Materials Report</i>			
Recreation Interpretations	1/02	1/02	SOI
<i>*Recreational Interpretations</i>			
Wildlife Interpretations	1/02	1/02	SOI
<i>*Wildlife Interpretations Report</i>			
Pastureland and Hayland Interpretations	1/02	1/02	SOI
<i>*Yields Per Acre of Pasture and Hayland</i>			

	Issue Date	Date of Last Review	Responsible Staff
Mined Land Interpretations <i>Use and Explanation of Mined Land Interpretations</i>	1/93	1/93	SOI
Windbreak Interpretations <i>*Conservation Tree and Shrub Management Report</i>	1/02	1/02	SOI
Engineering Interpretations <i>*Engineering Index Properties</i> <i>*Physical Properties of the Soils</i> <i>*Chemical Properties of the Soils</i> <i>*Water Features</i> <i>*Soil Features</i> <i>*Water Management Report</i>	1/02	1/02	SOI
Waste Disposal Interpretations <i>*Sanitary Facilities Report</i> <i>*Agricultural Waste Management Report</i>	1/02	1/02	SOI
Water Quantity and Quality Interpretations <i>Use and Explanation of Water Quantity and Quality Interpretations</i> <i>*Appendix A – Soils Potential For Surface Loss and Leaching</i> <i>*Appendix B – Pesticide Selected Properties Database</i> <i>*Appendix C – Herbicide Selected Properties Database</i> <i>*Soil-Pesticide Interaction Screening Procedure Worksheet (Blank)</i> <i>*WIN-PST SPISP II Soil Sensitivity to Pesticide Loss Rating Report</i>	1/02	1/02	SOI
Hydric Soil Interpretations <i>Use and Explanation of Hydric Soil Interpretations</i> <i>*Hydric Soils List</i>	1/02	1/02	SOI
HEL Interpretations <i>Use and Explanation of Highly Erodible Land Interpretations</i> <i>*Highly Erodible Lands Report</i> <i>*LS and Supporting Data for 1990 Frozen HEL List</i> <i>*CRP 20 Soil Supporting Data for 1990 Frozen HEL List</i>	7/95	1/00	SOI

**County specific computer generated reports.*

ACREAGE AND PROPORTIONATE EXTENT OF THE SOILS

McPherson County, Kansas: Published

Map symbol	Soil name	Acres	Percent
053ED	Edalgo Loam, 3 To 7 Percent Slopes-----	139	*
053MB	Mccook Loam, Occasionally Flooded-----	24	*
053MC	Mccook Silt Loam, Rarely Flooded-----	62	*
079GE	Geary Silt Loam, 3 To 6 Percent Slopes-----	14	*
079LB	Ladysmith Silty Clay Loam, 1 To 2 Percent Slopes-----	85	*
115IC	Irwin Silty Clay Loam, 3 To 6 Percent Slopes-----	10	*
159DP	Dillwyn-Plevna Loamy Fine Sands, 0 To 1 Percent Slopes-----	99	*
159QC	Geary-Clark Complex, 3 To 7 Percent Slopes, Eroded-----	5	*
159LA	Lancaster Loam, 1 To 3 Percent Slopes-----	18	*
159NF	Naron Fine Sandy Loam, 1 To 3 Percent Slopes-----	6	*
159PF	Pratt Loamy Fine Sand, 1 To 5 Percent Slopes-----	93	*
159PR	Pratt-Carwile Complex, 0 To 5 Percent Slopes-----	15	*
159SO	Smolan Soils, 2 To 7 Percent Slopes, Eroded-----	86	*
159WA	Waldeck Fine Sandy Loam, Occasionally Flooded-----	38	*
169CE	Clime Silty Clay Loam, 2 To 6 Percent Slopes-----	836	0.1
169ED	Edalgo Clay Loam, 3 To 7 Percent Slopes-----	68	*
169RO	Roxbury Silt Loam, Rarely Flooded-----	2,471	0.4
1191	Blazefork Silty Clay Loam, 0 To 1 Percent Slopes, Rarely Flooded-----	60	*
1200	Buhler-Blazefork Silty Clay Loams, 0 To 1 Percent Slopes, Rarely Flooded-----	139	*
1324	Carway And Carbika Soils, 0 To 1 Percent Slopes-----	190	*
1553	Darlow-Elmer Complex, 0 To 2 Percent Slopes-----	16	*
1555	Dillhut-Plev Complex, 0 To 2 Percent Slopes-----	403	*
1985	Hayes Fine Sandy Loam, 1 To 5 Percent Slopes-----	79	*
2391	Kaskan Silty Clay Loam, 0 To 1 Percent Slopes, Frequently Flooded, Channeled-----	2	*
2588	Longford Silty Clay Loam, 3 To 7 Percent Slopes, Moderately Eroded-----	141	*
3181	Pratt-Turon Fine Sands, 1 To 5 Percent Slopes-----	1	*
3641	Tivin-Dillhut Fine Sands, 0 To 15 Percent Slopes-----	132	*
AED	Arents, Earthen Dam-----	13	*
At	Attica Loamy Fine Sand, 1 To 4 Percent Slopes-----	2,015	0.3
Br	Bridgeport Silt Loam, Rarely Flooded-----	8,522	1.5
Ca	Carwile Fine Sandy Loam, 0 To 1 Percent Slopes-----	4,702	0.8
Cb	Cass Fine Sandy Loam, Rarely Flooded-----	3,152	0.5
Ce	Clime Silty Clay, 1 To 3 Percent Slopes-----	4,299	0.7
Cm	Clime Silty Clay, 3 To 6 Percent Slopes-----	7,529	1.3
Cr	Crete Silt Loam, 0 To 1 Percent Slopes-----	101,122	17.5
Cs	Crete Silt Loam, 1 To 3 Percent Slopes-----	90,867	15.8
Ct	Crete Silty Clay Loam, 1 To 3 Percent Slopes, Eroded-----	6,761	1.2
De	Detroit Silty Clay Loam, Rarely Flooded-----	4,600	0.8
Dr	Drummond Loam, 0 To 1 Percent Slopes-----	659	0.1
Ed	Edalgo Silt Loam, 5 To 12 Percent Slopes-----	11,752	2.0
Ee	Edalgo Silty Clay Loam, 3 To 9 Percent Slopes, Eroded-----	1,127	0.2
Fa	Farnum Loam, 1 To 3 Percent Slopes-----	11,757	2.0
Ge	Geary Silt Loam, 1 To 3 Percent Slopes-----	2,077	0.4
Go	Goessel Silty Clay, 0 To 2 Percent Slopes-----	19,062	3.3
Ho	Hord Silt Loam, Rarely Flooded-----	15,757	2.7
INT	Aquolls-----	148	*
Ir	Irwin Silty Clay Loam, 1 To 3 Percent Slopes-----	17,014	3.0
La	Ladysmith Silty Clay Loam, 0 To 1 Percent Slopes-----	79,774	13.8
Ld	Ladysmith-Drummond Complex, 0 To 1 Percent Slopes-----	668	0.1
Le	Lancaster Loam, 2 To 6 Percent Slopes-----	10,430	1.8
Lh	Lancaster-Hedville Loams, 6 To 12 Percent Slopes-----	35,500	6.2
LHH	Lancaster-Hedville Complex, 3 To 20 Percent Slopes-----	27,000	4.7
Ln	Longford Silty Clay Loam, 3 To 6 Percent Slopes-----	23,135	4.0
Lo	Longford Silty Clay Loam, 2 To 6 Percent Slopes, Eroded-----	5,860	1.0
Mc	Mccook Fine Sandy Loam, Rarely Flooded-----	2,229	0.4
Ns	Ness Silty Clay-----	604	0.1
Nw	New Cambria Silty Clay, Rarely Flooded-----	1,440	0.2
Pa	Plevna Fine Sandy Loam, Frequently Flooded-----	388	*
Pr	Pratt Loamy Fine Sand, 6 To 12 Percent Slopes-----	291	*
Ro	Roxbury Silty Clay Loam, Rarely Flooded-----	6,010	1.0
Sm	Smolan Silty Clay Loam, 1 To 3 Percent Slopes-----	27,206	4.7
To	Tobin Silt Loam, Occasionally Flooded-----	24,628	4.3
W	Water-----	926	0.2
Wb	Wells Loam, 1 To 3 Percent Slopes-----	5,628	1.0
Wc	Wells Loam, 3 To 6 Percent Slopes-----	6,455	1.1
	Total-----	576,339	100.0

* Less than 0.1 percent.

Nontechnical Soil Descriptions
McPherson County, Kansas

Nontechnical soil descriptions describe soil properties or management considerations specific to a soil map unit or group of map units, shown in the NonTechnical Descriptions report. These descriptions are written in terminology that Non-technical users of soil survey information can understand. Nontechnical soil descriptions are a powerful tool for creating reports. These high quality, easy to read reports can be generated by conservation planners and other NRCS employees for distribution to land users. Soil map unit descriptions and National Soil Information System records are the basis for these descriptions.

053ED Edalgo Loam, 3 To 7 Percent Slopes

Edalgo soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is very slow. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Clay Upland (pe26-30) range site. It is in the nonirrigated land capability classification 4e.

053MB Mccook Loam, Occasionally Flooded

Mccook soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Lowland (pe26-30) range site. It is in the nonirrigated land capability classification 2w.

053MC Mccook Silt Loam, Rarely Flooded

Mccook soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Terrace (pe26-30) range site. It is in the nonirrigated land capability classification 1.

079GE Geary Silt Loam, 3 To 6 Percent Slopes

Geary soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping upland, hillslope. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe25-34) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

079LB Ladysmith Silty Clay Loam, 1 To 2 Percent Slopes

Ladysmith soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on upland. The runoff class is low. The parent material consists of clayey alluvium. This soil is somewhat poorly drained. The slowest permeability is impermeable. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Clay Upland (pe25-34) range site. It is in the nonirrigated land capability classification 3e.

115IC Irwin Silty Clay Loam, 3 To 6 Percent Slopes

Irwin soil makes up 90 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is very high. The parent material consists of old clayey alluvium. This soil is moderately well drained. The slowest permeability is very slow. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Clay Upland (pe25-34) range site. It is in the nonirrigated land capability classification 4e.

159DP Dillwyn-Plevna Loamy Fine Sands, 0 To 1 Percent Slopes

Dillwyn soil makes up 60 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level interdune on paleoterrace on river valley, dune on paleoterrace on river valley. The runoff class is negligible. The parent material consists of sandy eolian deposits. This soil is somewhat poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 24 inches. This soil is in the Subirrigated (pe21-28) range site. It is in the nonirrigated land capability classification 4w.

Plevna soil makes up 40 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is poorly drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 12 inches. This soil is in the Subirrigated (pe21-28) range site. It is in the nonirrigated land capability classification 5w.

Nontechnical Soil Descriptions--Continued
McPherson County, Kansas

159GC Geary-Clark Complex, 3 To 7 Percent Slopes, Eroded

Geary soil makes up 75 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping upland, hillslope. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe25-34) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Clark soil makes up 25 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping paleoterrace on river valley. The runoff class is medium. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 45 percent calcium carbonate. This soil is in the Limy Upland (pe25-34) range site. It is in the nonirrigated land capability classification 4e.

159LA Lancaster Loam, 1 To 3 Percent Slopes

Lancaster soil makes up 90 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of loamy residuum weathered from sandstone and shale. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a moderate available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe26-30) range site. It is in the nonirrigated land capability classification 3e.

159NF Naron Fine Sandy Loam, 1 To 3 Percent Slopes

Naron soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy eolian deposits. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe21-28) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 3e.

159PF Pratt Loamy Fine Sand, 1 To 5 Percent Slopes

Pratt soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe21-28) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

159PR Pratt-Carwile Complex, 0 To 5 Percent Slopes

Pratt soil makes up 65 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping dune on paleoterrace on river valley. The runoff class is low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe21-28) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Carwile soil makes up 35 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level depression on paleoterrace on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is somewhat poorly drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. This soil is in the Sandy (pe21-28) range site. It is in the nonirrigated land capability classification 2w.

159SO Smolan Soils, 2 To 7 Percent Slopes, Eroded

Smolan soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping upland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe25-34) range site. It is in the nonirrigated land capability classification 3e.

159WA Waldeck Fine Sandy Loam, Occasionally Flooded

Waldeck soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 36 inches. This soil is in the Subirrigated (pe21-28) range site. It is in the nonirrigated land capability classification 3w.

Nontechnical Soil Descriptions--Continued
McPherson County, Kansas

169CE Clime Silty Clay Loam, 2 To 6 Percent Slopes

Clime soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of silty and clayey residuum weathered from shale, calcareous. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is slow. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Limy Upland (pe25-34) range site. It is in the nonirrigated land capability classification 4e.

169ED Edalgo Clay Loam, 3 To 7 Percent Slopes

Edalgo soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is very high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is impermeable. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Clay Upland (pe26-30) range site. It is in the nonirrigated land capability classification 4e.

169RO Roxbury Silt Loam, Rarely Flooded

Roxbury soil makes up 99 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of calcareous fine-silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Terrace (pe26-30) range site. It is in the nonirrigated land capability classification 1.

1191 Blazefork Silty Clay Loam, 0 To 1 Percent Slopes, Rarely Flooded

Blazefork soil makes up 90 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is very low. The parent material consists of silty alluvium. This soil is moderately well drained. The slowest permeability is slow. It has a high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 48 inches. This soil is in the Clay Lowland (pe25-34) range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2w.

1200 Buhler-Blazefork Silty Clay Loams, 0 To 1 Percent Slopes, Rarely Flooded

Buhler soil makes up 65 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is very low. The parent material consists of alluvium. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a high available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 60 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil contains a slightly saline horizon, it has a horizon that is strongly sodic. This soil is in the Saline Subirrigated (pe21-28) range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Blazefork soil makes up 30 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is very low. The parent material consists of silty alluvium. This soil is moderately well drained. The slowest permeability is slow. It has a high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 48 inches. This soil is in the Clay Lowland (pe25-34) range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2s.

1324 Carway And Carbika Soils, 0 To 1 Percent Slopes

Carway soil makes up 50 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level interdune on depression on paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy eolian deposits over alluvium. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is frequent ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Subirrigated (pe21-28) range site. It is in the nonirrigated land capability classification 2w.

Carbika soil makes up 30 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level interdune on depression on paleoterrace on river valley. The runoff class is very low. The parent material consists of loamy eolian deposits over alluvium. This soil is poorly drained. The slowest permeability is very slow. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is frequent ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Subirrigated (pe21-28) range site. It is in the nonirrigated land capability classification 2w.

Nontechnical Soil Descriptions--Continued
McPherson County, Kansas

1553 Darlow-Elmer Complex, 0 To 2 Percent Slopes

Darlow soil makes up 70 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping terrace on river valley. The runoff class is very low. The parent material consists of loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 2 percent calcium carbonate. This soil contains a moderately saline horizon, it has a horizon that is strongly sodic. This soil is in the Clay Pan (pe21-28) range site. This soil is in the irrigated land capability class 4s. It is in the nonirrigated land capability classification 4s.

Elmer soil makes up 20 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping terrace on river valley. The runoff class is very low. The parent material consists of loamy alluvium. This soil is moderately well drained. The slowest permeability is slow. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 2 percent calcium carbonate. This soil contains a very slightly saline horizon, it has a horizon that is strongly sodic. This soil is in the Loamy Terrace (pe21-28) range site. This soil is in the irrigated land capability class 3s. It is in the nonirrigated land capability classification 3s.

1555 Dillhut-Plev Complex, 0 To 2 Percent Slopes

Dillhut soil makes up 35 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping terrace on river valley. The runoff class is very low. The parent material consists of eolian deposits over alluvium. This soil is moderately well drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe21-28) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Plev soil makes up 35 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level depression on paleoterrace on river valley, interdune on paleoterrace on river valley. The runoff class is very low. The parent material consists of sandy eolian deposits over loamy alluvium. This soil is poorly drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. This soil is in the Subirrigated (pe21-28) range site. It is in the nonirrigated land capability classification 5w.

1985 Hayes Fine Sandy Loam, 1 To 5 Percent Slopes

Hayes soil makes up 60 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping dune on paleoterrace on river valley. The runoff class is low. The parent material consists of loamy eolian deposits over clayey alluvium. This soil is well drained. The slowest permeability is slow. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Sandy (pe21-28) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

2391 Kaskan Silty Clay Loam, 0 To 1 Percent Slopes, Frequently Flooded, Channeled

Kaskan soil makes up 75 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is very low. The parent material consists of loamy alluvium. This soil is well drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 60 inches. The soil contains a maximum amount of 1 percent calcium carbonate. This soil is in the Loamy Lowland (pe21-28) range site. It is in the nonirrigated land capability classification 5w.

2588 Longford Silty Clay Loam, 3 To 7 Percent Slopes, Moderately Eroded

Longford, Moderately Eroded, soil makes up 90 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of silty alluvium or loess. This soil is well drained. The slowest permeability is very slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe25-34) range site. It is in the nonirrigated land capability classification 4e.

3181 Pratt-Turon Fine Sands, 1 To 5 Percent Slopes

Pratt soil makes up 45 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping dune on paleoterrace on river valley. The runoff class is low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe21-28) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Nontechnical Soil Descriptions--Continued
McPherson County, Kansas

Turon soil makes up 30 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping dune on paleoterrace on river valley. The runoff class is low. The parent material consists of sandy eolian deposits over alluvium. This soil is well drained. The slowest permeability is very slow. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe21-28) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

3641 Tivin-Dillhut Fine Sands, 0 To 15 Percent Slopes

Tivin soil makes up 45 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately steep dune on paleoterrace on river valley. The runoff class is medium. The parent material consists of sandy eolian deposits. This soil is somewhat excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Choppy Sands (pe21-28) range site. It is in the nonirrigated land capability classification 6e.

Dillhut soil makes up 40 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level to moderately sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of eolian deposits over alluvium. This soil is moderately well drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe21-28) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

At Attica Loamy Fine Sand, 1 To 4 Percent Slopes

Attica soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping dune on paleoterrace on river valley. The runoff class is very low. The parent material consists of eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy (pe25-34) range site. It is in the nonirrigated land capability classification 2e.

Br Bridgeport Silt Loam, Rarely Flooded

Bridgeport soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on alluvial plain. The runoff class is negligible. The parent material consists of silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Terrace (pe26-30) range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 1.

Ca Carwile Fine Sandy Loam, 0 To 1 Percent Slopes

Carwile soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level depression on paleoterrace on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is somewhat poorly drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. This soil is in the Sandy (pe25-34) range site. It is in the nonirrigated land capability classification 2w.

Cb Cass Fine Sandy Loam, Rarely Flooded

Cass soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a nearly level flood plain. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderately rapid. It has a high available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland (pe26-30) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Ce Clime Silty Clay, 1 To 3 Percent Slopes

Clime soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is high. The parent material consists of silty and clayey residuum weathered from shale, calcareous. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is slow. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland (pe26-30) range site. It is in the nonirrigated land capability classification 3e.

Nontechnical Soil Descriptions--Continued
McPherson County, Kansas

Cm Clime Silty Clay, 3 To 6 Percent Slopes

Clime soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is very low. The parent material consists of silty and clayey residuum weathered from shale, calcareous. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is slow. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland (pe26-30) range site. It is in the nonirrigated land capability classification 4e.

Cr Crete Silt Loam, 0 To 1 Percent Slopes

Crete soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level upland. The runoff class is medium. The parent material consists of silty and clayey loess. This soil is moderately well drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Clay Upland (pe25-34) range site. This soil is in the irrigated land capability class 2s. It is in the nonirrigated land capability classification 2s.

Cs Crete Silt Loam, 1 To 3 Percent Slopes

Crete soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is very high. The parent material consists of silty and clayey loess. This soil is moderately well drained. The slowest permeability is impermeable. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Clay Upland (pe25-34) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Ct Crete Silty Clay Loam, 1 To 3 Percent Slopes, Eroded

Crete soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping upland. The runoff class is very high. The parent material consists of silty and clayey loess. This soil is well drained. The slowest permeability is impermeable. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Clay Upland (pe25-34) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

De Detroit Silty Clay Loam, Rarely Flooded

Detroit soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is very low. The parent material consists of alluvium. This soil is moderately well drained. The slowest permeability is slow. It has a high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Terrace (pe26-30) range site. It is in the nonirrigated land capability classification 1.

Dr Drummond Loam, 0 To 1 Percent Slopes

Drummond soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level terrace on river valley. The runoff class is negligible. The parent material consists of clayey and/or loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 48 inches. This soil contains a slightly saline horizon. This soil is in the Saline Lowland (pe25-34) range site. It is in the nonirrigated land capability classification 6s.

Ed Edalgo Silt Loam, 5 To 12 Percent Slopes

Edalgo soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hillslope on upland. The runoff class is very high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is impermeable. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Clay Upland (pe26-30) range site. It is in the nonirrigated land capability classification 4e.

Ee Edalgo Silty Clay Loam, 3 To 9 Percent Slopes, Eroded

Edalgo soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping upland. The runoff class is very high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is impermeable. It has a low available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Clay Upland (pe26-30) range site. It is in the nonirrigated land capability classification 4e.

Nontechnical Soil Descriptions--Continued
McPherson County, Kansas

Fa Farnum Loam, 1 To 3 Percent Slopes

Farnum soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on river valley. The runoff class is low. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe25-34) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Ge Geary Silt Loam, 1 To 3 Percent Slopes

Geary soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a gently sloping hillslope, upland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe25-34) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Go Goessel Silty Clay, 0 To 2 Percent Slopes

Goessel soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level paleoterrace on upland. The runoff class is low. The parent material consists of clayey alluvium. This soil is moderately well drained. The slowest permeability is very slow. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 30 inches. This soil is in the Clay Upland (pe25-34) range site. It is in the nonirrigated land capability classification 2s.

Ho Hord Silt Loam, Rarely Flooded

Hord soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a nearly level hillslope on upland. The runoff class is negligible. The parent material consists of loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Loamy Terrace (pe26-30) range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 1.

INT Aquolls

Aquolls soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level depression on terrace on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is very poorly drained. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is occasional ponded. The top of the seasonal high water table is at 0 inches. It is in the nonirrigated land capability classification 5w.

Ir Irwin Silty Clay Loam, 1 To 3 Percent Slopes

Irwin, bedrock substratum, soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping paleoterrace on upland. The runoff class is low. The parent material consists of residuum. The soil is 40 to 60 inches deep to bedrock (paralithic). This soil is moderately well drained. The slowest permeability is very slow. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Clay Upland (pe26-30) range site. It is in the nonirrigated land capability classification 3e.

La Ladysmith Silty Clay Loam, 0 To 1 Percent Slopes

Ladysmith soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level paleoterrace on upland. The runoff class is negligible. The parent material consists of clayey alluvium. This soil is somewhat poorly drained. The slowest permeability is impermeable. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Clay Upland (pe25-34) range site. It is in the nonirrigated land capability classification 2s.

Ld Ladysmith-Drummond Complex, 0 To 1 Percent Slopes

Ladysmith soil makes up 70 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level paleoterrace on upland. The runoff class is negligible. The parent material consists of clayey alluvium. This soil is somewhat poorly drained. The slowest permeability is impermeable. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Clay Upland (pe25-34) range site. It is in the nonirrigated land capability classification 4s.

Drummond soil makes up 30 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level terrace on river valley. The runoff class is negligible. The parent material consists of clayey and/or loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 48 inches. This soil contains a slightly saline horizon. This soil is in the Saline Lowland (pe25-34) range site. It is in the nonirrigated land capability classification 6s.

Nontechnical Soil Descriptions--Continued
McPherson County, Kansas

Le Lancaster Loam, 2 To 6 Percent Slopes

Lancaster soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe26-30) range site. It is in the nonirrigated land capability classification 4e.

Lh Lancaster-Hedville Loams, 6 To 12 Percent Slopes

Lancaster soil makes up 60 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hillslope on upland. The runoff class is very high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe26-30) range site. It is in the nonirrigated land capability classification 6e.

Hedville soil makes up 40 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping backslope hillslope on upland. The runoff class is very high. The parent material consists of loamy residuum weathered from sandstone and shale. The soil is 4 to 20 inches deep to bedrock (lithic). This soil is somewhat excessively drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow Sandstone (pe26-30) range site. It is in the nonirrigated land capability classification 7s.

LHH Lancaster-Hedville Complex, 3 To 20 Percent Slopes

Lancaster soil makes up 60 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hillslope on upland. The runoff class is very high. The parent material consists of residuum. The soil is 20 to 40 inches deep to bedrock (paralithic). This soil is well drained. The slowest permeability is moderate. It has a low available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe26-30) range site. It is in the nonirrigated land capability classification 6e.

Hedville soil makes up 40 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a moderately sloping to moderately steep backslope hillslope on upland. The runoff class is very high. The parent material consists of loamy residuum weathered from sandstone and shale. The soil is 4 to 20 inches deep to bedrock (lithic). This soil is somewhat excessively drained. The slowest permeability is moderate. It has a very low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow Sandstone (pe26-30) range site. It is in the nonirrigated land capability classification 6e.

Ln Longford Silty Clay Loam, 3 To 6 Percent Slopes

Longford soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of silty alluvium or loess. This soil is well drained. The slowest permeability is slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe26-30) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Lo Longford Silty Clay Loam, 2 To 6 Percent Slopes, Eroded

Longford soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a gently sloping to moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of silty alluvium or loess. This soil is well drained. The slowest permeability is impermeable. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe26-30) range site. It is in the nonirrigated land capability classification 4e.

Mc Mccook Fine Sandy Loam, Rarely Flooded

Mccook soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Terrace (pe26-30) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Nontechnical Soil Descriptions--Continued
McPherson County, Kansas

Ns Ness Silty Clay

Ness soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level playa. The runoff class is high. The parent material consists of alluvium. This soil is poorly drained. The slowest permeability is very slow. It has a moderate available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. This soil is in the Clay Lowland (pe25-34) range site. It is in the nonirrigated land capability classification 6w.

Nw New Cambria Silty Clay, Rarely Flooded

New Cambria soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a nearly level stream terrace on river valley. The runoff class is medium. The parent material consists of alluvium. This soil is moderately well drained. The slowest permeability is impermeable. It has a moderate available water capacity and a high shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Clay Terrace (pe20-26) range site. It is in the nonirrigated land capability classification 2s.

Pa Plevna Fine Sandy Loam, Frequently Flooded

Plevna soil makes up 100 percent of the map unit. This map unit is in the Great Bend Sand Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of alluvium. This soil is poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 12 inches. This soil is in the Subirrigated (pe25-34) range site. It is in the nonirrigated land capability classification 5w.

Pr Pratt Loamy Fine Sand, 6 To 12 Percent Slopes

Pratt soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping dune on paleoterrace on river valley. The runoff class is low. The parent material consists of sandy eolian deposits. This soil is well drained. The slowest permeability is rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands (pe25-34) range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

Ro Roxbury Silty Clay Loam, Rarely Flooded

Roxbury soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of calcareous fine-silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Loamy Terrace (pe26-30) range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 1.

Sm Smolan Silty Clay Loam, 1 To 3 Percent Slopes

Smolan soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe25-34) range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

To Tobin Silt Loam, Occasionally Flooded

Tobin soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on upland. The runoff class is negligible. The parent material consists of silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Lowland (pe26-30) range site. It is in the nonirrigated land capability classification 2w.

Wb Wells Loam, 1 To 3 Percent Slopes

Wells soil makes up 90 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of fine-loamy residuum weathered from sandstone. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe25-34) range site. It is in the nonirrigated land capability classification 2e.

Nontechnical Soil Descriptions--Continued
McPherson County, Kansas

Wc Wells Loam, 3 To 6 Percent Slopes

Wells soil makes up 100 percent of the map unit. This map unit is in the Central Kansas Sandstone Hills Major Land Resource Area. This soil occurs on a moderately sloping backslope hillslope on upland. The runoff class is low. The parent material consists of fine-loamy residuum. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Loamy Upland (pe26-30) range site. It is in the nonirrigated land capability classification 3e.

053ED—Edalgo loam, 3 to 7 percent slopes**Map Unit Composition**

Edalgo: 100 percent

Component Descriptions**Edalgo**

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Hillslope on upland

Parent material: Residuum

Slope: 3 to 7 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Low (About 5.2 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Clay Upland (pe26-30)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 8 inches; loam

H2—8 to 34 inches; clay

Cr—34 to 34 inches; weathered bedrock

Available water capacity: High (About 11.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Loamy Lowland (pe26-30)

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 12 inches; loam

H2—12 to 60 inches; silt loam

053MC—Mccook silt loam, rarely flooded**Map Unit Composition**

Mccook: 100 percent

Component Descriptions**Mccook**

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Flood plain on river valley

Parent material: Alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Loamy Terrace (pe26-30)

Land capability (nonirrigated): 1

Typical Profile:

H1—0 to 12 inches; silt loam

H2—12 to 60 inches; silt loam

053MB—Mccook loam, occasionally flooded**Map Unit Composition**

Mccook: 100 percent

Component Descriptions**Mccook**

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Flood plain on river valley

Parent material: Alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

079GE—Geary silt loam, 3 to 6 percent slopes**Map Unit Composition**

Geary: 100 percent

Component Descriptions

Geary

MLRA: 75 - Central Loess Plains

Landform: Upland, hillslope

Parent material: Loess

Slope: 3 to 6 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.1 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Loamy Upland (pe25-34)

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 9 inches; silt loam

H2—9 to 35 inches; silty clay loam

H3—35 to 60 inches; clay loam

079LB—Ladysmith silty clay loam, 1 to 2 percent slopes

Map Unit Composition

Ladysmith: 100 percent

Component Descriptions

Ladysmith

MLRA: 75 - Central Loess Plains

Landform: Paleoterrace on upland

Parent material: Clayey alluvium

Slope: 1 to 2 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Impermeable (About 0.00 in/hr)

Available water capacity: Moderate (About 8.8 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Clay Upland (pe25-34)

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 10 inches; silty clay loam

H2—10 to 45 inches; silty clay

H3—45 to 60 inches; silty clay loam

Minor Components

Unnamed Hydric Soils

Unnamed Hydric Soils

1151C—Irwin silty clay loam, 3 to 6 percent slopes

Map Unit Composition

Irwin: 90 percent

Minor components: 10 percent

Component Descriptions

Irwin

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Old clayey alluvium

Slope: 3 to 6 percent

Drainage class: Moderately well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Moderate (About 8.8 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Clay Upland (pe25-34)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 13 inches; silty clay loam

H2—13 to 40 inches; silty clay

H3—40 to 60 inches; silty clay

Minor Components

Clime

Composition: About 5 percent

Slope: 3 to 7 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Ecological site: Limy Upland (pe25-34)

Labette

Composition: About 5 percent

Slope: 1 to 4 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Drainage class: Well drained
Ecological site: Loamy Upland (pe25-34)

H3—33 to 60 inches; fine sand

159DP—Dillwyn-Plevna loamy fine sands, 0 to 1 percent slopes

Map Unit Composition

Dillwyn: 60 percent
 Plevna: 40 percent

Component Descriptions

Dillwyn

MLRA: 79 - Great Bend Sand Plains
Landform: Interdune on paleoterrace on river valley, dune on paleoterrace on river valley
Parent material: Sandy eolian deposits
Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 4.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: About 12 to 36 inches
Runoff class: Negligible
Ecological site: Subirrigated (pe21-28)
Land capability (nonirrigated): 4w

Typical Profile:

H1—0 to 9 inches; loamy fine sand
 H2—9 to 60 inches; loamy fine sand

Plevna

MLRA: 79 - Great Bend Sand Plains
Landform: Flood plain on river valley
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Poorly drained
Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: Low (About 5.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: About 0 to 24 inches
Runoff class: Negligible
Ecological site: Subirrigated (pe21-28)
Land capability (nonirrigated): 5w

Typical Profile:

H1—0 to 10 inches; loamy fine sand
 H2—10 to 33 inches; fine sandy loam

159GC—Geary-Clark complex, 3 to 7 percent slopes, eroded

Map Unit Composition

Geary: 75 percent
 Clark: 25 percent

Component Descriptions

Geary

MLRA: 75 - Central Loess Plains
Landform: Upland, hillslope
Parent material: Loess
Slope: 3 to 7 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.0 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Loamy Upland (pe25-34)
Land capability (irrigated): 3e
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 7 inches; silt loam
 H2—7 to 32 inches; silty clay loam
 H3—32 to 60 inches; silty clay loam

Clark

MLRA: 75 - Central Loess Plains
Landform: Paleoterrace on river valley
Parent material: Loamy alluvium
Slope: 3 to 7 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.4 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Limy Upland (pe25-34)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 11 inches; clay loam
H2—11 to 60 inches; clay loam

Irwin

Composition: About 5 percent

Slope: 1 to 3 percent

Drainage class: Moderately well drained

Ecological site: Clay Upland (pe25-34)

159LA—Lancaster loam, 1 to 3 percent slopes

Map Unit Composition

Lancaster: 90 percent

Minor components: 10 percent

Component Descriptions

Lancaster

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Hillslope on upland

Parent material: Loamy residuum weathered from sandstone and shale

Slope: 1 to 3 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Moderate (About 6.2 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Loamy Upland (pe26-30)

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 7 inches; loam
H2—7 to 25 inches; sandy clay loam
H3—25 to 35 inches; clay loam
Cr—35 to 35 inches; weathered bedrock

Minor Components

Hedville

Composition: About 5 percent

Geomorphic Position: hillslope on upland

Slope: 1 to 3 percent

Depth to restrictive feature: 4 to 20 inches to bedrock (lithic)

Drainage class: Somewhat excessively drained

Ecological site: Shallow Sandstone (pe26-30)

159NF—Naron fine sandy loam, 1 to 3 percent slopes

Map Unit Composition

Naron: 100 percent

Component Descriptions

Naron

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Loamy eolian deposits

Slope: 1 to 3 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 9.1 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sandy (pe21-28)

Land capability (irrigated): 2e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 14 inches; fine sandy loam
H2—14 to 40 inches; sandy clay loam
H3—40 to 60 inches; fine sandy loam

Minor Components

Carwile

Unnamed Wet Soils

Phase: Loamy, Depression

159PF—Pratt loamy fine sand, 1 to 5 percent slopes

Map Unit Composition

Pratt: 100 percent

Component Descriptions

Pratt

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 1 to 5 percent

Drainage class: Well drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Moderate (About 6.4 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sands (pe21-28)

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 13 inches; loamy fine sand

H2—13 to 38 inches;

H3—38 to 60 inches;

Minor Components

Carwile

159PR—Pratt-Carwile complex, 0 to 5 percent slopes

Map Unit Composition

Pratt: 65 percent

Carwile: 35 percent

Component Descriptions

Pratt

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 1 to 5 percent

Drainage class: Well drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Moderate (About 6.4 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sands (pe21-28)

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 13 inches; loamy fine sand

H2—13 to 38 inches; loamy fine sand

H3—38 to 60 inches; fine sand

Carwile

MLRA: 79 - Great Bend Sand Plains

Landform: Depression on paleoterrace on river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: High (About 9.4 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: About 0 to 0 inches

Runoff class: Negligible

Ecological site: Sandy (pe21-28)

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 12 inches; fine sandy loam

H2—12 to 17 inches; sandy clay loam

H3—17 to 60 inches; clay loam

Minor Components

Unnamed Wet Soils

Phase: Sandy, Depression

159SO—Smolan Soils, 2 to 7 percent slopes, eroded

Map Unit Composition

Smolan: 100 percent

Component Descriptions

Smolan

MLRA: 75 - Central Loess Plains

Landform: Upland

Parent material: Loess

Slope: 2 to 7 percent

Drainage class: Well drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: High (About 9.3 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Loamy Upland (pe25-34)

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 6 inches; silty clay loam

H2—6 to 60 inches; silty clay

159WA—Waldeck fine sandy loam, occasionally flooded

Map Unit Composition

Waldeck: 100 percent

Component Descriptions

Waldeck

MLRA: 79 - Great Bend Sand Plains

Landform: Flood plain on river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 7.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: About 24 to 48 inches

Runoff class: Negligible

Ecological site: Subirrigated (pe21-28)

Land capability (nonirrigated): 3w

Typical Profile:

H1—0 to 25 inches; fine sandy loam

H2—25 to 42 inches; fine sandy loam

H3—42 to 60 inches; sand

Minor Components

Unnamed Wet Soils

Phase: Sandy, Depression

169CE—Clime silty clay loam, 2 to 6 percent slopes

Map Unit Composition

Clime: 100 percent

Component Descriptions

Clime

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Silty and clayey residuum weathered from shale, calcareous

Slope: 2 to 6 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: Low (About 4.5 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Limy Upland (pe25-34)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 8 inches; silty clay loam

H2—8 to 20 inches; silty clay

H3—20 to 28 inches; silty clay

Cr—28 to 28 inches; weathered bedrock

169ED—Edalgo clay loam, 3 to 7 percent slopes

Map Unit Composition

Edalgo: 100 percent

Component Descriptions

Edalgo

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Residuum

Slope: 3 to 7 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Impermeable (About 0.00 in/hr)

Available water capacity: Low (About 4.8 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Clay Upland (pe26-30)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 9 inches; clay loam
H2—9 to 13 inches; clay loam
H3—13 to 28 inches; silty clay
Cr—28 to 28 inches; weathered bedrock

Minor Components

Sidehill Seep

169RO—Roxbury silt loam, rarely flooded

Map Unit Composition

Roxbury: 99 percent
Minor components: 1 percent

Component Descriptions

Roxbury

MLRA: 74 - Central Kansas Sandstone Hills
Landform: Flood plain on river valley
Parent material: Calcareous fine-silty alluvium
Slope: 0 to 2 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very high (About 12.6 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: Rare
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Loamy Terrace (pe26-30)
Land capability (nonirrigated): 1

Typical Profile:

H1—0 to 26 inches; silt loam
H2—26 to 60 inches; silt loam

Minor Components

Unnamed Hydric Soil

Composition: About 1 percent
Slope: 0 to 1 percent
Drainage class: Poorly drained

1191—Blazefork silty clay loam, 0 to 1 percent slopes, rarely flooded

Map Unit Composition

Blazefork: 90 percent
Minor components: 10 percent

Component Descriptions

Blazefork

MLRA: 75 - Central Loess Plains
Landform: Flood plain on river valley
Parent material: Silty alluvium
Slope: 0 to 1 percent
Drainage class: Moderately well drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: High (About 9.3 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: Rare
Depth to seasonal water saturation: About 48 to 48 inches
Runoff class: Very low
Ecological site: Clay Lowland (pe25-34)
Land capability (irrigated): 2s
Land capability (nonirrigated): 2w

Typical Profile:

Ap1—0 to 3 inches; silty clay loam
Ap2—3 to 7 inches; silty clay loam
Bt—7 to 14 inches; silty clay
Btss—14 to 22 inches; silty clay
Bt1—22 to 29 inches; silty clay
Bt2—29 to 34 inches; silty clay
Bt3—34 to 40 inches; silty clay
Bt4—40 to 48 inches; silty clay loam
2Bt5—48 to 61 inches; clay loam
2Bt6—61 to 80 inches; loam

Minor Components

Tobin

Composition: About 10 percent
Slope: 0 to 1 percent
Drainage class: Well drained
Ecological site: Loamy Lowland (pe25-34)

Unnamed Wet Soils

General Considerations: This map unit is well suited for the commonly grown crops such as wheat and grain sorghum. Most areas are cropped. The hazard of wind and water erosion is slight. The water table and high shrink-swell potential limit the engineering uses of this soil.

1200—Buhler-Blazefork silty clay loams, 0 to 1 percent slopes, rarely flooded

Map Unit Composition

Buhler: 65 percent
Blazefork: 30 percent
Minor components: 5 percent

Component Descriptions

Buhler

MLRA: 75 - Central Loess Plains
Landform: Flood plain on river valley
Parent material: Alluvium
Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: High (About 10.4 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: Rare
Depth to seasonal water saturation: About 60 to 60 inches
Runoff class: Very low
Ecological site: Saline Subirrigated (pe21-28)
Land capability (irrigated): 2w
Land capability (nonirrigated): 2w

Typical Profile:

Ay—0 to 3 inches; silty clay loam
Anyz—3 to 8 inches; silty clay loam
Eny—8 to 12 inches; silt loam
Bt_{ny}1—12 to 16 inches; silt loam
Bt_{ny}2—16 to 24 inches; clay loam
Bt_{kny}—24 to 36 inches; silty clay loam
Bt_{knyss}—36 to 42 inches; silty clay loam
Bk_{nyss}—42 to 50 inches; clay
2Bk_{ss}—50 to 58 inches; clay loam
2C1—58 to 76 inches; fine sandy loam
2C2—76 to 80 inches; loam

Blazefork

MLRA: 75 - Central Loess Plains
Landform: Flood plain on river valley
Parent material: Silty alluvium
Slope: 0 to 1 percent
Drainage class: Moderately well drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: High (About 9.3 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: Rare
Depth to seasonal water saturation: About 48 to 48 inches

Runoff class: Very low
Ecological site: Clay Lowland (pe25-34)
Land capability (irrigated): 2s
Land capability (nonirrigated): 2s

Typical Profile:

Ap1—0 to 3 inches; silty clay loam
Ap2—3 to 7 inches; silty clay loam
Bt—7 to 14 inches; silty clay
Bt_{ss}—14 to 22 inches; silty clay
Bt1—22 to 29 inches; silty clay
Bt2—29 to 34 inches; silty clay
Bt3—34 to 40 inches; silty clay
Bt4—40 to 48 inches; silty clay loam
2Bt5—48 to 61 inches; clay loam
2Bt6—61 to 80 inches; loam

Minor Components

Tobin

Composition: About 5 percent
Slope: 0 to 1 percent
Drainage class: Well drained
Ecological site: Loamy Lowland (pe25-34)

Unnamed Wet Soils

General Considerations: This map unit is somewhat poorly suited for the commonly grown crops such as wheat and grain sorghum, due to the sodic conditions and soluble salts. Most areas are cropped. The hazard of wind and water erosion is slight. Maintaining soil tilth and soil surface crusting are problems on the Buhler soils, but they can be improved by adding organic matter. The high sodium content, soluble salts, water tables, and high shrink-swell capacity limit most engineering uses, particularly in the area of the Buhler soils.

1324—Carway And Carbika Soils, 0 to 1 percent slopes

Map Unit Composition

Carway: 50 percent
Carbika: 30 percent
Minor components: 20 percent

Component Descriptions

Carway

MLRA: 79 - Great Bend Sand Plains
Landform: Interdune on depression on paleoterrace on river valley
Parent material: Loamy eolian deposits over alluvium
Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: High (About 9.0 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Ponding hazard: Frequent
Depth to seasonal water saturation: About 0 to 0 inches
Runoff class: Very low
Ecological site: Subirrigated (pe21-28)
Land capability (nonirrigated): 2w

Typical Profile:

Ap—0 to 7 inches; fine sandy loam
 Bt1—7 to 10 inches; sandy clay loam
 Bt2—10 to 15 inches; sandy clay loam
 Bt3—15 to 22 inches; fine sandy loam
 Bt4—22 to 35 inches; fine sandy loam
 2Btb1—35 to 40 inches; clay loam
 2Btb2—40 to 54 inches; clay loam
 2Btb3—54 to 63 inches; clay loam
 2Btb4—63 to 72 inches; clay loam
 2Btkb—72 to 80 inches; clay loam

Carbika

MLRA: 79 - Great Bend Sand Plains
Landform: Interdune on depression on paleoterrace on river valley
Parent material: Loamy eolian deposits over alluvium
Slope: 0 to 1 percent
Drainage class: Poorly drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: High (About 9.6 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Ponding hazard: Frequent
Depth to seasonal water saturation: About 0 to 0 inches
Runoff class: Very low
Ecological site: Subirrigated (pe21-28)
Land capability (nonirrigated): 2w

Typical Profile:

A—0 to 11 inches; silt loam
 Bt1—11 to 15 inches; clay
 Bt2—15 to 22 inches; clay loam
 Bt3—22 to 34 inches; clay loam
 Bt4—34 to 41 inches; clay loam
 Bt5—41 to 60 inches; clay loam
 Btk—60 to 80 inches; clay loam

Minor Components

Solvay

Composition: About 20 percent

Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Ecological site: Subirrigated (pe21-28)

General Considerations: Most areas are in cropland but, some are in pasture or range. This mapunit is poorly suited for the most commonly grown crops. Wheat and grain sorghum are the predominant crops. The hazard for wind and water erosion is slight. The presence of water tables and potential for high shrink-swell limit most engineering uses for this mapunit.

1553—Darlow-Elmer complex, 0 to 2 percent slopes

Map Unit Composition

Darlow: 70 percent
 Elmer: 20 percent
 Minor components: 10 percent

Component Descriptions

Darlow

MLRA: 79 - Great Bend Sand Plains
Landform: Terrace on river valley
Parent material: Loamy alluvium
Slope: 0 to 2 percent
Drainage class: Somewhat poorly drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Moderate (About 7.6 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Clay Pan (pe21-28)
Land capability (irrigated): 4s
Land capability (nonirrigated): 4s

Typical Profile:

Ap1—0 to 5 inches; loam
 Ap2—5 to 8 inches; loam
 Btn—8 to 14 inches; loam
 Btny—14 to 20 inches; clay loam
 Btknyz—20 to 26 inches; loam
 Btnz1—26 to 33 inches; loam
 Btnz2—33 to 44 inches; loam
 Btn1—44 to 53 inches; loam
 Btn2—53 to 68 inches; loam
 2Btn3—68 to 80 inches; sandy loam

Elmer*MLRA:* 79 - Great Bend Sand Plains*Landform:* Terrace on river valley*Parent material:* Loamy alluvium*Slope:* 0 to 2 percent*Drainage class:* Moderately well drained*Slowest permeability:* Slow (About 0.06 in/hr)*Available water capacity:* High (About 9.1 inches)*Shrink-swell potential:* High (About 7.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Very low*Ecological site:* Loamy Terrace (pe21-28)*Land capability (irrigated):* 3s*Land capability (nonirrigated):* 3s*Typical Profile:*

Ap1—0 to 6 inches; fine sandy loam

Ap2—6 to 9 inches; fine sandy loam

AB—9 to 19 inches; fine sandy loam

Btn1—19 to 26 inches; fine sandy loam

Btn2—26 to 37 inches; fine sandy loam

Btnk1—37 to 43 inches; loam

Btnk2—43 to 51 inches; clay loam

Btnk3—51 to 61 inches; fine sandy loam

Btn1'—61 to 72 inches; fine sandy loam

Btn2'—72 to 80 inches; fine sandy loam

Minor Components**Punkin***Composition:* About 10 percent*Slope:* 0 to 2 percent*Drainage class:* Moderately well drained*Ecological site:* Clay Pan (pe21-28)**Carway***Slope:* 0 to 1 percent*Drainage class:* Somewhat poorly drained*Ecological site:* Subirrigated (pe21-28)**Carbika***Slope:* 0 to 1 percent*Drainage class:* Poorly drained*Ecological site:* Subirrigated (pe21-28)

General Considerations: Most areas are in cropland, but some are in pasture or range. This mapunit is moderately well suited for the most commonly grown crops. Wheat and grain sorghum are the major crops. The hazard for wind erosion is severe and water erosion is slight. Maintaining soil tilth and soil crusting are problems but they can be improved by adding organic matter. The high sodium content, pH, and soluble salts can limit the engineering uses of this mapunit.

1555—Dillhut-Plev complex, 0 to 2 percent slopes**Map Unit Composition**

Dillhut: 35 percent

Plev: 35 percent

Minor components: 30 percent

Component Descriptions**Dillhut***MLRA:* 79 - Great Bend Sand Plains*Landform:* Dune on paleoterrace on river valley*Parent material:* Eolian deposits over alluvium*Slope:* 0 to 2 percent*Drainage class:* Moderately well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* Moderate (About 6.0 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Very low*Ecological site:* Sands (pe21-28)*Land capability (irrigated):* 3e*Land capability (nonirrigated):* 3e*Typical Profile:*

A—0 to 4 inches; fine sand

AC—4 to 9 inches; fine sand

C1—9 to 18 inches; fine sand

C2—18 to 26 inches; fine sand

2Btb1—26 to 41 inches; fine sandy loam

2Btb2—41 to 55 inches; fine sandy loam

2BCb1—55 to 65 inches; fine sandy loam

2BCb2—65 to 70 inches; fine sandy loam

2Cg—70 to 80 inches; fine sandy loam

Plev*MLRA:* 79 - Great Bend Sand Plains*Landform:* Depression on paleoterrace on river valley, interdune on paleoterrace on river valley*Parent material:* Sandy eolian deposits over loamy alluvium*Slope:* 0 to 1 percent*Drainage class:* Poorly drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* Low (About 3.9 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None

Depth to seasonal water saturation: About 6 to 6 inches

Runoff class: Very low

Ecological site: Subirrigated (pe21-28)

Land capability (nonirrigated): 5w

Typical Profile:

A1—0 to 4 inches; loamy fine sand
 A2—4 to 12 inches; fine sand
 Cg1—12 to 35 inches; fine sand
 Cg2—35 to 46 inches; fine sand
 2Btgb1—46 to 57 inches; fine sandy loam
 2Btgb2—57 to 75 inches; fine sandy loam
 2BCb—75 to 80 inches; loamy fine sand

Minor Components

Dillwyn

Composition: About 20 percent

Slope: 0 to 2 percent

Drainage class: Somewhat poorly drained

Ecological site: Subirrigated (pe21-28)

Walnut

Composition: About 10 percent

Slope: 0 to 1 percent

Drainage class: Poorly drained

Ecological site: Subirrigated (pe21-28)

General Considerations: Most areas are in pasture or range. This mapunit is poorly suited for most commonly grown crops. The hazard for wind erosion is severe and water erosion is slight. The presence of water tables and sandy textures limits many of the engineering uses of this soil.

1985—Hayes fine sandy loam, 1 to 5 percent slopes

Map Unit Composition

Hayes: 60 percent

Minor components: 40 percent

Component Descriptions

Hayes

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Loamy eolian deposits over clayey alluvium

Slope: 1 to 5 percent

Drainage class: Well drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: Moderate (About 8.1 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sandy (pe21-28)

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

Ap—0 to 8 inches; fine sandy loam
 Bt1—8 to 14 inches; fine sandy loam
 Bt2—14 to 23 inches; fine sandy loam
 Bt3—23 to 34 inches; fine sandy loam
 Bt4—34 to 42 inches; fine sandy loam
 Ab—42 to 47 inches; fine sandy loam
 2Btb1—47 to 56 inches; sandy clay loam
 2Btb2—56 to 69 inches; silty clay
 2Btb3—69 to 80 inches; clay loam

Minor Components

Attica

Composition: About 25 percent

Slope: 1 to 5 percent

Drainage class: Well drained

Ecological site: Sandy (pe21-28)

Saltcreek

Composition: About 15 percent

Slope: 1 to 5 percent

Drainage class: Well drained

Ecological site: Sandy (pe21-28)

General Considerations: Most areas are used for cropland, but some areas are in pasture or range. This mapunit is moderately well suited for most commonly grown crops. Wheat, grain sorghum, and irrigated corn are the predominant crops. The hazard for wind erosion is moderate and water erosion is slight. The high shrink-swell potential may limit some of the engineering uses of the soil.

2391—Kaskan silty clay loam, 0 to 1 percent slopes, Frequently flooded, channeled

Map Unit Composition

Kaskan: 75 percent

Minor components: 25 percent

Component Descriptions

Kaskan

MLRA: 79 - Great Bend Sand Plains

Landform: Flood plain on river valley

Parent material: Loamy alluvium

Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Moderate (About 7.8 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: About 60 to 60 inches
Runoff class: Very low
Ecological site: Loamy Lowland (pe21-28)
Land capability (nonirrigated): 5w

Typical Profile:

A1—0 to 9 inches; silty clay loam
 A2—9 to 13 inches; silty clay loam
 Bw1—13 to 17 inches; fine sandy loam
 Bw2—17 to 21 inches; fine sandy loam
 Bw3—21 to 27 inches; fine sandy loam
 C1—27 to 43 inches; stratified fine sand to loamy fine sand
 C2—43 to 57 inches; stratified fine sand to fine sandy loam
 C3—57 to 80 inches; stratified fine sand to fine sandy loam

Minor Components

Tobin

Composition: About 25 percent
Slope: 0 to 1 percent
Drainage class: Well drained
Ecological site: Loamy Lowland (pe25-34)

Unnamed Wet Soils

General Considerations: Most areas are in pasture or range. This map unit is poorly suited for most commonly grown crops. The hazard for wind and water erosion is slight. The water table and occasional chance of flooding may limit some of the engineering practices.

2588—Longford silty clay loam, 3 to 7 percent slopes, moderately eroded

Map Unit Composition

Longford: 90 percent
 Minor components: 10 percent

Component Descriptions

Longford

MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Silty alluvium or loess
Slope: 3 to 7 percent
Drainage class: Well drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: High (About 10.3 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Loamy Upland (pe25-34)
Land capability (nonirrigated): 4e

Typical Profile:

Ap—0 to 6 inches; silty clay loam
 Bt1—6 to 11 inches; silty clay loam
 Bt2—11 to 28 inches; silty clay
 Bt3—28 to 43 inches; silty clay
 BC1—43 to 60 inches; silty clay loam
 BC2—60 to 80 inches; silty clay loam

Minor Components

Geary

Phase: Moderately Eroded
Composition: About 10 percent
Slope: 3 to 7 percent
Drainage class: Well drained
Ecological site: Loamy Upland (pe25-34)

General Considerations: Most areas are used as cropland, but some are used for pasture or range. This mapunit is moderately well suited to all commonly grown crops. Wheat, grain sorghum, and soybeans are the major crops. The hazard of water erosion is severe and wind erosion is moderate. Ephemeral gully erosion potential is also severe. This problem can be overcome by using a conservation tillage, tall grass barriers, contour farming, terraces and waterways, and residue management. This mapunit is moderately well suited for most engineering uses, but has some limits due to high clay content and potential for shrink-swell.

3181—Pratt-Turon fine sands, 1 to 5 percent slopes

Map Unit Composition

Pratt: 45 percent
 Turon: 30 percent

Minor components: 25 percent

Component Descriptions

Pratt

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 1 to 5 percent

Drainage class: Well drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Moderate (About 6.3 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sands (pe21-28)

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

Ap—0 to 8 inches; fine sand

Bt—8 to 24 inches; loamy fine sand

E&Bt—24 to 64 inches; stratified fine sand to loamy fine sand

C—64 to 80 inches; fine sand

Turon

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits over alluvium

Slope: 1 to 5 percent

Drainage class: Well drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Moderate (About 7.1 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sands (pe21-28)

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

Ap—0 to 8 inches; fine sand

Bt—8 to 28 inches; loamy fine sand

E&Bt—28 to 40 inches; stratified loamy fine sand to fine sandy loam

2Btb1—40 to 58 inches; silty clay

2Btb2—58 to 75 inches; silty clay

2Btb3—75 to 80 inches; silty clay

Minor Components

Hayes

Composition: About 25 percent

Slope: 1 to 5 percent

Drainage class: Well drained

Ecological site: Sandy (pe21-28)

Carway

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Ecological site: Subirrigated (pe21-28)

Warnut

Slope: 0 to 1 percent

Drainage class: Poorly drained

Ecological site: Subirrigated (pe21-28)

General Considerations: Most areas are in cropland, but some are in pasture and range. Some areas are also in the Conservation Reserve Program. This mapunit is somewhat poorly suited for the most commonly grown crops. Wheat, grain sorghum, and irrigated corn are the predominant crops. The hazard for wind erosion is severe and water erosion is slight. Wind erosion can be controlled by plant residue management, conservation tillage, and tall grass barriers. This mapunit is moderately well suited for most engineering practices.

3641—Tivin-Dillhut fine sands, 0 to 15 percent slopes

Map Unit Composition

Tivin: 45 percent

Dillhut: 40 percent

Minor components: 15 percent

Component Descriptions

Tivin

MLRA: 79 - Great Bend Sand Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 1 to 15 percent

Drainage class: Somewhat excessively drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Low (About 3.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Ponding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Choppy Sands (pe21-28)
Land capability (nonirrigated): 6e

Typical Profile:

A—0 to 7 inches; fine sand
 AC—7 to 18 inches; fine sand
 C—18 to 80 inches; fine sand

Dillhut

MLRA: 79 - Great Bend Sand Plains
Landform: Dune on paleoterrace on river valley
Parent material: Eolian deposits over alluvium
Slope: 0 to 7 percent
Drainage class: Moderately well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Moderate (About 6.0 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sands (pe21-28)
Land capability (irrigated): 3e
Land capability (nonirrigated): 3e

Typical Profile:

A—0 to 4 inches; fine sand
 AC—4 to 9 inches; fine sand
 C1—9 to 18 inches; fine sand
 C2—18 to 26 inches; fine sand
 2Btb1—26 to 41 inches; fine sandy loam
 2Btb2—41 to 55 inches; fine sandy loam
 2BCb1—55 to 65 inches; fine sandy loam
 2BCb2—65 to 70 inches; fine sandy loam
 2Cg—70 to 80 inches; fine sandy loam

Minor Components

Solvay

Composition: About 15 percent
Slope: 0 to 2 percent
Drainage class: Somewhat poorly drained
Ecological site: Subirrigated (pe21-28)

Carway

Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Ecological site: Subirrigated (pe21-28)

Warnut

Slope: 0 to 1 percent
Drainage class: Poorly drained
Ecological site: Subirrigated (pe21-28)

Plev

Slope: 0 to 1 percent
Drainage class: Poorly drained
Ecological site: Subirrigated (pe21-28)

General Considerations: Most areas are in pasture or range. This mapunit is poorly suited for most commonly grown crops. The hazard for wind erosion is severe and water erosion is slight. This mapunit is poorly suited for most engineering practices.

AED—Arents, Earthen Dam

At—Attica loamy fine sand, 1 to 4 percent slopes

Map Unit Composition

Attica: 100 percent

Component Descriptions

Attica

MLRA: 79 - Great Bend Sand Plains
Landform: Dune on paleoterrace on river valley
Parent material: Eolian deposits
Slope: 1 to 4 percent
Drainage class: Well drained
Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: Moderate (About 7.5 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy (pe25-34)
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 10 inches; loamy fine sand
 H2—10 to 25 inches; fine sandy loam
 H3—25 to 60 inches; loamy fine sand

Minor Components

Carwile

Plevna

Unnamed Wet Soils

Phase: Sandy, Depression

Br—Bridgeport silt loam, rarely flooded**Map Unit Composition**

Bridgeport: 100 percent

Component Descriptions**Bridgeport**

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Flood plain on alluvial plain

Parent material: Silty alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Very high (About 13.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Loamy Terrace (pe26-30)

Land capability (irrigated): 1

Land capability (nonirrigated): 1

Typical Profile:

H1—0 to 14 inches; silt loam

H2—14 to 60 inches; silt loam

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: About 0 to 0 inches

Runoff class: Negligible

Ecological site: Sandy (pe25-34)

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 16 inches; fine sandy loam

H2—16 to 21 inches; loam

H3—21 to 45 inches; clay loam

H4—45 to 60 inches; clay loam

Cb—Cass fine sandy loam, rarely flooded**Map Unit Composition**

Cass: 100 percent

Component Descriptions**Cass**

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Flood plain

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 9.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Sandy Lowland (pe26-30)

Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 7 inches; fine sandy loam

H2—7 to 51 inches; fine sandy loam

H3—51 to 60 inches; loamy fine sand

Ca—Carwile fine sandy loam, 0 to 1 percent slopes**Map Unit Composition**

Carwile: 100 percent

Component Descriptions**Carwile**

MLRA: 75 - Central Loess Plains

Landform: Depression on paleoterrace on river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: High (About 9.4 inches)

Minor Components
Bridgeport

Slope: 0 to 1 percent
Drainage class: Well drained
Ecological site: Overflow

Carwile

Ce—Clime silty clay, 1 to 3 percent slopes

Map Unit Composition

Clime: 100 percent

Component Descriptions

Clime

MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Silty and clayey residuum weathered from shale, calcareous
Slope: 1 to 3 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: Low (About 3.9 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: High
Ecological site: Limy Upland (pe26-30)
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 9 inches; silty clay
H2—9 to 27 inches; silty clay
Cr—27 to 27 inches; unweathered bedrock

Cm—Clime silty clay, 3 to 6 percent slopes

Map Unit Composition

Clime: 100 percent

Component Descriptions

Clime

MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Silty and clayey residuum weathered from shale, calcareous
Slope: 3 to 6 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: Low (About 3.9 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Limy Upland (pe26-30)
Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 9 inches; silty clay
H2—9 to 27 inches; silty clay
Cr—27 to 27 inches; unweathered bedrock

Cr—Crete silt loam, 0 to 1 percent slopes

Map Unit Composition

Crete: 100 percent

Component Descriptions

Crete

MLRA: 75 - Central Loess Plains
Landform: Upland
Parent material: Silty and clayey loess
Slope: 0 to 1 percent
Drainage class: Moderately well drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: High (About 10.9 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Clay Upland (pe25-34)
Land capability (irrigated): 2s
Land capability (nonirrigated): 2s

Typical Profile:

Ap—0 to 5 inches; silt loam
BA—5 to 9 inches; silty clay loam

Bt1—9 to 19 inches; silty clay loam
 Bt2—19 to 27 inches; silty clay
 Bt3—27 to 38 inches; silty clay
 BC—38 to 48 inches; silty clay loam
 C—48 to 80 inches; silty clay loam

Minor Components**Unnamed Wet Soils**

Phase: Clayey, Drainageway

Unnamed Wet Soils

Phase: Clayey, Depression

General Considerations: Most areas are used as cropland. This mapunit is well suited to all commonly grown crops. Wheat, grain sorghum, and soybeans are the major crops. A few areas are planted to irrigated corn. The hazard of water erosion is slight and wind erosion is moderate. This problem can be overcome by using a conservation tillage and residue management. This mapunit is moderately well suited for most engineering uses.

Cs—Crete silt loam, 1 to 3 percent slopes**Map Unit Composition**

Crete: 100 percent

Component Descriptions**Crete**

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Silty and clayey loess

Slope: 1 to 3 percent

Drainage class: Moderately well drained

Slowest permeability: Impermeable (About 0.00 in/hr)

Available water capacity: High (About 10.9 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Clay Upland (pe25-34)

Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 6 inches; silt loam

H2—6 to 13 inches; silty clay loam
 H3—13 to 43 inches; silty clay
 H4—43 to 60 inches; silty clay loam

Minor Components**Unnamed Wet Soils**

Phase: Clayey, Drainageway

Ct—Crete silty clay loam, 1 to 3 percent slopes, eroded**Map Unit Composition**

Crete: 100 percent

Component Descriptions**Crete**

MLRA: 75 - Central Loess Plains

Landform: Upland

Parent material: Silty and clayey loess

Slope: 1 to 3 percent

Drainage class: Well drained

Slowest permeability: Impermeable (About 0.00 in/hr)

Available water capacity: High (About 11.2 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Clay Upland (pe25-34)

Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 6 inches; silty clay loam

H2—6 to 13 inches; silty clay loam

H3—13 to 35 inches; silty clay

H4—35 to 60 inches; silty clay loam

De—Detroit silty clay loam, rarely flooded**Map Unit Composition**

Detroit: 100 percent

Component Descriptions

Detroit*MLRA:* 74 - Central Kansas Sandstone Hills*Landform:* Flood plain on river valley*Parent material:* Alluvium*Slope:* 0 to 1 percent*Drainage class:* Moderately well drained*Slowest permeability:* Slow (About 0.06 in/hr)*Available water capacity:* High (About 11.1 inches)*Shrink-swell potential:* High (About 7.5 LEP)*Flooding hazard:* Rare*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Very low*Ecological site:* Loamy Terrace (pe26-30)*Land capability (nonirrigated):* 1*Typical Profile:*

Ap—0 to 16 inches; silty clay loam

H2—16 to 37 inches; silty clay loam

H3—37 to 60 inches; silt loam

Minor Components**Unnamed Hydric Soils****Dr—Drummond loam, 0 to 1 percent slopes****Map Unit Composition**

Drummond: 100 percent

Component Descriptions**Drummond***MLRA:* 75 - Central Loess Plains*Landform:* Terrace on river valley*Parent material:* Clayey and/or loamy alluvium*Slope:* 0 to 1 percent*Drainage class:* Somewhat poorly drained*Slowest permeability:* Very slow (About 0.00 in/hr)*Available water capacity:* Moderate (About 8.1 inches)*Shrink-swell potential:* High (About 7.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* About 24 to 72 inches*Runoff class:* Negligible*Ecological site:* Saline Lowland (pe25-34)*Land capability (nonirrigated):* 6s*Typical Profile:*

H1—0 to 19 inches; loam

H2—19 to 60 inches; clay loam

Minor Components**Carwile****Unnamed Wet Soils***Phase:* Clayey, Depression**Ed—Edalgo silt loam, 5 to 12 percent slopes****Map Unit Composition**

Edalgo: 100 percent

Component Descriptions**Edalgo***MLRA:* 74 - Central Kansas Sandstone Hills*Landform:* Hillslope on upland*Parent material:* Residuum*Slope:* 5 to 12 percent*Depth to restrictive feature:* 20 to 40 inches to bedrock (paralithic)*Drainage class:* Well drained*Slowest permeability:* Impermeable (About 0.00 in/hr)*Available water capacity:* Low (About 5.1 inches)*Shrink-swell potential:* High (About 7.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Very high*Ecological site:* Clay Upland (pe26-30)*Land capability (nonirrigated):* 4e*Typical Profile:*

H1—0 to 6 inches; silt loam

H2—6 to 15 inches; silty clay loam

H3—15 to 30 inches; silty clay loam

Cr—30 to 30 inches; weathered bedrock

Minor Components**Unnamed Wet Soils***Phase:* Clayey, Drainageway**Clime***Depth to restrictive feature:* 20 to 40 inches to bedrock (paralithic)*Drainage class:* Well drained*Ecological site:* Limy Upland (pe25-34)**Lancaster***Depth to restrictive feature:* 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained
Ecological site: Loamy Upland (pe26-30)

Ee—Edalgo silty clay loam, 3 to 9 percent slopes, eroded

Map Unit Composition

Edalgo: 100 percent

Component Descriptions

Edalgo

MLRA: 74 - Central Kansas Sandstone Hills
Landform: Upland
Parent material: Residuum
Slope: 3 to 9 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Impermeable (About 0.00 in/hr)
Available water capacity: Low (About 3.8 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Clay Upland (pe26-30)
Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 8 inches; silty clay loam
 H2—8 to 23 inches; silty clay loam
 Cr—23 to 23 inches; weathered bedrock

Minor Components

Unnamed Wet Soils

Phase: Clayey, Drainageway

Fa—Farnum loam, 1 to 3 percent slopes

Map Unit Composition

Farnum: 100 percent

Component Descriptions

Farnum

MLRA: 79 - Great Bend Sand Plains
Landform: Paleoterrace on river valley
Parent material: Alluvium
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 10.3 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe25-34)
Land capability (irrigated): 2e
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 12 inches; loam
 H2—12 to 49 inches; sandy clay loam
 H3—49 to 60 inches; fine sandy loam

Ge—Geary silt loam, 1 to 3 percent slopes

Map Unit Composition

Geary: 100 percent

Component Descriptions

Geary

MLRA: 74 - Central Kansas Sandstone Hills
Landform: Hillslope, upland
Parent material: Loess
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.0 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Loamy Upland (pe25-34)
Land capability (irrigated): 2e
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 7 inches; silt loam
 H2—7 to 32 inches; silty clay loam
 H3—32 to 60 inches; silty clay loam

Go—Goessel silty clay, 0 to 2 percent slopes**Map Unit Composition**

Goessel: 100 percent

Component Descriptions**Goessel**

MLRA: 75 - Central Loess Plains
Landform: Paleoterrace on upland
Parent material: Clayey alluvium
Slope: 0 to 1 percent
Drainage class: Moderately well drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Moderate (About 7.7 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: About 24 to 36 inches
Runoff class: Low
Ecological site: Clay Upland (pe25-34)
Land capability (nonirrigated): 2s

Typical Profile:

H1—0 to 14 inches; silty clay
 H2—14 to 50 inches; silty clay
 H3—50 to 60 inches; silty clay

Minor Components**Unnamed Wet Soils**

Phase: Clayey, Depression

Ho—Hord silt loam, rarely flooded**Map Unit Composition**

Hord: 100 percent

Component Descriptions**Hord**

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Hillslope on upland

Parent material: Loess

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.9 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Loamy Terrace (pe26-30)

Land capability (irrigated): 1

Land capability (nonirrigated): 1

Typical Profile:

H1—0 to 6 inches; silt loam
 H2—6 to 43 inches; silty clay loam
 H3—43 to 60 inches; silt loam

Minor Components**Unnamed Wet Soils**

Phase: Loamy, Depression

Unnamed Wet Soils

Phase: Loamy, Drainageway

INT—Aquolls

General Considerations: This map unit was formerly labeled as an Intermittent Water spot symbol. These depressional areas contain soils that are occasionally ponded for long duration.

Ir—Irwin silty clay loam, 1 to 3 percent slopes**Map Unit Composition**

Irwin: 100 percent

Component Descriptions**Irwin**

MLRA: 75 - Central Loess Plains

Landform: Paleoterrace on upland

Parent material: Residuum
Slope: 1 to 3 percent
Depth to restrictive feature: 40 to 60 inches to bedrock (paralithic)
Drainage class: Moderately well drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Moderate (About 6.4 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Clay Upland (pe26-30)
Land capability (nonirrigated): 3e

Typical Profile:
 H1—0 to 11 inches; silty clay loam
 H2—11 to 42 inches; clay
 Cr—42 to 42 inches; unweathered bedrock

Minor Components
Unnamed Wet Soils
Phase: Clayey, Drainageway

La—Ladysmith silty clay loam, 0 to 1 percent slopes

Map Unit Composition

Ladysmith: 100 percent

Component Descriptions

Ladysmith
MLRA: 75 - Central Loess Plains
Landform: Paleoterrace on upland
Parent material: Clayey alluvium
Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Slowest permeability: Impermeable (About 0.00 in/hr)
Available water capacity: Moderate (About 8.6 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Clay Upland (pe25-34)
Land capability (nonirrigated): 2s

Typical Profile:

H1—0 to 8 inches; silty clay loam
 H2—8 to 48 inches; clay
 H3—48 to 60 inches; silty clay

Minor Components
Unnamed Wet Soils
Phase: Clayey, Drainageway

Unnamed Wet Soils
Phase: Clayey, Depression

Ld—Ladysmith-Drummond complex, 0 to 1 percent slopes

Map Unit Composition

Ladysmith: 70 percent
 Drummond: 30 percent

Component Descriptions

Ladysmith
MLRA: 75 - Central Loess Plains
Landform: Paleoterrace on upland
Parent material: Clayey alluvium
Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Slowest permeability: Impermeable (About 0.00 in/hr)
Available water capacity: Moderate (About 8.6 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Clay Upland (pe25-34)
Land capability (nonirrigated): 4s

Typical Profile:
 H1—0 to 8 inches; silty clay loam
 H2—8 to 48 inches; clay
 H3—48 to 60 inches; silty clay

Drummond
MLRA: 75 - Central Loess Plains
Landform: Terrace on river valley
Parent material: Clayey and/or loamy alluvium
Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: Moderate (About 7.8 inches)

Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: About 24 to 72 inches

Runoff class: Negligible
Ecological site: Saline Lowland (pe25-34)
Land capability (nonirrigated): 6s

Typical Profile:
 H1—0 to 7 inches; loam
 H2—7 to 60 inches; clay loam

Minor Components

Unnamed Wet Soils

Phase: Clayey, Depression

Le—Lancaster loam, 2 to 6 percent slopes

Map Unit Composition

Lancaster: 100 percent

Component Descriptions

Lancaster

MLRA: 74 - Central Kansas Sandstone Hills
Landform: Hillslope on upland
Parent material: Residuum
Slope: 2 to 6 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Low (About 5.7 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Loamy Upland (pe26-30)
Land capability (nonirrigated): 4e

Typical Profile:
 H1—0 to 10 inches; loam
 H2—10 to 32 inches; clay loam
 Cr—32 to 32 inches; weathered bedrock

Lh—Lancaster-Hedville loams, 6 to 12 percent slopes

Map Unit Composition

Lancaster: 60 percent
 Hedville: 40 percent

Component Descriptions

Lancaster

MLRA: 74 - Central Kansas Sandstone Hills
Landform: Hillslope on upland
Parent material: Residuum
Slope: 6 to 12 percent
Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Low (About 5.7 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Loamy Upland (pe26-30)
Land capability (nonirrigated): 6e

Typical Profile:
 H1—0 to 10 inches; loam
 H2—10 to 32 inches; clay loam
 Cr—32 to 32 inches; weathered bedrock

Hedville

MLRA: 74 - Central Kansas Sandstone Hills
Landform: Hillslope on upland
Hillslope position: Backslope
Parent material: Loamy residuum weathered from sandstone and shale
Slope: 6 to 12 percent
Depth to restrictive feature: 4 to 20 inches to bedrock (lithic)
Drainage class: Somewhat excessively drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very low (About 2.8 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very high
Ecological site: Shallow Sandstone (pe26-30)

Land capability (nonirrigated): 7s

Typical Profile:

H1—0 to 15 inches; loam
R—15 to 15 inches; unweathered bedrock

LHH—Lancaster-Hedville complex, 3 to 20 percent slopes

Map Unit Composition

Lancaster: 60 percent

Hedville: 40 percent

Component Descriptions

Lancaster

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Hillslope on upland

Parent material: Residuum

Slope: 3 to 12 percent

Depth to restrictive feature: 20 to 40 inches to bedrock (paralithic)

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Low (About 5.4 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Loamy Upland (pe26-30)

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 9 inches; loam
H2—9 to 24 inches; clay loam
H3—24 to 30 inches; sandy clay loam
Cr—30 to 30 inches; weathered bedrock

Hedville

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Hillslope on upland

Hillslope position: Backslope

Parent material: Loamy residuum weathered from sandstone and shale

Slope: 3 to 20 percent

Depth to restrictive feature: 4 to 20 inches to bedrock (lithic)

Drainage class: Somewhat excessively drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Very low (About 2.7 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very high

Ecological site: Shallow Sandstone (pe26-30)

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 8 inches; loam
H2—8 to 17 inches; gravelly loam
R—17 to 17 inches; unweathered bedrock

Minor Components

Sidehill Seep

Ln—Longford silty clay loam, 3 to 6 percent slopes

Map Unit Composition

Longford: 100 percent

Component Descriptions

Longford

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Hillslope on upland

Parent material: Silty alluvium or loess

Slope: 3 to 6 percent

Drainage class: Well drained

Slowest permeability: Slow (About 0.06 in/hr)

Available water capacity: High (About 11.4 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Loamy Upland (pe26-30)

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 15 inches; silty clay loam
H2—15 to 29 inches; silty clay
H4—29 to 47 inches; silty clay loam
H3—47 to 60 inches; silty clay loam

Minor Components

Unnamed Wet Soils

Phase: Clayey, Drainageway

Lo—Longford silty clay loam, 2 to 6 percent slopes, eroded

Map Unit Composition

Longford: 100 percent

Component Descriptions

Longford

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Hillslope on upland

Parent material: Silty alluvium or loess

Slope: 2 to 6 percent

Drainage class: Well drained

Slowest permeability: Impermeable (About 0.00 in/hr)

Available water capacity: High (About 10.5 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Loamy Upland (pe26-30)

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 6 inches; silty clay loam

H2—6 to 15 inches; silty clay loam

H3—15 to 47 inches; silty clay

H4—47 to 60 inches; silty clay loam

Minor Components

Unnamed Wet Soils

Phase: Clayey, Drainageway

Mc—Mccook fine sandy loam, rarely flooded

Map Unit Composition

Mccook: 100 percent

Component Descriptions

Mccook

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Flood plain on river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 10.9 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Sandy Terrace (pe26-30)

Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 14 inches; fine sandy loam

H2—14 to 60 inches; very fine sandy loam

Ns—Ness silty clay

Map Unit Composition

Ness: 95 percent

Minor components: 5 percent

Component Descriptions

Ness

MLRA: 75 - Central Loess Plains

Landform: Playa

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Poorly drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: Moderate (About 7.7 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: About 0 to 0 inches

Runoff class: High

Ecological site: Clay Lowland (pe25-34)

Land capability (nonirrigated): 6w

Typical Profile:

H1—0 to 60 inches; silty clay

Minor Components

Unnamed Wet Soils

Phase: Clayey, Depression

Composition: About 5 percent

Nw—New Cambria silty clay, rarely flooded

Map Unit Composition

New Cambria: 100 percent

Component Descriptions

New Cambria

MLRA: 74 - Central Kansas Sandstone Hills

Landform: Stream terrace on river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Moderately well drained

Slowest permeability: Impermeable (About 0.00 in/hr)

Available water capacity: Moderate (About 8.6 inches)

Shrink-swell potential: High (About 7.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Clay Terrace (pe20-26)

Land capability (nonirrigated): 2s

Typical Profile:

H1—0 to 12 inches; silty clay

H2—12 to 34 inches; silty clay

H3—34 to 60 inches; silty clay

Minor Components

Unnamed Hydric Soils

Unnamed Hydric Soils

Pa—Plevna fine sandy loam, Frequently flooded

Map Unit Composition

Plevna: 100 percent

Component Descriptions

Plevna

MLRA: 79 - Great Bend Sand Plains

Landform: Flood plain on river valley

Parent material: Alluvium

Slope: 0 to 1 percent

Drainage class: Poorly drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 6.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Frequent

Depth to seasonal water saturation: About 0 to 24 inches

Runoff class: Negligible

Ecological site: Subirrigated (pe25-34)

Land capability (nonirrigated): 5w

Typical Profile:

H1—0 to 18 inches; fine sandy loam

H2—18 to 36 inches; fine sandy loam

H3—36 to 60 inches; loamy fine sand

Pr—Pratt loamy fine sand, 6 to 12 percent slopes

Map Unit Composition

Pratt: 100 percent

Component Descriptions

Pratt

MLRA: 75 - Central Loess Plains

Landform: Dune on paleoterrace on river valley

Parent material: Sandy eolian deposits

Slope: 6 to 12 percent

Drainage class: Well drained

Slowest permeability: Rapid (About 5.95 in/hr)

Available water capacity: Moderate (About 6.4 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sands (pe25-34)

Land capability (irrigated): 3e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 11 inches; loamy fine sand

H2—11 to 36 inches; loamy fine sand

H3—36 to 60 inches; loamy fine sand

Minor Components

Carwile

Plevna**Unnamed Wet Soils***Phase: Sandy, Depression***Ro—Roxbury silty clay loam,
rarely flooded****Map Unit Composition**

Roxbury: 100 percent

Component Descriptions**Roxbury***MLRA: 74 - Central Kansas Sandstone Hills**Landform: Flood plain on river valley**Parent material: Calcareous fine-silty alluvium**Slope: 0 to 1 percent**Drainage class: Well drained**Slowest permeability: Moderate (About 0.60 in/hr)**Available water capacity: Very high (About 12.2 inches)**Shrink-swell potential: Moderate (About 4.5 LEP)**Flooding hazard: Rare**Depth to seasonal water saturation: More than 6 feet**Runoff class: Negligible**Ecological site: Loamy Terrace (pe26-30)**Land capability (irrigated): 1**Land capability (nonirrigated): 1**Typical Profile:**H1—0 to 21 inches; silty clay loam**H2—21 to 50 inches; silty clay loam**H3—50 to 60 inches; silt loam***Sm—Smolan silty clay loam, 1 to 3
percent slopes****Map Unit Composition**

Smolan: 100 percent

Component Descriptions**Smolan***MLRA: 75 - Central Loess Plains**Landform: Hillslope on upland**Parent material: Loess**Slope: 1 to 3 percent**Drainage class: Well drained**Slowest permeability: Slow (About 0.06 in/hr)**Available water capacity: High (About 10.0 inches)**Shrink-swell potential: High (About 7.5 LEP)**Flooding hazard: None**Depth to seasonal water saturation: More than 6 feet**Runoff class: Low**Ecological site: Loamy Upland (pe25-34)**Land capability (irrigated): 2e**Land capability (nonirrigated): 2e**Typical Profile:**H1—0 to 11 inches; silty clay loam**H2—11 to 16 inches;**H3—16 to 60 inches;***To—Tobin silt loam, occasionally
flooded****Map Unit Composition**

Tobin: 100 percent

Component Descriptions**Tobin***MLRA: 75 - Central Loess Plains**Landform: Flood plain on upland**Parent material: Silty alluvium**Slope: 0 to 2 percent**Drainage class: Well drained**Slowest permeability: Moderate (About 0.60 in/hr)**Available water capacity: Very high (About 12.1 inches)**Shrink-swell potential: Moderate (About 4.5 LEP)**Flooding hazard: Occasional**Depth to seasonal water saturation: More than 6 feet**Runoff class: Negligible**Ecological site: Loamy Lowland (pe26-30)**Land capability (nonirrigated): 2w**Typical Profile:**H1—0 to 20 inches; silt loam**H2—20 to 32 inches; silt loam**H3—32 to 60 inches; silt loam*

Minor Components**Unnamed Hydric Soils***Slope:* 0 to 2 percent*Drainage class:* Poorly drained**Unnamed Hydric Soil***Slope:* 0 to 2 percent*Drainage class:* Poorly drained**Unnamed Wet Soils***Phase:* Loamy, Depression**Unnamed Wet Soils***Phase:* Loamy, Drainageway**Clime***Composition:* About 5 percent*Slope:* 1 to 3 percent*Depth to restrictive feature:* 20 to 40 inches to bedrock (paralitric)*Drainage class:* Well drained*Ecological site:* Limy Upland (pe25-34)**Irwin***Composition:* About 5 percent*Slope:* 1 to 3 percent*Drainage class:* Moderately well drained*Ecological site:* Clay Upland (pe25-34)**W—Water****Wc—Wells loam, 3 to 6 percent slopes****Wb—Wells loam, 1 to 3 percent slopes****Map Unit Composition**

Wells: 100 percent

Map Unit Composition

Wells: 90 percent

Minor components: 10 percent

Component Descriptions**Wells***MLRA:* 74 - Central Kansas Sandstone Hills*Landform:* Hillslope on upland*Parent material:* Fine-loamy residuum weathered from sandstone*Slope:* 1 to 3 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 10.2 inches)*Shrink-swell potential:* Moderate (About 4.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Low*Ecological site:* Loamy Upland (pe25-34)*Land capability (nonirrigated):* 2e*Typical Profile:*

H1—0 to 15 inches; loam

H2—15 to 36 inches; sandy clay loam

H3—36 to 60 inches; sandy loam

Minor Components**Component Descriptions****Wells***MLRA:* 74 - Central Kansas Sandstone Hills*Landform:* Hillslope on upland*Hillslope position:* Backslope*Parent material:* Fine-loamy residuum*Slope:* 3 to 6 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 10.3 inches)*Shrink-swell potential:* Moderate (About 4.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Low*Ecological site:* Loamy Upland (pe26-30)*Land capability (nonirrigated):* 3e*Typical Profile:*

H1—0 to 11 inches; loam

H2—11 to 15 inches; sandy clay loam

H3—15 to 49 inches; sandy clay loam

H4—49 to 60 inches; sandy loam

Minor Components**Unnamed Wet Soils***Phase:* Loamy, Drainageway

Prime farmland is one of several kinds of important farmland defined by the U.S. Department of Agriculture. It is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil qualities, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. It is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

A recent trend in land use in some parts of the survey area has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

The map units in the survey area that are considered prime farmland are listed in the following table. This list does not constitute a recommendation for a particular land use. On some soils included in the list, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures. The extent of each listed map unit is shown in the "Acres and Proportionate Extent of Soils" table. The location is shown on the detailed soil maps. The soil qualities that affect use and management are described in other tables in this document."

Map symbol	Mapunit name	Farmland Classification
053MB	Mccook loam, occasionally flooded	All areas are prime farmland
053MC	Mccook silt loam, rarely flooded	All areas are prime farmland
079GE	Geary silt loam, 3 to 6 percent slopes	All areas are prime farmland
079LB	Ladysmith silty clay loam, 1 to 2 percent slopes	All areas are prime farmland
115IC	Irwin silty clay loam, 3 to 6 percent slopes	All areas are prime farmland
159GC	Geary-clark complex, 3 to 7 percent slopes, eroded	All areas are prime farmland
159LA	Lancaster loam, 1 to 3 percent slopes	All areas are prime farmland
159NF	Naron fine sandy loam, 1 to 3 percent slopes	All areas are prime farmland
159WA	Waldeck fine sandy loam, occasionally flooded	All areas are prime farmland
169RO	Roxbury silt loam, rarely flooded	All areas are prime farmland
1191	Blazefork silty clay loam, 0 to 1 percent slopes, rarely flooded	All areas are prime farmland
1985	Hayes fine sandy loam, 1 to 5 percent slopes	All areas are prime farmland
Br	Bridgeport silt loam, rarely flooded	All areas are prime farmland
Cb	Cass fine sandy loam, rarely flooded	All areas are prime farmland
Cr	Crete silt loam, 0 to 1 percent slopes	All areas are prime farmland
Cs	Crete silt loam, 1 to 3 percent slopes	All areas are prime farmland
Ct	Crete silty clay loam, 1 to 3 percent slopes, eroded	All areas are prime farmland
De	Detroit silty clay loam, rarely flooded	All areas are prime farmland
Fa	Farnum loam, 1 to 3 percent slopes	All areas are prime farmland
Ge	Geary silt loam, 1 to 3 percent slopes	All areas are prime farmland
Go	Goessel silty clay, 0 to 2 percent slopes	All areas are prime farmland
Ho	Hord silt loam, rarely flooded	All areas are prime farmland
Ir	Irwin silty clay loam, 1 to 3 percent slopes	All areas are prime farmland
La	Ladysmith silty clay loam, 0 to 1 percent slopes	All areas are prime farmland
Le	Lancaster loam, 2 to 6 percent slopes	All areas are prime farmland
Ln	Longford silty clay loam, 3 to 6 percent slopes	All areas are prime farmland
Mc	Mccook fine sandy loam, rarely flooded	All areas are prime farmland
Nw	New cambria silty clay, rarely flooded	All areas are prime farmland
Ro	Roxbury silty clay loam, rarely flooded	All areas are prime farmland
Sm	Smolan silty clay loam, 1 to 3 percent slopes	All areas are prime farmland
To	Tobin silt loam, occasionally flooded	All areas are prime farmland
Wb	Wells loam, 1 to 3 percent slopes	All areas are prime farmland
Wc	Wells loam, 3 to 6 percent slopes	All areas are prime farmland

The "Soil Rating for Plant Growth, modified 1998" (SRPG) is a relative rating of the capacity of a soil to produce a specific plant under a defined management system. The index is determined from yield data on a few benchmark soils and is used to calculate yields, the net returns from crops, land assessment values, and taxes and to perform risk analysis when land management decisions are made. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

Map symbol	Soil name	Crop Index
053ED	Edalgo Loam, 3 To 7 Percent Slopes-----	45
053MB	Mccook Loam, Occasionally Flooded-----	59
053MC	Mccook Silt Loam, Rarely Flooded-----	61
079GE	Geary Silt Loam, 3 To 6 Percent Slopes-----	78
079LB	Ladysmith Silty Clay Loam, 1 To 2 Percent Slopes-----	67
115IC	Irwin Silty Clay Loam, 3 To 6 Percent Slopes-----	63
119I	Blazefork Silty Clay Loam, 0 To 1 Percent Slopes, Rarely Flooded-----	55
1200	Buhler-Blazefork Silty Clay Loams, 0 To 1 Percent Slopes, Rarely Flooded-----	46
1324	Carway And Carbika Soils, 0 To 1 Percent Slopes-----	33
1553	Darlow-Elmer Complex, 0 To 2 Percent Slopes-----	26
1555	Dillhut-Plevna Complex, 0 To 2 Percent Slopes-----	30
159DP	Dillwyn-Plevna Loamy Fine Sands, 0 To 1 Percent Slopes-----	29
159GC	Geary-Clark Complex, 3 To 7 Percent Slopes, Eroded-----	64
159LA	Lancaster Loam, 1 To 3 Percent Slopes-----	61
159NF	Naron Fine Sandy Loam, 1 To 3 Percent Slopes-----	68
159PF	Pratt Loamy Fine Sand, 1 To 5 Percent Slopes-----	40
159PR	Pratt-Carwile Complex, 0 To 5 Percent Slopes-----	33
159SO	Smolan Soils, 2 To 7 Percent Slopes, Eroded-----	66
159WA	Waldeck Fine Sandy Loam, Occasionally Flooded-----	45
169CE	Clime Silty Clay Loam, 2 To 6 Percent Slopes-----	30
169ED	Edalgo Clay Loam, 3 To 7 Percent Slopes-----	38
169RO	Roxbury Silt Loam, Rarely Flooded-----	62
1985	Hayes Fine Sandy Loam, 1 To 5 Percent Slopes-----	57
2391	Kaskan Silty Clay Loam, 0 To 1 Percent Slopes, Frequently Flooded, Channeled-----	54
2588	Longford Silty Clay Loam, 3 To 7 Percent Slopes, Moderately Eroded-----	72
3181	Pratt-Turon Fine Sands, 1 To 5 Percent Slopes-----	43
3641	Tivin-Dillhut Fine Sands, 0 To 15 Percent Slopes-----	33
AED	Arents, Earthen Dam-----	0
At	Attica Loamy Fine Sand, 1 To 4 Percent Slopes-----	51
Br	Bridgeport Silt Loam, Rarely Flooded-----	63
Ca	Carwile Fine Sandy Loam, 0 To 1 Percent Slopes-----	22
Cb	Cass Fine Sandy Loam, Rarely Flooded-----	56
Ce	Clime Silty Clay, 1 To 3 Percent Slopes-----	29
Cm	Clime Silty Clay, 3 To 6 Percent Slopes-----	28
Cr	Crete Silt Loam, 0 To 1 Percent Slopes-----	70
Cs	Crete Silt Loam, 1 To 3 Percent Slopes-----	71
Ct	Crete Silty Clay Loam, 1 To 3 Percent Slopes, Eroded-----	70
De	Detroit Silty Clay Loam, Rarely Flooded-----	71
Dr	Drummond Loam, 0 To 1 Percent Slopes-----	52
Ed	Edalgo Silt Loam, 5 To 12 Percent Slopes-----	38
Ee	Edalgo Silty Clay Loam, 3 To 9 Percent Slopes, Eroded-----	24
Fa	Farnum Loam, 1 To 3 Percent Slopes-----	75
Ge	Geary Silt Loam, 1 To 3 Percent Slopes-----	77
Go	Goessel Silty Clay, 0 To 2 Percent Slopes-----	54
Ho	Hord Silt Loam, Rarely Flooded-----	79
INT	Aquolls-----	12
Ir	Irwin Silty Clay Loam, 1 To 3 Percent Slopes-----	60
LHH	Lancaster-Hedville Complex, 3 To 20 Percent Slopes-----	31
La	Ladysmith Silty Clay Loam, 0 To 1 Percent Slopes-----	66
Ld	Ladysmith-Drummond Complex, 0 To 1 Percent Slopes-----	62
Le	Lancaster Loam, 2 To 6 Percent Slopes-----	52
Lh	Lancaster-Hedville Loams, 6 To 12 Percent Slopes-----	33
Ln	Longford Silty Clay Loam, 3 To 6 Percent Slopes-----	72
Lo	Longford Silty Clay Loam, 2 To 6 Percent Slopes, Eroded-----	73
Mc	Mccook Fine Sandy Loam, Rarely Flooded-----	61
Ns	Ness Silty Clay-----	13
Nw	New Cambria Silty Clay, Rarely Flooded-----	45
Pa	Plevna Fine Sandy Loam, Frequently Flooded-----	31
Pr	Pratt Loamy Fine Sand, 6 To 12 Percent Slopes-----	37
Ro	Roxbury Silty Clay Loam, Rarely Flooded-----	62
Sm	Smolan Silty Clay Loam, 1 To 3 Percent Slopes-----	74
To	Tobin Silt Loam, Occasionally Flooded-----	65
W	Water-----	0
Wb	Wells Loam, 1 To 3 Percent Slopes-----	78
Wc	Wells Loam, 3 To 6 Percent Slopes-----	75

McPherson County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
053ED:EDALGO----	100	N/A	4e	Not prime farmland	C	Clay Upland (pe26-30)	7	.32	.32	3	6	48
053MB:MCCOOK----	100	N/A	2w	All areas are prime farmland	B	Loamy Lowland (pe26-30)	5	.32	.32	5	4L	86
053MC:MCCOOK----	100	N/A	1	All areas are prime farmland	B	Loamy Terrace (pe26-30)	5	.32	.32	5	4L	86
079GE:GEARY-----	100	3e-	3e	All areas are prime farmland	B	Loamy Upland (pe25-34)	7	.32	.32	5	6	48
079LB:LADYSMITH-	100	N/A	3e	All areas are prime farmland	D	Clay Upland (pe25-34)	8	.37	.37	5	7	38
115IC:IRWIN-----	90	N/A	4e	All areas are prime farmland	D	Clay Upland (pe25-34)	8	.37	.37	5	7	38
1191:BLAZEFORK--	90	2s-	2w	All areas are prime farmland	D	Clay Lowland (pe25-34)	8	.37	.37	5	7	38
1200:BUHLER-----	65	2w-	2w	Not prime farmland	D	Saline Subirrigated (pe21-28)	8	.43	.43	2	7	38
1200:BLAZEFORK--	30	2s-	2s	Not prime farmland	D	Clay Lowland (pe25-34)	8	.37	.37	5	7	38
1324:CARWAY-----	50	N/A	2w	Not prime farmland	D	Subirrigated (pe21-28)	3	.20	.20	5	3	86
1324:CARBIKA----	30	N/A	2w	Not prime farmland	D	Subirrigated (pe21-28)	6	.24	.24	5	5	56
1553:DARLOW-----	70	4s-	4s	Not prime farmland	C	Clay Pan (pe21- 28)	6	.43	.43	2	5	56
1553:ELMER-----	20	3s-	3s	Not prime farmland	C	Loamy Terrace (pe21-28)	3	.32	.32	2	3	86
1555:DILLHUT----	35	3e-	3e	Not prime farmland	B	Sands (pe21-28)	1	.15	.15	5	1	220
1555:PLEV-----	35	N/A	5w	Not prime farmland	B	Subirrigated (pe21-28)	2	.17	.17	5	2	134
159DP:DILLWYN---	60	N/A	4w	Not prime farmland	A	Subirrigated (pe21-28)	2	.17	.17	5	2	134
159DP:PLEVNA----	40	N/A	5w	Not prime farmland	D	Subirrigated (pe21-28)	2	.17	.17	5	2	134
159GC:GEARY-----	75	3e-	3e	All areas are prime farmland	B	Loamy Upland (pe25-34)	7	.32	.32	5	6	48

McPherson County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning--Continued

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
159GC:CLARK-----	25	N/A	4e	All areas are prime farmland	B	Limy Upland (pe25-34)	5	.28	.28	5	4L	86
159LA:LANCASTER--	90	N/A	3e	All areas are prime farmland	B	Loamy Upland (pe26-30)	7	.28	.28	3	6	48
159NF:NARON-----	100	2e-	3e	All areas are prime farmland	B	Sandy (pe21-28)	3	.20	.20	5	3	86
159PF:PRATT-----	100	3e-	3e	Not prime farmland	A	Sands (pe21-28)	2	.17	.17	5	2	134
159PR:PRATT-----	65	3e-	3e	Not prime farmland	A	Sands (pe21-28)	2	.17	.17	5	2	134
159PR:CARWILE---	35	N/A	2w	Not prime farmland	D	Sandy (pe21-28)	3	.24	.24	5	3	86
159SO:SMOLAN----	100	N/A	3e	Not prime farmland	C	Loamy Upland (pe25-34)	8	.37	.37	5	7	38
159WA:WALDECK---	100	N/A	3w	All areas are prime farmland	C	Subirrigated (pe21-28)	3	.20	.20	4	3	86
169CE:CLIME-----	100	N/A	4e	Not prime farmland	C	Limy Upland (pe25-34)	4	.37	.37	3	4	86
169ED:EDALGO----	100	N/A	4e	Not prime farmland	C	Clay Upland (pe26-30)	8	.37	.37	3	7	38
169RO:ROXBURY---	99	N/A	1	All areas are prime farmland	B	Loamy Terrace (pe26-30)	5	.32	.32	5	4L	86
1985:HAYES-----	60	3e-	3e	All areas are prime farmland	B	Sandy (pe21-28)	3	.20	.20	5	3	86
2391:KASKAN-----	75	N/A	5w	Not prime farmland	B	Loamy Lowland (pe21-28)	8	.37	.37	5	7	38
2588:LONGFORD---	90	N/A	4e	Not prime farmland	C	Loamy Upland (pe25-34)	8	.37	.37	5	7	38
3181:PRATT-----	45	3e-	3e	Not prime farmland	A	Sands (pe21-28)	1	.15	.15	5	1	220
3181:TURON-----	30	3e-	3e	Not prime farmland	A	Sands (pe21-28)	1	.15	.15	5	1	220
3641:TIVIN-----	45	N/A	6e	Not prime farmland	A	Choppy Sands (pe21-28)	1	.15	.15	5	1	220
3641:DILLHUT----	40	3e-	3e	Not prime farmland	B	Sands (pe21-28)	1	.15	.15	5	1	220
AED:ARENTS, EARTHEN DAM----	100	N/A	8	Not prime farmland		Unspecified		---	---	-	---	---
At:ATTICA-----	100	N/A	2e	Not prime farmland	B	Sandy (pe25-34)	2	.17	.17	5	2	134

McPherson County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning--Continued

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Br:BRIDGEPORT---	100	1-	1	All areas are prime farmland	B	Loamy Terrace (pe26-30)	5	.32	.32	5	4L	86
Ca:CARWILE-----	100	N/A	2w	Not prime farmland	D	Sandy (pe25-34)	3	.24	.24	5	3	86
Cb:CASS-----	100	2e-	2e	All areas are prime farmland	B	Sandy Lowland (pe26-30)	3	.20	.20	4	3	86
Ce:CLIME-----	100	N/A	3e	Not prime farmland	C	Limy Upland (pe26-30)	4	.28	.28	3	4	86
Cm:CLIME-----	100	N/A	4e	Not prime farmland	C	Limy Upland (pe26-30)	4	.28	.28	3	4	86
Cr:CRETE-----	100	2s-	2s	All areas are prime farmland	C	Clay Upland (pe25-34)	7	.37	.37	5	6	48
Cs:CRETE-----	100	2e-	2e	All areas are prime farmland	C	Clay Upland (pe25-34)	7	.37	.37	5	6	48
Ct:CRETE-----	100	2e-	2e	All areas are prime farmland	C	Clay Upland (pe25-34)	8	.37	.37	5	7	38
De:DETROIT-----	100	N/A	1	All areas are prime farmland	C	Loamy Terrace (pe26-30)	8	.37	.37	5	7	38
Dr:DRUMMOND----	100	N/A	6s	Not prime farmland	D	Saline Lowland (pe25-34)	5	.49	.49	2	4L	48
Ed:EDALGO-----	100	N/A	4e	Not prime farmland	C	Clay Upland (pe26-30)	7	.37	.37	3	6	48
Ee:EDALGO-----	100	N/A	4e	Not prime farmland	C	Clay Upland (pe26-30)	8	.37	.37	3	7	38
Fa:FARNUM-----	100	2e-	2e	All areas are prime farmland	B	Loamy Upland (pe25-34)	7	.28	.28	5	6	48
Ge:GEARY-----	100	2e-	2e	All areas are prime farmland	B	Loamy Upland (pe25-34)	7	.32	.32	5	6	48
Go:GOESSEL-----	100	N/A	2s	All areas are prime farmland	D	Clay Upland (pe25-34)	4	.28	.28	5	4	86
Ho:HORD-----	100	1-	1	All areas are prime farmland	B	Loamy Terrace (pe26-30)	7	.32	.32	5	6	48
INT:AQUOLLS----	100	N/A	5w	Not prime farmland	C	Unspecified		---	---	-	---	0
Ir:IRWIN-----	100	N/A	3e	All areas are prime farmland	D	Clay Upland (pe26-30)	8	.37	.37	4	7	38

McPherson County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning--Continued

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
LHH:LANCASTER----	60	N/A	6e	Not prime farmland	B	Loamy Upland (pe26-30)	7	.28	.43	3	6	48
LHH:HEDVILLE----	40	N/A	6e	Not prime farmland	D	Shallow Sandstone (pe26- 30)	6	.32	.43	1	5	56
La:LADYSMITH----	100	N/A	2s	All areas are prime farmland	D	Clay Upland (pe25-34)	8	.37	.37	5	7	38
Ld:LADYSMITH----	70	N/A	4s	Not prime farmland	D	Clay Upland (pe25-34)	8	.37	.37	5	7	38
Ld:DRUMMOND-----	30	N/A	6s	Not prime farmland	D	Saline Lowland (pe25-34)	7	.49	.49	2	6	48
Le:LANCASTER----	100	N/A	4e	All areas are prime farmland	B	Loamy Upland (pe26-30)	7	.28	.28	3	6	48
Lh:LANCASTER----	60	N/A	6e	Not prime farmland	B	Loamy Upland (pe26-30)	7	.28	.28	3	6	48
Lh:HEDVILLE-----	40	N/A	7s	Not prime farmland	D	Shallow Sandstone (pe26- 30)	6	.32	.32	2	5	56
Ln:LONGFORD-----	100	3e-	3e	All areas are prime farmland	C	Loamy Upland (pe26-30)	8	.32	.32	5	7	38
Lo:LONGFORD-----	100	N/A	4e	Not prime farmland	C	Loamy Upland (pe26-30)	8	.32	.32	5	7	38
Mc:MCCOOK-----	100	2e-	2e	All areas are prime farmland	B	Sandy Terrace (pe26-30)	3	.20	.20	5	3	86
Ns:NESS-----	100	N/A	6w	Not prime farmland	D	Clay Lowland (pe25-34)	4	.28	.28	5	4	86
Nw:NEW CAMBRIA--	100	N/A	2s	All areas are prime farmland	C	Clay Terrace (pe20-26)	4	.28	.28	5	4	86
Pa:PLEVNA-----	100	N/A	5w	Not prime farmland	D	Subirrigated (pe25-34)	3	.20	.20	5	3	86
Pr:PRATT-----	100	3e-	4e	Not prime farmland	A	Sands (pe25-34)	2	.17	.17	5	2	134
Ro:ROXBURY-----	100	1-	1	All areas are prime farmland	B	Loamy Terrace (pe26-30)	5	.43	.43	5	4L	86
Sm:SMOLAN-----	100	2e-	2e	All areas are prime farmland	C	Loamy Upland (pe25-34)	8	.37	.37	5	7	38
To:TOBIN-----	100	N/A	2w	All areas are prime farmland	B	Loamy Lowland (pe26-30)	7	.32	.32	5	6	48
W:WATER-----	100	N/A	N/A			Unspecified		---	---	-	---	---

McPherson County, Kansas: Published
Field Office Thunderbook: Soils Properties for Conservation Planning--Continued

Map symbol and soil name	Percent	Irr Cap Class	Nonirr Cap Class	Prime Farmland	Hydro- logic Group	Range site name	Windbreak suitability group	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
								K	Kf	T		
Wb:WELLS-----	90	N/A	2e	All areas are prime farmland	B	Loamy Upland (pe25-34)	7	.28	.28	5	6	48
Wc:WELLS-----	100	N/A	3e	All areas are prime farmland	B	Loamy Upland (pe26-30)	7	.28	.28	5	6	48

RANGELAND PRODUCTIVITY
McPherson County, Kansas

Use and Explanation of Rangeland, Grazed Forest Land, Native Pastureland Interpretations

Information in this subsection can be used to plan the use and management of soils for rangeland, grazed forest land, and native pasture. Different kinds of soils vary in their capacity to produce native grasses and other plants suitable for grazing. Information in this subsection provides groupings of similar soils and estimates of potential forage production, which can be used to determine livestock stocking rates.

Rangeland. Range is land on which the native vegetation (climax or natural potential plant community) is predominantly grasses, grasslike plants, forbs, and shrubs suitable for grazing and browsing. Range includes natural grasslands, savannas, many wetlands, some deserts, tundra, and certain shrub and forb communities. Rangeland receives no regular or frequent cultural treatment. The composition and production of the plant community are determined by soil, climate, topography, overstory canopy, and grazing management.

Grazed Forest Land. Includes land on which the understory includes, as an integral part of the forest plant community, plants that can be grazed without significantly impairing other forest values.

Native Pasture. Includes land on which the native vegetation (climax or natural potential plant community) is forest but which is used and managed primarily for production of native plants for forage. Native pasture includes cut-over forest land and forest land cleared and now managed for native or naturalized forage plants.

Rangeland

In areas that have similar climate and topography, differences in the kind and amount of vegetation produced on rangeland are closely related to the kind of soil. Effective management based on the relationship between the soils and vegetation and water.

The Rangeland, Grazed Forest land, Native Pastureland Interpretations shows, for each soil that supports rangeland vegetation, the ecological site and the potential annual production of vegetation in favorable, normal, unfavorable years. An explanation of the column headings in this table follows.

An ecological site is the product of all the environmental factors responsible for its development. It has characteristic soils that have developed over time throughout the soil development process; a characteristic hydrology, particularly infiltration and runoff, that has developed over time; and a characteristic plant community (kind and amount of vegetation). The hydrology of a site is influenced by development of the soil and plant community. The vegetation, soils, and hydrology are all interrelated. Each is influenced by the others and influences the development of the others. The plant community on an ecological site is typified by an association of species that differs from that of other ecological sites in the kind and/or proportion of species or in total production. Descriptions of ecological sites are provided in the Field Office Technical Guide, which is available in local offices of the Natural Resources Conservation Service.

Total dry-weight production is the amount of vegetation that can be expected to grow annually on well managed rangeland that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, average, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are about average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Yields are adjusted to a common percent of air-dry moisture content.

Range management requires a knowledge of the kinds of soil and of the potential natural plant community. It also requires an evaluation of the present range similarity index and rangeland trend. Range similarity index is determined by comparing the present plant community with the potential natural plant community on a particular rangeland ecological site. The more closely the existing community resembles the potential community, the higher the range similarity index. Rangeland trend is defined as the direction of change in an existing plant community relative to the potential natural plant community. Further information about the range similarity index and rangeland trend is available in chapter 4 of the National Range and Pasture Handbook, which is available in local offices of the Natural Resources Conservation Service. The objective in range management is to control grazing so that the plants growing on a site are about the same in kind and amount as the potential natural plant community for that site. Such management generally results in the optimum production of vegetation, control of undesirable brush species, conservation of water, and control of erosion. Sometimes, however, an area with a range similarity index somewhat below the potential meets grazing needs, provides wildlife habitat, and protects soil and water resources.

RANGELAND PRODUCTIVITY--Continued
McPherson County, Kansas

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
053ED: Edalgo-----	Clay Upland (pe26-30)	5,000	3,500	2,000
053MB: Mccook-----	Loamy Lowland (pe26-30)	3,800	3,300	2,800
053MC: Mccook-----	Loamy Terrace (pe26-30)	3,800	3,300	2,800
079GE: Geary-----	Loamy Upland (pe25-34)	6,000	4,000	3,000
079LB: Ladysmith-----	Clay Upland (pe25-34)	5,000	3,500	2,000
115IC: Irwin-----	Clay Upland (pe25-34)	5,000	3,500	2,500
159DP: Dillwyn-----	Subirrigated (pe21-28)	9,000	8,000	7,000
159GC: Plevna-----	Subirrigated (pe21-28)	9,000	8,000	7,000
159LA: Geary-----	Loamy Upland (pe25-34)	6,000	4,000	3,000
159NF: Clark-----	Limy Upland (pe25-34)	5,000	4,000	3,000
159PF: Lancaster-----	Loamy Upland (pe26-30)	5,250	4,000	2,750
159PR: Naron-----	Sandy (pe21-28)	4,500	3,000	2,000
159SO: Pratt-----	Sands (pe21-28)	4,500	3,500	2,500
159WA: Pratt-----	Sands (pe21-28)	4,500	3,500	2,500
169CE: Carwile-----	Sandy (pe21-28)	5,000	3,800	3,000
169ED: Smolan-----	Loamy Upland (pe25-34)	5,500	4,000	3,000
169RO: Waldeck-----	Subirrigated (pe21-28)	9,000	8,000	7,000
1985: Clime-----	Limy Upland (pe25-34)	5,000	3,500	2,500
2391: Edalgo-----	Clay Upland (pe26-30)	5,000	3,500	2,000
2588: Roxbury-----	Loamy Terrace (pe26-30)	5,000	4,000	3,000
3181: Blazefork-----	Clay Lowland (pe25-34)	6,500	5,000	4,000
3641: Buhler-----	Saline Subirrigated (pe21-28)	3,500	2,500	1,800
AED: Blazefork-----	Clay Lowland (pe25-34)	6,500	5,000	4,000
At: Carway-----	Subirrigated (pe21-28)	9,500	8,500	7,500
Br: Carbika-----	Subirrigated (pe21-28)	9,500	8,500	7,500
Ca: Darlow-----	Clay Pan (pe21-28)	3,500	2,500	1,800
Ch: Elmer-----	Loamy Terrace (pe21-28)	5,500	5,000	3,400
Cs: Dillhut-----	Sands (pe21-28)	4,500	3,500	2,500
Ct: Plev-----	Subirrigated (pe21-28)	9,500	8,500	7,500
De: Hayes-----	Sandy (pe21-28)	4,000	3,000	2,000
Ed: Kaskan-----	Loamy Lowland (pe21-28)	7,000	5,500	4,500
El: 2588: Longford, Moderately Eroded-----	Loamy Upland (pe25-34)	5,000	3,500	2,500
Em: 3181: Pratt-----	Sands (pe21-28)	4,500	3,500	2,500
En: Turon-----	Sands (pe21-28)	4,500	3,500	2,500
Er: 3641: Tivin-----	Choppy Sands (pe21-28)	3,000	2,150	1,550
Es: Dillhut-----	Sands (pe21-28)	4,500	3,500	2,500
Et: AED: Arents, Earthen Dam-----	---	---	---	---
Eu: At: Attica-----	Sandy (pe25-34)	4,500	3,000	2,000
Ev: Br: Bridgeport-----	Loamy Terrace (pe26-30)	5,000	4,000	3,000
Ex: Ca: Carwile-----	Sandy (pe25-34)	5,000	3,800	3,000
Ey: Ch: Cass-----	Sandy Lowland (pe26-30)	6,000	4,750	3,500
Fa: Ce: Clime-----	Limy Upland (pe26-30)	5,000	3,500	2,500
Fb: Cm: Clime-----	Limy Upland (pe26-30)	5,000	3,500	2,500
Fc: Cr: Crete-----	Clay Upland (pe25-34)	5,000	3,500	2,500
Fd: Cs: Crete-----	Clay Upland (pe25-34)	4,500	4,100	3,700
Fe: Ct: Crete-----	Clay Upland (pe25-34)	4,500	4,100	3,700
Fg: De: Detroit-----	Loamy Terrace (pe26-30)	6,000	4,500	3,000

RANGELAND PRODUCTIVITY--Continued
McPherson County, Kansas

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
Dr: Drummond-----	Saline Lowland (pe25-34)	7,000	5,800	5,000
Ed: Edalgo-----	Clay Upland (pe26-30)	5,000	3,500	2,500
Ee: Edalgo-----	Clay Upland (pe26-30)	5,000	3,500	2,000
Fa: Farnum-----	Loamy Upland (pe25-34)	5,500	4,000	2,500
Ge: Geary-----	Loamy Upland (pe25-34)	6,000	4,000	3,000
Go: Goessel-----	Clay Upland (pe25-34)	5,500	3,500	2,000
Ho: Hord-----	Loamy Terrace (pe26-30)	4,500	4,200	3,800
INT: Aquolls-----	---	---	---	---
Ir: Irwin, bedrock substratum-----	Clay Upland (pe26-30)	5,000	3,500	2,000
La: Ladysmith-----	Clay Upland (pe25-34)	5,000	3,500	2,000
Ld: Ladysmith-----	Clay Upland (pe25-34)	5,000	3,500	2,000
Drummond-----	Saline Lowland (pe25-34)	7,000	5,800	5,000
Le: Lancaster-----	Loamy Upland (pe26-30)	5,000	3,500	2,000
Lh: Lancaster-----	Loamy Upland (pe26-30)	5,000	3,500	2,000
Hedville-----	Shallow Sandstone (pe26-30)	4,000	3,000	2,000
LHH: Lancaster-----	Loamy Upland (pe26-30)	5,000	3,500	2,000
Hedville-----	Shallow Sandstone (pe26-30)	4,000	3,000	2,000
Ln: Longford-----	Loamy Upland (pe26-30)	5,500	4,000	3,000
Lo: Longford-----	Loamy Upland (pe26-30)	5,000	3,500	2,500
Mc: Mccook-----	Sandy Terrace (pe26-30)	3,800	3,300	2,800
Ns: Ness-----	Clay Lowland (pe25-34)	2,000	1,500	500
Nw: New Cambria-----	Clay Terrace (pe20-26)	5,000	4,000	2,500
Pa: Plevna-----	Subirrigated (pe25-34)	9,000	8,000	7,000
Pr: Pratt-----	Sands (pe25-34)	4,500	3,500	2,500
Ro: Roxbury-----	Loamy Terrace (pe26-30)	5,000	4,000	3,000
Sm: Smolan-----	Loamy Upland (pe25-34)	5,500	4,000	3,000
To: Tobin-----	Loamy Lowland (pe26-30)	6,000	5,000	4,000
W: Water-----	---	---	---	---
Wb: Wells-----	Loamy Upland (pe25-34)	5,250	4,000	2,750
Wc: Wells-----	Loamy Upland (pe26-30)	5,500	4,000	3,000

BUILDING SITE DEVELOPMENT
McPherson County, Kansas

Building Site Development

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. The following tables show the degree and kind of soil limitations that affect dwellings with and without basements, small commercial buildings, local roads and streets, shallow excavations, and lawns and landscaping.

The ratings in the tables are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Local roads and streets have an all-weather surface and carry automobile and light truck traffic all year. They have a subgrade of cut or fill soil material; a base of gravel, crushed rock, or soil material stabilized by lime or cement; and a surface of flexible material (asphalt), rigid material (concrete), or gravel with a binder. The ratings are based on the soil properties that affect the ease of excavation and grading and the traffic-supporting capacity. The properties that affect the ease of excavation and grading are depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, depth to a water table, ponding, flooding, the amount of large stones, and slope. The properties that affect the traffic-supporting capacity are soil strength (as inferred from the AASHTO group index number), subsidence, linear extensibility (shrink-swell potential), the potential for frost action, depth to a water table, and ponding.

Shallow excavations are trenches or holes dug to a maximum depth of 5 or 6 feet for graves, utility lines, open ditches, or other purposes. The ratings are based on the soil properties that influence the ease of digging and the resistance to sloughing. Depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, the amount of large stones, and dense layers influence the ease of digging, filling, and compacting. Depth to the seasonal high water table, flooding, and ponding may restrict the period when excavations can be made. Slope influences the ease of using machinery. Soil texture, depth to the water table, and linear extensibility (shrink-swell potential) influence the resistance to sloughing.

Lawns and landscaping require soils on which turf and ornamental trees and shrubs can be established and maintained. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer.

BUILDING SITE DEVELOPMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
053ED: Edalgo-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to soft bedrock	1.00 0.15	Very limited Shrink-swell Slope	1.00 0.12
053MB: Mccook-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
053MC: Mccook-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
079GE: Geary-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
079LB: Ladysmith-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
115IC: Irwin-----	90	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell Slope	1.00 0.12
159DP: Dillwyn-----	60	Somewhat limited Depth to saturated zone	0.44	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.44
Plevna-----	40	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
159GC: Geary-----	75	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
Clark-----	25	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
159LA: Lancaster-----	90	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Depth to soft bedrock	0.50 0.10	Somewhat limited Shrink-swell	0.50
159NF: Naron-----	100	Not limited		Not limited		Not limited	
159PF: Pratt-----	100	Not limited		Not limited		Not limited	
159PR: Pratt-----	65	Not limited		Not limited		Not limited	
Carwile-----	35	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00
159SO: Smolan-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell Slope	1.00 0.12
159WA: Waldeck-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.95	Very limited Flooding	1.00
169CE: Clime-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Shrink-swell	0.64 0.50	Somewhat limited Shrink-swell Slope	0.50 0.00
169ED: Edalgo-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to soft bedrock	1.00 0.42	Very limited Shrink-swell Slope	1.00 0.12
169RO: Roxbury-----	99	Very limited Flooding	1.00	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding	1.00

BUILDING SITE DEVELOPMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
1191: Blazefork-----	90	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 0.61 0.50	Very limited Flooding Shrink-swell	1.00 1.00
1200: Buhler-----	65	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.16	Very limited Flooding Shrink-swell	1.00 1.00
Blazefork-----	30	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 0.61 0.50	Very limited Flooding Shrink-swell	1.00 1.00
1324: Carway-----	50	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
Carbika-----	30	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
1553: Darlow-----	70	Not limited		Not limited		Not limited	
Elmer-----	20	Not limited		Not limited		Not limited	
1555: Dillhut-----	35	Not limited		Not limited		Not limited	
Plev-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
1985: Hayes-----	60	Not limited		Very limited Shrink-swell	1.00	Not limited	
2391: Kaskan-----	75	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.16	Very limited Flooding	1.00
2588: Longford, Moderately Eroded-----	90	Very limited Shrink-swell	 1.00	Very limited Shrink-swell	 1.00	Very limited Shrink-swell Slope	 1.00 0.00
3181: Pratt-----	45	Not limited		Not limited		Not limited	
Turon-----	30	Not limited		Not limited		Not limited	
3641: Tivin-----	45	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
Dillhut-----	40	Not limited		Not limited		Not limited	
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
At: Attica-----	100	Not limited		Not limited		Not limited	
Br: Bridgeport-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Ca: Carwile-----	100	Very limited Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00 1.00
Cb: Cass-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Ce: Clime-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Shrink-swell	0.71 0.50	Somewhat limited Shrink-swell	0.50

BUILDING SITE DEVELOPMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Cm: Cline-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Shrink-swell	0.71 0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
Cr: Crete-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
Cs: Crete-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
Ct: Crete-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
De: Detroit-----	100	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 1.00
Dr: Drummond-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to saturated zone	1.00 0.61	Very limited Shrink-swell	1.00
Ed: Edalgo-----	100	Very limited Shrink-swell Slope	1.00 0.04	Very limited Shrink-swell Depth to soft bedrock Slope	1.00 0.42 0.04	Very limited Shrink-swell Slope	1.00 1.00
Ee: Edalgo-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to soft bedrock	1.00 0.95	Very limited Shrink-swell Slope	1.00 0.48
Fa: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Ge: Geary-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Go: Goessel-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to saturated zone	1.00 1.00	Very limited Shrink-swell	1.00
Ho: Hord-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
INT: Aquolls-----	100	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
Ir: Irwin, bedrock substratum-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
La: Ladysmith-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
Ld: Ladysmith-----	70	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
Drummond-----	30	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to saturated zone	1.00 0.61	Very limited Shrink-swell	1.00
Le: Lancaster-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Depth to soft bedrock	0.50 0.29	Somewhat limited Shrink-swell Slope	0.50 0.00

BUILDING SITE DEVELOPMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Lh: Lancaster-----	60	Somewhat limited Shrink-swell Slope	0.50 0.04	Somewhat limited Shrink-swell Depth to soft bedrock Slope	0.50 0.29 0.04	Very limited Slope Shrink-swell	1.00 0.50
Hedville-----	40	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 1.00
LHH: Lancaster-----	60	Somewhat limited Shrink-swell Slope	0.50 0.00	Somewhat limited Shrink-swell Depth to soft bedrock Slope	0.50 0.46 0.00	Very limited Slope Shrink-swell	1.00 0.50
Hedville-----	40	Very limited Depth to hard bedrock Slope	1.00 0.63	Very limited Depth to hard bedrock Slope	1.00 0.63	Very limited Depth to hard bedrock Slope	1.00 1.00
Ln: Longford-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12
Lo: Longford-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell Slope	1.00 0.00
Mccook-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
Ns: Ness-----	100	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00
Nw: New Cambria-----	100	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Shrink-swell	1.00 1.00	Very limited Flooding Shrink-swell	1.00 1.00
Pa: Plevna-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Pr: Pratt-----	100	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Ro: Roxbury-----	100	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 0.50
Sm: Smolan-----	100	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00	Very limited Shrink-swell	1.00
To: Tobin-----	100	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 0.50
W: Water-----	100	Not rated		Not rated		Not rated	
Wb: Wells-----	90	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
Wc: Wells-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Slope	0.50 0.12

BUILDING SITE DEVELOPMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
053ED: Edalgo-----	100	Very limited Shrink-swell	1.00	Somewhat limited Too clayey Depth to soft bedrock Cutbanks cave	0.50 0.15 0.10	Somewhat limited Depth to bedrock	0.16
053MB: Mccook-----	100	Very limited Flooding	1.00	Somewhat limited Flooding Cutbanks cave	0.60 0.10	Somewhat limited Flooding	0.60
053MC: Mccook-----	100	Somewhat limited Flooding	0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
079GE: Geary-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
079LB: Ladysmith-----	100	Very limited Shrink-swell	1.00	Somewhat limited Too clayey Cutbanks cave	0.50 0.10	Not limited	
115IC: Irwin-----	90	Very limited Shrink-swell Frost action	1.00 0.50	Somewhat limited Too clayey Cutbanks cave	0.50 0.10	Not limited	
159DP: Dillwyn-----	60	Somewhat limited Depth to saturated zone	0.19	Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.21
Plevna-----	40	Very limited Flooding	1.00	Depth to saturated zone	1.00	Depth to saturated zone	0.19
		Depth to saturated zone	1.00	Very limited Depth to saturated zone Cutbanks cave	1.00	Very limited Flooding	1.00
				Flooding	0.80	Depth to saturated zone	1.00
159GC: Geary-----	75	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Clark-----	25	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
159LA: Lancaster-----	90	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave Depth to soft bedrock	0.10 0.10	Somewhat limited Depth to bedrock	0.10
159NF: Naron-----	100	Not limited		Somewhat limited Cutbanks cave	0.10	Not limited	
159PF: Pratt-----	100	Not limited		Very limited Cutbanks cave	1.00	Not limited	
159PR: Pratt-----	65	Not limited		Very limited Cutbanks cave	1.00	Not limited	
Carwile-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Shrink-swell	1.00	Too clayey Cutbanks cave	0.28 0.10		
159SO: Smolan-----	100	Very limited Shrink-swell	1.00	Somewhat limited Cutbanks cave Too clayey	0.10 0.03	Not limited	
159WA: Waldeck-----	100	Very limited Flooding	1.00	Very limited Cutbanks cave Depth to saturated zone Flooding	1.00 0.95 0.60	Somewhat limited Flooding	0.60
169CE: Cline-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Too clayey Cutbanks cave	0.64 0.28 0.10	Somewhat limited Depth to bedrock	0.65

BUILDING SITE DEVELOPMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
169ED: Edalgo-----	100	Very limited Shrink-swell	1.00	Somewhat limited Too clayey Depth to soft bedrock Cutbanks cave	0.50 0.42 0.10	Somewhat limited Depth to bedrock	0.42
169RO: Roxbury-----	99	Somewhat limited Flooding	0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
1191: Blazefork-----	90	Very limited Low strength	1.00	Somewhat limited Depth to saturated zone	0.61	Not limited	
		Shrink-swell	1.00	Too clayey	0.12		
		Flooding	0.40	Cutbanks cave	0.10		
1200: Buhler-----	65	Very limited Low strength	1.00	Somewhat limited Depth to saturated zone	0.16	Very limited Sodium content	1.00
		Shrink-swell	1.00	Cutbanks cave	0.10	Salinity	0.13
		Flooding	0.40	Too clayey	0.00		
Blazefork-----	30	Very limited Low strength	1.00	Somewhat limited Depth to saturated zone	0.61	Not limited	
		Shrink-swell	1.00	Too clayey	0.12		
		Flooding	0.40	Cutbanks cave	0.10		
1324: Carway-----	50	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Cutbanks cave	1.00 1.00 0.10	Very limited Ponding Depth to saturated zone	1.00 1.00
Carbika-----	30	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Cutbanks cave Too clayey	1.00 1.00 0.10 0.00	Very limited Ponding Depth to saturated zone	1.00 1.00
1553: Darlow-----	70	Very limited Low strength	1.00	Somewhat limited Cutbanks cave	0.10	Very limited Sodium content	1.00
Elmer-----	20	Not limited		Somewhat limited Cutbanks cave	0.10	Very limited Sodium content	1.00
1555: Dillhut-----	35	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.15
Plev-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone Droughty	1.00 0.92
1985: Hayes-----	60	Not limited		Somewhat limited Cutbanks cave Too clayey	0.10 0.02	Not limited	
2391: Kaskan-----	75	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding Depth to saturated zone	1.00 0.80 0.16	Very limited Flooding	1.00
2588: Longford, Moderately Eroded-----	90	Very limited Low strength Shrink-swell Frost action	 1.00 1.00 0.50	Somewhat limited Cutbanks cave	 0.10	Not limited	
3181: Pratt-----	45	Not limited		Very limited Cutbanks cave	1.00	Not limited	
Turon-----	30	Not limited		Very limited Cutbanks cave Too clayey	1.00 0.01	Not limited	
3641: Tivin-----	45	Somewhat limited Slope	0.16	Very limited Cutbanks cave Slope	1.00 0.16	Somewhat limited Droughty Slope	0.98 0.16
Dillhut-----	40	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.15

BUILDING SITE DEVELOPMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
At: Attica-----	100	Not limited		Very limited Cutbanks cave	1.00	Not limited	
Br: Bridgeport-----	100	Somewhat limited Flooding	0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
Ca: Carwile-----	100	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Too clayey Cutbanks cave	1.00 0.28 0.10	Very limited Depth to saturated zone	1.00
Cb: Cass-----	100	Somewhat limited Flooding	0.40	Very limited Cutbanks cave	1.00	Not limited	
Ce: Cline-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Too clayey Cutbanks cave	0.71 0.28 0.10	Very limited Too clayey Depth to bedrock Droughty	1.00 0.71 0.00
Cm: Cline-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Too clayey Cutbanks cave	0.71 0.28 0.10	Very limited Too clayey Depth to bedrock Droughty	1.00 0.71 0.00
Cr: Crete-----	100	Very limited Low strength Shrink-swell	1.00 1.00	Somewhat limited Too clayey Cutbanks cave	0.12 0.10	Not limited	
Cs: Crete-----	100	Very limited Shrink-swell	1.00	Somewhat limited Too clayey Cutbanks cave	0.12 0.10	Not limited	
Ct: Crete-----	100	Very limited Shrink-swell	1.00	Somewhat limited Too clayey Cutbanks cave	0.12 0.10	Not limited	
De: Detroit-----	100	Very limited Shrink-swell Flooding	1.00 0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
Dr: Drummond-----	100	Very limited Shrink-swell	1.00	Somewhat limited Depth to saturated zone Too clayey Cutbanks cave	0.61 0.28 0.10	Not limited	
Ed: Edalgo-----	100	Very limited Shrink-swell Slope	1.00 0.04	Somewhat limited Too clayey Depth to soft bedrock Cutbanks cave Slope	0.50 0.42 0.10 0.04	Somewhat limited Depth to bedrock Slope	0.42 0.04
Ee: Edalgo-----	100	Very limited Shrink-swell	1.00	Somewhat limited Depth to soft bedrock Too clayey Cutbanks cave	0.95 0.50 0.10	Somewhat limited Depth to bedrock Droughty	0.95 0.01
Fa: Farnum-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Ge: Geary-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Go: Goessel-----	100	Very limited Shrink-swell	1.00	Very limited Cutbanks cave Depth to saturated zone Too clayey	1.00 1.00 0.28	Very limited Too clayey	1.00

BUILDING SITE DEVELOPMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ho: Hord-----	100	Somewhat limited Flooding	0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
INT: Aguolls-----	100	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding Cutbanks cave	1.00 1.00 0.10	Very limited Depth to saturated zone Ponding	1.00 1.00
Ir: Irwin, bedrock substratum-----	100	Very limited Shrink-swell	 1.00	Somewhat limited Too clayey Cutbanks cave	 0.28 0.10	Not limited	
La: Ladysmith-----	100	Very limited Shrink-swell	1.00	Somewhat limited Too clayey Cutbanks cave	0.50 0.10	Not limited	
Ld: Ladysmith-----	70	Very limited Shrink-swell	1.00	Somewhat limited Too clayey Cutbanks cave	0.50 0.10	Not limited	
Drummond-----	30	Very limited Shrink-swell	1.00	Somewhat limited Depth to saturated zone Too clayey Cutbanks cave	0.61 0.28 0.10	Not limited	
Le: Lancaster-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to soft bedrock Cutbanks cave	0.29 0.10	Somewhat limited Depth to bedrock	0.29
Lh: Lancaster-----	60	Somewhat limited Shrink-swell Slope	0.50 0.04	Somewhat limited Depth to soft bedrock Cutbanks cave Slope	0.29 0.10 0.04	Somewhat limited Depth to bedrock Slope	0.29 0.04
Hedville-----	40	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Cutbanks cave Slope	1.00 0.10 0.04	Very limited Depth to bedrock Droughty Slope Content of large stones	1.00 0.62 0.04 0.03
LHH: Lancaster-----	60	Somewhat limited Shrink-swell Slope	0.50 0.00	Somewhat limited Depth to soft bedrock Cutbanks cave Slope	0.46 0.10 0.00	Somewhat limited Depth to bedrock Slope	0.46 0.00
Hedville-----	40	Very limited Depth to hard bedrock Slope	1.00 0.63	Very limited Depth to hard bedrock Slope Cutbanks cave	1.00 0.63 0.10	Very limited Depth to bedrock Droughty Slope Content of large stones	1.00 0.75 0.63 0.03
Ln: Longford-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
Lo: Longford-----	100	Very limited Shrink-swell	1.00	Somewhat limited Cutbanks cave	0.10	Not limited	
Mc: Mccook-----	100	Somewhat limited Flooding	0.40	Somewhat limited Cutbanks cave	0.10	Not limited	
Ns: Ness-----	100	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Too clayey	1.00 1.00 0.50	Very limited Depth to saturated zone Too clayey	1.00 1.00
Nw: New Cambria-----	100	Very limited Shrink-swell Flooding	1.00 0.40	Somewhat limited Too clayey Cutbanks cave	0.50 0.10	Very limited Too clayey	1.00

BUILDING SITE DEVELOPMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Pa: Plevna-----	100	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
Pr: Pratt-----	100	Somewhat limited Slope	0.04	Flooding	0.80		
				Very limited Cutbanks cave	1.00	Somewhat limited Slope	0.04
				Slope	0.04		
Ro: Roxbury-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
		Flooding	0.40				
Sm: Smolan-----	100	Very limited Shrink-swell	1.00	Somewhat limited Cutbanks cave	0.10	Not limited	
				Too clayey	0.03		
To: Tobin-----	100	Very limited Flooding	1.00	Somewhat limited Flooding	0.60	Somewhat limited Flooding	0.60
		Shrink-swell	0.50	Cutbanks cave	0.10		
W: Water-----	100	Not rated		Not rated		Not rated	
Wb: Wells-----	90	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
		Frost action	0.50				
Wc: Wells-----	100	Somewhat limited Shrink-swell	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	

CONSTRUCTION MATERIALS
McPherson County, Kansas

Construction Materials

The following tables give information about the soils as potential sources of gravel, sand, topsoil, reclamation material, and roadfill. Normal compaction, minor processing, and other standard construction practices are assumed.

The soils are rated good, fair, or poor as potential sources of topsoil, reclamation material, and roadfill. The features that limit the soils as sources of these materials are specified in the tables. The numerical ratings given after the specified features indicate the degree to which the features limit the soils as sources of topsoil, reclamation material, or roadfill. The lower the number, the greater the limitation.

The soils are rated as a probable or improbable source of sand and gravel. A rating of probable means that the source material is likely to be in or below the soil. The numerical ratings in these columns indicate the degree of probability. The number 0.00 indicates that the soil is an improbable source. A number between 0.00 and 1.00 indicates the degree to which the soil is a probable source of sand or gravel.

Sand and gravel are natural aggregates suitable for commercial use with a minimum of processing. They are used in many kinds of construction. Specifications for each use vary widely. In these tables, only the probability of finding material in suitable quantity is evaluated. The suitability of the material for specific purposes is not evaluated, nor are factors that affect excavation of the material. The properties used to evaluate the soil as a source of sand or gravel are gradation of grain sizes (as indicated by the Unified classification of the soil), the thickness of suitable material, and the content of rock fragments. If the lowest layer of the soil contains sand or gravel, the soil is rated as a probable source regardless of thickness. The assumption is that the sand or gravel layer below the depth of observation exceeds the minimum thickness.

Topsoil is used to cover an area so that vegetation can be established and maintained. The upper 40 inches of a soil is evaluated for use as topsoil. Also evaluated is the reclamation potential of the borrow area. The ratings are based on the soil properties that affect plant growth; the ease of excavating, loading, and spreading the material; and reclamation of the borrow area. Toxic substances, soil reaction, and the properties that are inferred from soil texture, such as available water capacity and fertility, affect plant growth. The ease of excavating, loading, and spreading is affected by rock fragments, slope, depth to a water table, soil texture, and thickness of suitable material. Reclamation of the borrow area is affected by slope, depth to a water table, rock fragments, depth to bedrock or a cemented pan, and toxic material.

The surface layer of most soils is generally preferred for topsoil because of its organic matter content. Organic matter greatly increases the absorption and retention of moisture and nutrients for plant growth.

Reclamation material is used in areas that have been drastically disturbed by surface mining or similar activities. When these areas are reclaimed, layers of soil material or unconsolidated geological material, or both, are replaced in a vertical sequence. The reconstructed soil favors plant growth. The ratings in the table do not apply to quarries and other mined areas that require an offsite source of reconstruction material. The ratings are based on the soil properties that affect erosion and stability of the surface and the productive potential of the reconstructed soil. These properties include the content of sodium, salts, and calcium carbonate; reaction; available water capacity; erodibility; texture; content of rock fragments; and content of organic matter and other features that affect fertility.

Roadfill is soil material that is excavated in one place and used in road embankments in another place. In this table, the soils are rated as a source of roadfill for low embankments, generally less than 6 feet high and less exacting in design than higher embankments.

The ratings are for the whole soil, from the surface to a depth of about 5 feet. It is assumed that soil layers will be mixed when the soil material is excavated and spread.

The ratings are based on the amount of suitable material and on soil properties that affect the ease of excavation and the performance of the material after it is in place. The thickness of the suitable material is a major consideration. The ease of excavation is affected by large stones, depth to a water table, and slope. How well the soil performs in place after it has been compacted and drained is determined by its strength (as inferred from the AASHTO classification of the soil) and linear extensibility (shrink-swell potential).

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
053ED: Edalgo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
053MB: Mccook-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
053MC: Mccook-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
079GE: Geary-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
079LB: Ladysmith-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
115IC: Irwin-----	90	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
159DP: Dillwyn-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.18 0.18
Plevna-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
159GC: Geary-----	75	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Clark-----	25	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
159LA: Lancaster-----	90	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
159NF: Naron-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.05 0.08
159PF: Pratt-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.49
159PR: Pratt-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.49
Carwile-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.07
159SO: Smolan-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
159WA: Waldeck-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Good Thickest layer	0.09
169CE: Cline-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
169ED: Edalgo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
169RO: Roxbury-----	99	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
1191: Blazefork-----	90	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
1200: Buhler-----	65	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.09
Blazefork-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
1324: Carway-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Carbika-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
1553: Darlow-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.12
Elmer-----	20	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.10
1555: Dillhut-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.10 0.13
Plev-----	35	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.41 0.43
1985: Hayes-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
2391: Kaskan-----	75	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.36 0.40
2588: Longford, Moderately Eroded-----	90	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
3181: Pratt-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.84 0.86
Turon-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.00 0.30
3641: Tivin-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Good	
Dillhut-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.10 0.13
AED: Arents, Earthen Dam-	100	Not rated		Not rated	
At: Attica-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.72
Br: Bridgeport-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ca: Carwile-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Cb: Cass-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.08 0.70
Ce: Clime-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Cm: Clime-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Cr: Crete-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Cs: Crete-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ct: Crete-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
De: Detroit-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Dr: Drummond-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Ed: Edalgo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ee: Edalgo-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Fa: Farnum-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.03 0.06
Ge: Geary-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Go: Goessel-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ho: Hord-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
INT: Aquolls-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ir: Irwin, bedrock substratum-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
La: Ladysmith-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Ld: Ladysmith-----	70	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Drummond-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Le: Lancaster-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Lh: Lancaster-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Hedville-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
LHH: Lancaster-----	60	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Hedville-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
Ln: Longford-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Lo: Longford-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Mc: Mccook-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.05 0.09
Ns: Ness-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Nw: New Cambria-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Pa: Plevna-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.09 0.20
Pr: Pratt-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.19 0.47
Ro: Roxbury-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Sm: Smolan-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
To: Tobin-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
W: Water-----	100	Not rated		Not rated	
Wb: Wells-----	90	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.06
Wc: Wells-----	100	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.03 0.08

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
053ED: Edalgo-----	100	Poor Too clayey Depth to bedrock Too acid Droughty No water erosion limitation	0.00 0.84 0.84 0.89 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Depth to bedrock	0.00 0.84
053MB: Mccook-----	100	Poor Low content of organic matter Water erosion	0.00 0.90	Good		Good	
053MC: Mccook-----	100	Poor Low content of organic matter Water erosion	0.00 0.90	Good		Good	
079GE: Geary-----	100	Poor Low content of organic matter Water erosion Too acid Too clayey	0.00 0.90 0.95 0.98	Fair Shrink-swell	0.87	Fair Too Clayey	0.49
079LB: Ladysmith-----	100	Poor Too clayey No water erosion limitation	0.00 0.99	Fair Shrink-swell	0.31	Poor Too Clayey	0.00
115IC: Irwin-----	90	Poor Too clayey No water erosion limitation	0.00 0.99	Fair Shrink-swell	0.16	Poor Too Clayey	0.00
159DP: Dillwyn-----	60	Poor Wind erosion Low content of organic matter Too sandy Droughty	0.00 0.00 0.36 0.80	Fair Depth to saturated zone	0.53	Fair Too sandy Depth to saturated zone	0.36 0.53
Plevna-----	40	Poor Wind erosion Low content of organic matter	0.00 0.00	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone	0.00
159GC: Geary-----	75	Poor Low content of organic matter Water erosion Too acid Too clayey	0.00 0.90 0.95 0.98	Fair Shrink-swell	0.87	Fair Too Clayey	0.49
Clark-----	25	Poor Low content of organic matter Carbonate content	0.00 0.68	Fair Shrink-swell	0.87	Fair Carbonate content	0.68
159LA: Lancaster-----	90	Fair Depth to bedrock Too acid	0.90 0.95	Poor Depth to bedrock Shrink-swell	0.00 0.99	Fair Depth to bedrock	0.90
159NF: Naron-----	100	Poor Low content of organic matter	0.00	Good		Good	

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
159PF: Pratt-----	100	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.00 0.00	Good		Poor Too sandy	0.00
159PR: Pratt-----	65	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.00 0.00	Good		Poor Too sandy	0.00
Carwile-----	35	Poor Low content of organic matter Too clayey Too acid No water erosion limitation	0.00 0.00 0.00 0.97 0.99	Poor Depth to saturated zone Shrink-swell	0.00 0.23	Poor Depth to saturated zone Too Clayey	0.00 0.00
159SO: Smolan-----	100	Poor Low content of organic matter Too clayey No water erosion limitation	0.00 0.00 0.99	Fair Shrink-swell	0.12	Poor Too Clayey	0.00
159WA: Waldeck-----	100	Good		Good		Good	
169CE: Clime-----	100	Poor Too clayey Depth to bedrock Droughty No water erosion limitation	0.00 0.35 0.55 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.87	Poor Too Clayey Depth to bedrock	0.00 0.35
169ED: Edalgo-----	100	Poor Too clayey Depth to bedrock Droughty Too acid No water erosion limitation	0.00 0.58 0.72 0.84 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.23	Poor Too Clayey Depth to bedrock	0.00 0.58
169RO: Roxbury-----	99	Fair Low content of organic matter Water erosion	0.50 0.90	Fair Shrink-swell	0.98	Good	
1191: Blazefork-----	90	Poor Too clayey Low content of organic matter Too acid Water erosion	0.00 0.06 0.32 0.90	Poor Low strength Shrink-swell	0.00 0.50	Poor Too Clayey	0.00
1200: Buhler-----	65	Poor Sodium content Too alkaline Low content of organic matter Too clayey Too acid Water erosion	0.00 0.00 0.05 0.23 0.88 0.90	Fair Shrink-swell	0.30	Poor Sodium content Too Clayey Salinity	0.00 0.16 0.50

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Blazefork-----	30	Poor Too clayey Low content of organic matter Too acid Water erosion	0.00 0.06 0.32 0.90	Poor Low strength Shrink-swell	0.00 0.50	Poor Too Clayey	0.00
1324: Carway-----	50	Fair Low content of organic matter Too acid No water erosion limitation	0.12 0.95 0.99	Poor Depth to saturated zone Low strength Shrink-swell	0.00 0.00 0.89	Poor Depth to saturated zone	0.00
Carbika-----	30	Fair Too clayey Low content of organic matter Too acid No water erosion limitation	0.74 0.88 0.95 0.99	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone Too Clayey	0.00 0.53
1553: Darlow-----	70	Poor Sodium content Too alkaline Low content of organic matter Too acid Salinity Water erosion	0.00 0.00 0.08 0.16 0.88 0.90	Good		Poor Sodium content Salinity	0.00 0.00
Elmer-----	20	Poor Too alkaline Too acid Low content of organic matter Sodium content No water erosion limitation	0.00 0.16 0.46 0.78 0.99	Fair Shrink-swell	0.99	Poor Sodium content	0.00
1555: Dillhut-----	35	Poor Wind erosion Low content of organic matter Too acid	0.00 0.00 0.99	Good		Good	
Plev-----	35	Poor Too sandy Wind erosion Low content of organic matter Too acid Droughty	0.00 0.00 0.00 0.95 0.99	Poor Depth to saturated zone	0.00	Poor Too sandy Depth to saturated zone	0.00 0.00
1985: Hayes-----	60	Fair Low content of organic matter Too acid	0.12 0.97	Poor Low strength	0.00	Good	
2391: Kaskan-----	75	Fair Too sandy Low content of organic matter No water erosion limitation	0.01 0.12 0.99	Good		Fair Too sandy	0.01

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
2588: Longford, Moderately Eroded-----	90	Poor Too clayey Low content of organic matter Water erosion	0.00 0.88 0.90	Poor Low strength Shrink-swell	0.00 0.36	Poor Too Clayey	0.00
3181: Pratt-----	45	Poor Wind erosion Too sandy Low content of organic matter Too acid	0.00 0.00 0.00 0.74	Good		Poor Too sandy	0.00
Turon-----	30	Poor Too sandy Wind erosion Too acid Low content of organic matter	0.00 0.00 0.39 0.88	Good		Poor Too sandy Too acid	0.00 0.92
3641: Tivin-----	45	Poor Too sandy Wind erosion Low content of organic matter Droughty Too acid	0.00 0.00 0.00 0.36 0.99	Good		Poor Too sandy Slope	0.00 0.84
Dillhut-----	40	Poor Wind erosion Low content of organic matter Too acid	0.00 0.00 0.99	Good		Good	
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
At: Attica-----	100	Poor Wind erosion Low content of organic matter Too acid	0.00 0.00 0.95	Good		Good	
Br: Bridgeport-----	100	Poor Low content of organic matter Water erosion	0.00 0.90	Good		Good	
Ca: Carwile-----	100	Poor Low content of organic matter Too clayey Too acid No water erosion limitation	0.00 0.00 0.97 0.99	Poor Depth to saturated zone Shrink-swell	0.00 0.36	Poor Depth to saturated zone Too Clayey	0.00 0.00
Cb: Cass-----	100	Poor Low content of organic matter	0.00	Good		Good	
Ce: Clime-----	100	Poor Too clayey Droughty Depth to bedrock	0.00 0.20 0.29	Poor Depth to bedrock Shrink-swell	0.00 0.87	Poor Too Clayey Depth to bedrock	0.00 0.29
Cm: Clime-----	100	Poor Too clayey Droughty Depth to bedrock	0.00 0.20 0.29	Poor Depth to bedrock Shrink-swell	0.00 0.87	Poor Too Clayey Depth to bedrock	0.00 0.29

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Cr: Crete-----	100	Poor Too clayey Too acid Low content of organic matter Water erosion	0.00 0.84 0.88 0.90	Poor Low strength Shrink-swell	0.00 0.12	Poor Too Clayey	0.00
Cs: Crete-----	100	Poor Low content of organic matter Too clayey Too acid No water erosion limitation	0.00 0.00 0.84 0.99	Fair Shrink-swell	0.12	Poor Too Clayey	0.00
Ct: Crete-----	100	Poor Low content of organic matter Too clayey Too acid No water erosion limitation	0.00 0.00 0.84 0.99	Fair Shrink-swell	0.12	Poor Too Clayey	0.00
De: Detroit-----	100	Poor Too clayey Low content of organic matter No water erosion limitation	0.00 0.88 0.99	Fair Shrink-swell	0.57	Poor Too Clayey	0.00
Dr: Drummond-----	100	Poor Low content of organic matter Too clayey Water erosion	0.00 0.00 0.37	Fair Shrink-swell	0.38	Poor Too Clayey Salinity	0.00 0.88
Ed: Edalgo-----	100	Poor Too clayey Depth to bedrock Too acid Droughty No water erosion limitation	0.00 0.58 0.84 0.85 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.28	Poor Too Clayey Depth to bedrock Slope	0.00 0.58 0.96
Ee: Edalgo-----	100	Poor Too clayey Depth to bedrock Droughty Too acid No water erosion limitation	0.00 0.05 0.17 0.84 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.12	Poor Too Clayey Depth to bedrock	0.00 0.05
Fa: Farnum-----	100	Poor Low content of organic matter	0.00	Fair Shrink-swell	0.97	Good	
Ge: Geary-----	100	Poor Low content of organic matter Water erosion Too acid Too clayey	0.00 0.90 0.95 0.98	Fair Shrink-swell	0.87	Fair Too Clayey	0.49
Go: Goessel-----	100	Poor Too clayey Low content of organic matter	0.00 0.00	Fair Shrink-swell Depth to saturated zone	0.12 0.89	Poor Too Clayey Depth to saturated zone	0.00 0.89

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ho: Hord-----	100	Poor Low content of organic matter	0.00	Good		Good	
INT: Aguolls-----	100	Poor Low content of organic matter	0.00	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone	0.00
Ir: Irwin, bedrock substratum-----	100	Poor Too clayey Low content of organic matter No water erosion limitation	0.00 0.00 0.99	Fair Shrink-swell Depth to bedrock	0.14 0.58	Poor Too Clayey	0.00
La: Ladysmith-----	100	Poor Too clayey No water erosion limitation	0.00 0.99	Fair Shrink-swell	0.26	Poor Too Clayey	0.00
Ld: Ladysmith-----	70	Poor Too clayey No water erosion limitation	0.00 0.99	Fair Shrink-swell	0.26	Poor Too Clayey	0.00
Drummond-----	30	Poor Low content of organic matter Too clayey Water erosion	0.00 0.00 0.37	Fair Shrink-swell	0.12	Poor Too Clayey Salinity	0.00 0.88
Le: Lancaster-----	100	Poor Low content of organic matter Depth to bedrock Too acid Droughty	0.00 0.71 0.95 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.87	Fair Depth to bedrock	0.71
Lh: Lancaster-----	60	Poor Low content of organic matter Depth to bedrock Too acid Droughty	0.00 0.71 0.95 0.99	Poor Depth to bedrock Shrink-swell	0.00 0.87	Fair Depth to bedrock Slope	0.71 0.96
Hedville-----	40	Poor Depth to bedrock Droughty	0.00 0.00	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope	0.00 0.95 0.96
LHH: Lancaster-----	60	Poor Low content of organic matter Depth to bedrock Droughty Too acid	0.00 0.54 0.93 0.95	Poor Depth to bedrock Shrink-swell	0.00 0.98	Fair Depth to bedrock	0.54
Hedville-----	40	Poor Low content of organic matter Depth to bedrock Droughty	0.00 0.00 0.00	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope	0.00 0.00 0.37
Ln: Longford-----	100	Fair Too clayey	0.98	Fair Shrink-swell	0.72	Fair Too Clayey	0.97

CONSTRUCTION MATERIALS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of reclamation material		Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Lo: Longford-----	100	Poor Too clayey Low content of organic matter Water erosion	0.00 0.88 0.90	Fair Shrink-swell	0.36	Poor Too Clayey	0.00
Mc: Mccook-----	100	Poor Low content of organic matter Water erosion	0.00 0.90	Good		Good	
Ns: Ness-----	100	Poor Too clayey	0.00	Poor Depth to saturated zone Shrink-swell	0.00 0.12	Poor Too Clayey Depth to saturated zone	0.00 0.00
Nw: New Cambria-----	100	Poor Low content of organic matter Too clayey	0.00 0.00	Fair Shrink-swell	0.12	Poor Too Clayey	0.00
Pa: Plevna-----	100	Poor Low content of organic matter	0.00	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone	0.00
Pr: Pratt-----	100	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.00 0.00	Good		Poor Too sandy Slope	0.00 0.96
Ro: Roxbury-----	100	Fair Water erosion Too clayey	0.90 0.98	Fair Shrink-swell	0.87	Fair Too Clayey	0.98
Sm: Smolan-----	100	Poor Low content of organic matter Too clayey No water erosion limitation	0.00 0.00 0.99	Fair Shrink-swell	0.20	Poor Too Clayey	0.00
To: Tobin-----	100	Fair Low content of organic matter Water erosion	0.50 0.90	Fair Shrink-swell	0.96	Good	
W: Water-----	100	Not rated		Not rated		Not rated	
Wb: Wells-----	90	Fair Too acid Too clayey	0.95 0.98	Good		Fair Too Clayey	0.93
Wc: Wells-----	100	Poor Low content of organic matter Too acid Too clayey	0.00 0.95 0.98	Fair Shrink-swell	0.96	Fair Too Clayey	0.49

RECREATIONAL INTERPRETATIONS
McPherson County, Kansas

Recreation

The soils of the survey area are rated in the following tables according to limitations that affect their suitability for recreation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the recreational uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The ratings in the tables are based on restrictive soil features, such as wetness, slope, and texture of the surface layer. Susceptibility to flooding is considered. Not considered in the ratings, but important in evaluating a site, are the location and accessibility of the area, the size and shape of the area and its scenic quality, vegetation, access to water, potential water impoundment sites, and access to public sewer lines. The capacity of the soil to absorb septic tank effluent and the ability of the soil to support vegetation also are important. Soils that are subject to flooding are limited for recreational uses by the duration and intensity of flooding and the season when flooding occurs. In planning recreational facilities, onsite assessment of the height, duration, intensity, and frequency of flooding is essential.

The information in this table can be supplemented by other information in this survey, for example, interpretations for building site development, construction materials, sanitary facilities, and water management.

Camp areas require site preparation, such as shaping and leveling the tent and parking areas, stabilizing roads and intensively used areas, and installing sanitary facilities and utility lines. Camp areas are subject to heavy foot traffic and some vehicular traffic. The ratings are based on the soil properties that affect the ease of developing camp areas and the performance of the areas after development. Slope, stoniness, and depth to bedrock or a cemented pan are the main concerns affecting the development of camp areas.

The soil properties that affect the performance of the areas after development are those that influence trafficability and promote the growth of vegetation, especially in heavily used areas. For good trafficability, the surface of camp areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Picnic areas are subject to heavy foot traffic. Most vehicular traffic is confined to access roads and parking areas. The ratings are based on the soil properties that affect the ease of developing picnic areas and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of picnic areas. For good trafficability, the surface of picnic areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Playgrounds require soils that are nearly level, are free of stones, and can withstand intensive foot traffic. The ratings are based on the soil properties that affect the ease of developing playgrounds and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of playgrounds. For good trafficability, the surface of the playgrounds should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Paths and trails for hiking and horseback riding should require little or no slope modification through cutting and filling. The ratings are based on the soil properties that affect trafficability and erodibility. These properties are stoniness, depth to a water table, ponding, flooding, slope, and texture of the surface layer.

Golf fairways are subject to heavy foot traffic and some light vehicular traffic. Cutting or filling may be required. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer. The suitability of the soil for traps, tees, roughs, and greens is not considered in the ratings.

RECREATIONAL INTERPRETATIONS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
053ED: Edalgo-----	100	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45	Somewhat limited Slope Restricted permeability Depth to bedrock	0.87 0.45 0.16
053MB: Mccook-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
053MC: Mccook-----	100	Very limited Flooding	1.00	Not limited		Not limited	
079GE: Geary-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
079LB: Ladysmith-----	100	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability Slope	0.45 0.00
115IC: Irwin-----	90	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45	Somewhat limited Slope Restricted permeability	0.87 0.45
159DP: Dillwyn-----	60	Somewhat limited Depth to saturated zone Too sandy	0.44 0.37	Somewhat limited Too sandy Depth to saturated zone	0.37 0.19	Somewhat limited Depth to saturated zone Too sandy	0.44 0.37
Plevna-----	40	Very limited Flooding Depth to saturated zone Too sandy	1.00 1.00 0.37	Very limited Depth to saturated zone Flooding Too sandy	1.00 0.40 0.37	Very limited Flooding Depth to saturated zone Too sandy	1.00 1.00 0.37
159GC: Geary-----	75	Not limited		Not limited		Somewhat limited Slope	0.87
Clark-----	25	Not limited		Not limited		Somewhat limited Slope	0.87
159LA: Lancaster-----	90	Not limited		Not limited		Somewhat limited Slope	0.00
159NF: Naron-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
159PF: Pratt-----	100	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy Slope	0.37 0.13
159PR: Pratt-----	65	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy	0.37	Somewhat limited Too sandy Slope	0.37 0.13
Carwile-----	35	Very limited Depth to saturated zone Restricted permeability	1.00 0.94	Very limited Depth to saturated zone Restricted permeability	1.00 0.94	Very limited Depth to saturated zone Restricted permeability	1.00 0.94
159SO: Smolan-----	100	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39	Somewhat limited Slope Restricted permeability	0.87 0.39
159WA: Waldeck-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
169CE: Cline-----	100	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39	Somewhat limited Depth to bedrock Slope Restricted permeability	0.65 0.50 0.39
169ED: Edalgo-----	100	Somewhat limited		Somewhat limited		Somewhat limited	

RECREATIONAL INTERPRETATIONS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
169RO: Roxbury-----	99	Restricted permeability	0.45	Restricted permeability	0.45	Slope	0.87
						Restricted permeability	0.45
						Depth to bedrock	0.42
1191: Blazefork-----	90	Very limited Flooding	1.00	Not limited		Not limited	
1200: Buhler-----	65	Very limited Flooding	1.00	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39
		Restricted permeability	0.39				
		Very limited Sodium content Flooding	1.00	Very limited Sodium content Restricted permeability Salinity	1.00 0.45	Very limited Sodium content Restricted permeability Salinity	1.00 0.45
Blazefork-----	30	Restricted permeability Salinity	0.45 0.13				
		Very limited Flooding	1.00	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39
		Restricted permeability	0.39				
1324: Carway-----	50	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding	1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
Carbika-----	30	Restricted permeability	1.00	Depth to saturated zone Restricted permeability	1.00	Restricted permeability	1.00
		Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding	1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
		Restricted permeability	1.00	Depth to saturated zone Restricted permeability	1.00	Restricted permeability	1.00
1553: Darlow-----	70	Very limited Sodium content Restricted permeability	1.00 0.45	Very limited Sodium content Restricted permeability	1.00 0.45	Very limited Sodium content Restricted permeability	1.00 0.45
Elmer-----	20	Very limited Sodium content Restricted permeability	1.00 0.39	Very limited Sodium content Restricted permeability	1.00 0.39	Very limited Sodium content Restricted permeability	1.00 0.39
1555: Dillhut-----	35	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy	1.00
Plev-----	35					Slope	0.00
		Very limited Depth to saturated zone Too sandy	1.00 0.94	Very limited Depth to saturated zone Too sandy	1.00 0.94	Very limited Depth to saturated zone Too sandy	1.00 0.94
1985: Hayes-----	60	Not limited		Not limited		Somewhat limited Slope	0.13
2391: Kaskan-----	75	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
2588: Longford, Moderately Eroded-----	90	Somewhat limited		Somewhat limited		Somewhat limited	
		Restricted permeability	0.05	Restricted permeability	0.05	Slope	0.50
3181: Pratt-----	45					Restricted permeability	0.05
		Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Too sandy	1.00
Turon-----	30					Slope	0.13
		Somewhat limited Too sandy	0.98	Somewhat limited Too sandy	0.98	Somewhat limited Too sandy	0.98
3641: Tivin-----	45	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Slope	0.13
						Very limited Too sandy	1.00

RECREATIONAL INTERPRETATIONS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Dillhut-----	40	Slope Very limited Too sandy	0.16 1.00	Slope Very limited Too sandy	0.16 1.00	Slope Very limited Too sandy Slope	1.00 1.00 0.13
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
At: Attica-----	100	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy	0.96	Somewhat limited Too sandy Slope	0.96 0.13
Br: Bridgeport-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Ca: Carwile-----	100	Very limited Depth to saturated zone Restricted permeability	1.00 0.94	Very limited Depth to saturated zone Restricted permeability	1.00 0.94	Very limited Depth to saturated zone Restricted permeability	1.00 0.94
Cb: Cass-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Ce: Clime-----	100	Somewhat limited Too clayey Restricted permeability	0.50 0.39	Somewhat limited Too clayey Restricted permeability	0.50 0.39	Somewhat limited Too clayey Restricted permeability Slope	0.50 0.39 0.00
Cm: Clime-----	100	Somewhat limited Too clayey Restricted permeability	0.50 0.39	Somewhat limited Too clayey Restricted permeability	0.50 0.39	Somewhat limited Slope Depth to bedrock Too clayey Restricted permeability	0.87 0.71 0.50 0.39
Cr: Crete-----	100	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39
Cs: Crete-----	100	Somewhat limited Restricted permeability	0.05	Somewhat limited Restricted permeability	0.05	Somewhat limited Restricted permeability Slope	0.05 0.00
Ct: Crete-----	100	Somewhat limited Restricted permeability	0.05	Somewhat limited Restricted permeability	0.05	Somewhat limited Restricted permeability Slope	0.05 0.00
De: Detroit-----	100	Very limited Flooding Restricted permeability	1.00 0.39	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39
Dr: Drummond-----	100	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45
Ed: Edalgo-----	100	Somewhat limited Restricted permeability Slope	0.45 0.04	Somewhat limited Restricted permeability Slope	0.45 0.04	Very limited Slope Restricted permeability Depth to bedrock	1.00 0.45 0.42
Ee: Edalgo-----	100	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45	Very limited Slope Depth to bedrock Restricted permeability	1.00 0.95 0.45
Fa: Farnum-----	100	Not limited		Not limited		Somewhat limited Slope	0.00
Ge: Geary-----	100	Not limited		Not limited		Somewhat limited Slope	0.00

RECREATIONAL INTERPRETATIONS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Go: Goessel-----	100	Somewhat limited Too clayey Restricted permeability	0.50 0.45	Somewhat limited Too clayey Restricted permeability	0.50 0.45	Somewhat limited Too clayey Restricted permeability	0.50 0.45
Ho: Hord-----	100	Very limited Flooding	1.00	Not limited		Not limited	
INT: Aquolls-----	100	Very limited Depth to saturated zone Restricted permeability Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Restricted permeability Ponding	1.00 1.00 1.00	Very limited Restricted permeability Depth to saturated zone Ponding	1.00 1.00 1.00
Ir: Irwin, bedrock substratum-----	100	Somewhat limited Restricted permeability	 0.45	Somewhat limited Restricted permeability	 0.45	Somewhat limited Restricted permeability Slope	 0.45 0.00
La: Ladysmith-----	100	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45
Ld: Ladysmith-----	70	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45
Drummond-----	30	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45	Somewhat limited Restricted permeability	0.45
Le: Lancaster-----	100	Not limited		Not limited		Somewhat limited Slope Depth to bedrock	0.50 0.29
Lh: Lancaster-----	60	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope Depth to bedrock	1.00 0.29
Hedville-----	40	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Slope Content of large stones Gravel content	1.00 1.00 0.03 0.02
LHH: Lancaster-----	60	Somewhat limited Slope	0.00	Somewhat limited Slope	0.00	Very limited Slope Depth to bedrock	1.00 0.46
Hedville-----	40	Very limited Depth to bedrock Slope	1.00 0.63	Very limited Depth to bedrock Slope	1.00 0.63	Very limited Depth to bedrock Slope Content of large stones Gravel content	1.00 1.00 0.03 0.02
Ln: Longford-----	100	Not limited		Not limited		Somewhat limited Slope	0.87
Lo: Longford-----	100	Somewhat limited Restricted permeability	0.05	Somewhat limited Restricted permeability	0.05	Somewhat limited Slope Restricted permeability	0.50 0.05
Mc: Mccook-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Ns: Ness-----	100	Very limited Depth to saturated zone Too clayey Restricted permeability	1.00 0.50 0.45	Very limited Depth to saturated zone Too clayey Restricted permeability	1.00 0.50 0.45	Very limited Depth to saturated zone Too clayey Restricted permeability	1.00 0.50 0.45
Nw: New Cambria-----	100	Very limited Flooding Too clayey Restricted permeability	1.00 0.50 0.39	Somewhat limited Too clayey Restricted permeability	0.50 0.39	Somewhat limited Too clayey Restricted permeability	0.50 0.39

RECREATIONAL INTERPRETATIONS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Pa: Plevna-----	100	Very limited Flooding	1.00	Very limited Depth to saturated zone Flooding	1.00	Very limited Flooding	1.00
		Depth to saturated zone	1.00		0.40	Depth to saturated zone	1.00
Pr: Pratt-----	100	Somewhat limited Too sandy Slope	0.37 0.04	Somewhat limited Too sandy Slope	0.37 0.04	Very limited Slope Too sandy	1.00 0.37
Ro: Roxbury-----	100	Very limited Flooding	1.00	Not limited		Not limited	
Sm: Smolan-----	100	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability	0.39	Somewhat limited Restricted permeability Slope	0.39 0.00
To: Tobin-----	100	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60
W: Water-----	100	Not rated		Not rated		Not rated	
Wb: Wells-----	90	Not limited		Not limited		Somewhat limited Slope	0.00
Wc: Wells-----	100	Not limited		Not limited		Somewhat limited Slope	0.87

RECREATIONAL INTERPRETATIONS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
053ED: Edalgo-----	100	Not limited		Somewhat limited Depth to bedrock	0.16
053MB: Mccook-----	100	Not limited		Somewhat limited Flooding	0.60
053MC: Mccook-----	100	Not limited		Not limited	
079GE: Geary-----	100	Not limited		Not limited	
079LB: Ladysmith-----	100	Not limited		Not limited	
115IC: Irwin-----	90	Not limited		Not limited	
159DP: Dillwyn-----	60	Somewhat limited Too sandy	0.37	Somewhat limited Droughty Depth to saturated zone	0.21 0.19
Plevna-----	40	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 1.00
		Too sandy	0.37		
159GC: Geary-----	75	Not limited		Not limited	
Clark-----	25	Not limited		Not limited	
159LA: Lancaster-----	90	Not limited		Somewhat limited Depth to bedrock	0.10
159NF: Naron-----	100	Not limited		Not limited	
159PF: Pratt-----	100	Somewhat limited Too sandy	0.37	Not limited	
159PR: Pratt-----	65	Somewhat limited Too sandy	0.37	Not limited	
Carwile-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
159SO: Smolan-----	100	Not limited		Not limited	
159WA: Waldeck-----	100	Not limited		Somewhat limited Flooding	0.60
169CE: Clime-----	100	Not limited		Somewhat limited Depth to bedrock	0.65
169ED: Edalgo-----	100	Not limited		Somewhat limited Depth to bedrock	0.42
169RO: Roxbury-----	99	Not limited		Not limited	
1191: Blazefork-----	90	Not limited		Not limited	
1200: Buhler-----	65	Not limited		Very limited Sodium content Salinity	1.00 0.13
Blazefork-----	30	Not limited		Not limited	
1324: Carway-----	50	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
Carbika-----	30	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
1553: Darlow-----	70	Not limited		Very limited Sodium content	1.00
Elmer-----	20	Not limited		Very limited Sodium content	1.00
1555: Dillhut-----	35	Very limited Too sandy	1.00	Somewhat limited Droughty	0.15

RECREATIONAL INTERPRETATIONS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Plev-----	35	Very limited Depth to saturated zone Too sandy	1.00 0.94	Very limited Depth to saturated zone Droughty	1.00 0.92
1985: Hayes-----	60	Not limited		Not limited	
2391: Kaskan-----	75	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
2588: Longford, Moderately Eroded-----	90	Not limited		Not limited	
3181: Pratt-----	45	Very limited Too sandy	1.00	Not limited	
Turon-----	30	Somewhat limited Too sandy	0.98	Not limited	
3641: Tivin-----	45	Very limited Too sandy	1.00	Somewhat limited Droughty Slope	0.98 0.16
Dillhut-----	40	Very limited Too sandy	1.00	Somewhat limited Droughty	0.15
AED: Arents, Earthen Dam-	100	Not rated		Not rated	
At: Attica-----	100	Somewhat limited Too sandy	0.96	Not limited	
Br: Bridgeport-----	100	Not limited		Not limited	
Ca: Carwile-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Cb: Cass-----	100	Not limited		Not limited	
Ce: Clime-----	100	Somewhat limited Too clayey	0.50	Very limited Too clayey Depth to bedrock Droughty	1.00 0.71 0.00
Cm: Clime-----	100	Somewhat limited Too clayey	0.50	Very limited Too clayey Depth to bedrock Droughty	1.00 0.71 0.00
Cr: Crete-----	100	Not limited		Not limited	
Cs: Crete-----	100	Not limited		Not limited	
Ct: Crete-----	100	Not limited		Not limited	
De: Detroit-----	100	Not limited		Not limited	
Dr: Drummond-----	100	Not limited		Not limited	
Ed: Edalgo-----	100	Not limited		Somewhat limited Depth to bedrock Slope	0.42 0.04
Ee: Edalgo-----	100	Not limited		Somewhat limited Depth to bedrock Droughty	0.95 0.01
Fa: Farnum-----	100	Not limited		Not limited	
Ge: Geary-----	100	Not limited		Not limited	
Go: Goessel-----	100	Somewhat limited Too clayey	0.50	Very limited Too clayey	1.00
Ho: Hord-----	100	Not limited		Not limited	
INT: Aguolls-----	100	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00

RECREATIONAL INTERPRETATIONS--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Paths and trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Ir: Irwin, bedrock substratum-----	100	Not limited		Not limited	
La: Ladysmith-----	100	Not limited		Not limited	
Ld: Ladysmith-----	70	Not limited		Not limited	
Drummond-----	30	Not limited		Not limited	
Le: Lancaster-----	100	Not limited		Somewhat limited Depth to bedrock	0.29
Lh: Lancaster-----	60	Not limited		Somewhat limited Depth to bedrock Slope	0.29 0.04
Hedville-----	40	Not limited		Very limited Depth to bedrock Droughty Slope Content of large stones	1.00 0.62 0.04 0.03
LHH: Lancaster-----	60	Not limited		Somewhat limited Depth to bedrock Slope	0.46 0.00
Hedville-----	40	Not limited		Very limited Depth to bedrock Droughty Slope Content of large stones	1.00 0.75 0.63 0.03
Ln: Longford-----	100	Not limited		Not limited	
Lo: Longford-----	100	Not limited		Not limited	
Mc: Mccook-----	100	Not limited		Not limited	
Ns: Ness-----	100	Very limited Depth to saturated zone Too clayey	1.00 0.50	Very limited Depth to saturated zone Too clayey	1.00 1.00
Nw: New Cambria-----	100	Somewhat limited Too clayey	0.50	Very limited Too clayey	1.00
Pa: Plevna-----	100	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 1.00
Pr: Pratt-----	100	Somewhat limited Too sandy	0.37	Somewhat limited Slope	0.04
Ro: Roxbury-----	100	Not limited		Not limited	
Sm: Smolan-----	100	Not limited		Not limited	
To: Tobin-----	100	Not limited		Somewhat limited Flooding	0.60
W: Water-----	100	Not rated		Not rated	
Wb: Wells-----	90	Not limited		Not limited	
Wc: Wells-----	100	Not limited		Not limited	

WILDLIFE INTERPRETATIONS
McPherson County, Kansas

Use and Explanation of Wildlife Interpretations

Soils directly affect the kind and amount of vegetation that is available to wildlife as food and cover. They also affect the development of water impoundments. The kind and abundance of wildlife that populate an area depend largely on the amount and distribution of food, cover, water, and living space. If any one of these elements is missing, inadequate, or inaccessible, wildlife will be scarce or will not inhabit the area. If the soils have the potential, wildlife habitat can be created or improved by planting appropriate vegetation, properly managing the existing plant cover, and fostering the natural establishment of desirable plants.

In the Wildlife Interpretations table, the soils in the survey area are rated according to their potential for providing habitat for various kinds of wildlife. This information can be used in planning parks, wildlife refuges, nature study areas, and other developments for wildlife; in selecting soils that are suitable for establishing, improving, or maintaining specific elements of wildlife habitat; and in determining the intensity of management needed for each element of the habitat.

Suitability Ratings

The potential of the soil is rated good, fair, poor, or very poor.

Good - means that the element of wildlife habitat or the kind of habitat is easily created, improved, or maintained. Few or no limitations affect management, and satisfactory results can be expected if the soil is used for the designated purpose.

Fair - means that the element of wildlife habitat or kind of habitat can be created, improved, or maintained in most places. Moderately intensive management is required for satisfactory results.

Poor - means that limitations are severe for the designated element or kind of wildlife habitat. Habitat can be created, improved, or maintained in most places, but management is difficult and requires intensive effort.

Very Poor - means that limitations are very severe for the designated element or kind of wildlife habitat. Habitat is difficult to create, improve, or maintain in most places, and management is difficult and requires intensive effort.

Description of Wildlife Habitat Elements

Openland habitat consists of croplands, pastures, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. The kind of wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, killdeer, cottontail rabbit, red fox, and coyote.

Woodland habitat consists of hardwood or conifers, or a mixture of these and associated grasses, legumes and wild herbaceous plants. Examples of wildlife attracted to this habitat are wild turkey, thrushes, woodpeckers, owl, tree squirrels, raccoon, and deer.

Wetland habitat consists of water-tolerant plants in open, marshy or swampy, shallow water areas. Examples of wildlife attracted to this habitat are ducks, geese, herons, bitterns, rails, kingfishers, shorebirds, muskrat, mink, and beaver.

The elements of wildlife habitat are described in the following paragraphs.

Grain and seed crops are domestic grains and seed-producing herbaceous plants. Soil properties and features that affect the growth of grain and seed crops are depth of the root zone, texture of the surface layer, available water capacity, wetness, slope, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of grain and seed crops are corn, wheat, oats, and barley.

Grasses and legumes are domestic perennial grasses and herbaceous legumes. Soil properties and features that affect the growth of grasses and legumes are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, flooding, and slope. Soil temperature and soil moisture also are considerations. Examples of grasses and legumes are fescue, lovegrass, bromegrass, clover, and alfalfa.

Wild herbaceous plants are native or naturally established grasses and forbs, including weeds. Soil properties and features that affect the growth of these plants are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of wild herbaceous plants are bluestem, goldenrod, beggarweed, wheatgrass, and grama.

Hardwood trees and woody understory produce nuts or other fruit, buds, catkins, twigs, bark, and foliage. Soil properties and features that affect the growth of hardwood trees and shrubs are depth of the root zone, available water capacity, and wetness. Examples of these plants are oak, poplar, cherry, sweetgum, apple, hawthorn, dogwood, hickory, blackberry, and blueberry. Examples of fruit-producing shrubs that are suitable for planting on soils rated good are Russian-olive, autumn-olive, and crabapple.

Coniferous plants furnish browse and seeds. Soil properties and features that affect the growth of coniferous trees, shrubs, and ground cover are depth of the root zone, available water capacity, and wetness. Examples of coniferous plants are pine, spruce, fir, cedar, and juniper.

Shrubs are bushy woody plants that produce fruit, buds, twigs, bark, and foliage. Soil properties and features that affect the growth of shrubs are depth of the root zone, available water capacity, salinity, and soil moisture. Examples of shrubs are fragrant sumac, chokecherry, American plum, sand plum, and gorden currant.

Wetland plants are annual and perennial wild herbaceous plants that grow on moist or wet sites. Submerged or floating aquatic plants are excluded. Soil properties and features affecting wetland plants are texture of the surface layer, wetness, reaction, salinity, slope, and surface stoniness. Examples of wetland plants are smartweed, wild millet, saltgrass, cordgrass, rushes, sedges, and cattails.

Shallow water areas have an average depth of less than 5 feet. Some are naturally wet areas. Others are created by dams, levees, or other water-control structures. Soil properties and features affecting shallow water areas are depth to bedrock, wetness, surface stoniness, slope, and permeability. Examples of shallow water areas are marshes, waterfowl feeding areas, and ponds.

The habitat for various kinds of wildlife is described in the following paragraphs.

Habitat for openland wildlife consists of cropland, pasture, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. Wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, cottontail, red fox and coyote.

Habitat for woodland wildlife consists of areas of deciduous and/or coniferous plants and associated grasses, legumes, and wild herbaceous plants. Wildlife attracted to these areas include wild turkey, thrushes, woodpeckers, squirrels, gray fox, raccoon, and deer.

Habitat for wetland wildlife consists of open, marshy or swampy shallow water areas. Some of the wildlife attracted to such areas are ducks, geese, herons, shore birds, muskrat, mink, and beaver.

Habitat for rangeland wildlife consists of areas of shrubs and wild herbaceous plants. Wildlife attracted to rangeland include antelope, deer, cottontail rabbit, prairie chicken, meadowlark, quail, and pheasant.

WILDLIFE INTERPRETATIONS
McPherson County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
053ED: EDALGO-----	Good	Good	Good	---	---	Fair	Very poor	Very poor	Good	---	Very poor	Good
053MB: MCCOOK-----	Good	Good	Good	Good	Fair	Good	Very poor	Very poor	Good	Fair	Very poor	Good
053MC: MCCOOK-----	Good	Good	Good	Good	Fair	Good	Very poor	Very poor	Good	Fair	Very poor	Good
079GE: GEARY-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Good
079LB: LADYSMITH-----	Fair	Good	Good	---	---	Good	Poor	Fair	Good	---	Poor	Good
115IC: IRWIN-----	Good	Good	Good	---	---	Fair	Poor	Poor	Good	---	Poor	Fair
159DP: DILLWYN-----	Poor	Fair	Good	---	---	Fair	Fair	Fair	Fair	---	Fair	Fair
PLEVNA-----	Poor	Fair	Fair	---	---	Fair	Good	Good	Fair	---	Good	Fair
159GC: GEARY-----	Fair	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Good
CLARK-----	Fair	Good	Fair	Fair	Fair	Fair	Poor	Very poor	Fair	---	Very poor	Fair
159LA: LANCASTER-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
159NF: NARON-----	Good	Good	Good	---	---	Fair	Poor	Very poor	Good	---	Very poor	Fair
159PF: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
159PR: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
CARWILE-----	Fair	Good	Good	---	---	Good	Good	Fair	Good	---	Fair	Good
159SO: SMOLAN-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
159WA: WALDECK-----	Fair	Good	Good	---	---	Good	Fair	Fair	Good	---	Fair	Good
169CE: CLIME-----	Fair	Fair	Good	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
169ED: EDALGO-----	Good	Good	Good	---	---	Fair	Very poor	Very poor	Good	---	Very poor	Good
169RO: ROXBURY-----	Good	Good	Good	Fair	Fair	Fair	Poor	Fair	Good	Fair	Poor	Fair
1191: BLAZEFORK-----	Good	Good	Fair	Good	Good	Good	Good	Poor	Fair	Good	Fair	Fair
1200: BUHLER-----	Poor	Poor	Poor	Poor	Poor	Poor	Poor	Poor	Poor	Poor	Poor	Poor
BLAZEFORK-----	Good	Good	Fair	Good	Good	Good	Good	Poor	Fair	Good	Fair	Fair
1324: CARWAY-----	Fair	Good	Good	Good	Good	Good	Good	Good	Good	Good	Good	Good
CARBIKA-----	Fair	Good	Good	Good	Good	Good	Good	Good	Good	Good	Good	Good

WILDLIFE INTERPRETATIONS--Continued
McPherson County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
1553: DARLOW-----	Fair	Fair	Poor	Fair	Poor	Poor	Good	Fair	Fair	Fair	Fair	Poor
ELMER-----	Fair	Fair	Poor	Fair	Poor	Poor	Poor	Poor	Fair	Fair	Poor	Poor
1555: DILLHUT-----	Fair	Good	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor	Fair
PLEV-----	Fair	Good	Good	Good	Good	Good	Good	Fair	Good	Good	Fair	Good
1985: HAYES-----	Fair	Fair	Good	Good	Good	Good	Poor	Very poor	Fair	Good	Very poor	Fair
2391: KASKAN-----	Good	Good	Good	Good	Fair	Good	Fair	Good	Good	Good	Fair	Poor
2588: LONGFORD-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
3181: PRATT-----	Fair	Good	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor	Fair
TURON-----	Fair	Good	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor	Fair
3641: TIVIN-----	Poor	Poor	Fair	Fair	Fair	Poor	Very poor	Very poor	Poor	Poor	Very poor	Poor
DILLHUT-----	Fair	Good	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor	Fair
AED: ARENTS, EARTHEN DAM-----	---	---	---	---	---	---	---	---	---	---	---	---
At: ATTICA-----	Fair	Fair	Good	---	---	Fair	Poor	Very poor	Fair	---	Very poor	Fair
Br: BRIDGEPORT-----	Good	Good	Good	---	---	Fair	Poor	Poor	Good	---	Poor	Fair
Ca: CARWILE-----	Fair	Good	Good	---	---	Good	Good	Fair	Good	---	Fair	Good
Cb: CASS-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
Ce: CLIME-----	Fair	Fair	Good	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Cm: CLIME-----	Fair	Fair	Good	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Cr: CRETE-----	Good	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Good
Cs: CRETE-----	Good	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Good
Ct: CRETE-----	Good	Good	Good	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor	Good
De: DETROIT-----	Good	Good	Good	---	---	Good	Poor	Poor	Good	Poor	Poor	Good
Dr: DRUMMOND-----	Poor	Fair	Fair	---	Poor	Poor	Fair	Fair	Fair	---	Fair	Poor
Ed: EDALGO-----	Good	Good	Good	---	---	Fair	Very poor	Very poor	Good	---	Very poor	Good

WILDLIFE INTERPRETATIONS--Continued
McPherson County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
Ee: EDALGO-----	Good	Good	Good	---	---	Fair	Very poor	Very poor	Good	---	Very poor	Good
Fa: FARNUM-----	Good	Good	Good	---	---	Good	Poor	Poor	Good	---	Poor	Good
Ge: GEARY-----	Good	Good	Good	Fair	Fair	Fair	Poor	Poor	Good	Fair	Poor	Good
Go: GOESSEL-----	Fair	Fair	Fair	---	---	Fair	Poor	Fair	Fair	---	Poor	Fair
Ho: HORD-----	Good	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor	Good
INT: AQUOLLS-----	---	---	---	---	---	---	---	---	---	---	---	---
Ir: IRWIN-----	Good	Good	Good	---	---	Fair	Poor	Poor	Good	---	Poor	Fair
La: LADYSMITH-----	Fair	Good	Good	---	---	Good	Poor	Fair	Good	---	Poor	Good
Ld: LADYSMITH-----	Fair	Good	Good	---	---	Good	Poor	Fair	Good	---	Poor	Good
DRUMMOND-----	Poor	Fair	Fair	---	Poor	Poor	Fair	Fair	Fair	---	Fair	Poor
Le: LANCASTER-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Lh: LANCASTER-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
HEDVILLE-----	Very poor	Poor	Poor	---	---	Poor	Very poor	Very poor	Poor	---	Very poor	Poor
LHH: LANCASTER-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
HEDVILLE-----	Very poor	Poor	Poor	---	---	Poor	Very poor	Very poor	Poor	---	Very poor	Poor
Ln: LONGFORD-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Lo: LONGFORD-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Mc: MCCOOK-----	Good	Good	Good	Good	Fair	Good	Very poor	Very poor	Good	Fair	Very poor	Good
Ns: NESS-----	Poor	Poor	Poor	---	---	Poor	Fair	Good	Poor	---	Good	Poor
Nw: NEW CAMBRIA-----	Fair	Fair	Poor	Good	Good	Fair	Poor	Poor	Fair	Good	Poor	Poor
Pa: PLEVNA-----	Poor	Fair	Fair	---	---	Fair	Good	Good	Fair	---	Good	Fair
Pr: PRATT-----	Fair	Good	Fair	---	---	Fair	Very poor	Very poor	Fair	---	Very poor	Fair
Ro: ROXBURY-----	Good	Good	Good	Fair	Fair	Fair	Poor	Fair	Good	Fair	Poor	Fair
Sm: SMOLAN-----	Good	Good	Fair	---	---	Fair	Poor	Fair	Good	---	Poor	Fair
To: TOBIN-----	Good	Good	Good	Good	Good	Good	Poor	Fair	Good	Good	Poor	Good

WILDLIFE INTERPRETATIONS--Continued
McPherson County, Kansas

Map symbol and soil name	Potential for habitat elements								Potential as habitat for--			
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Shrubs	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life	Range- land wild- life
W: WATER-----	---	---	---	---	---	---	---	---	---	---	---	---
Wb: WELLS-----	Good	Good	Good	---	---	Fair	Very poor	Very poor	Good	---	Very poor	Fair
Wc: WELLS-----	Good	Good	Good	---	---	Fair	Very poor	Very poor	Good	---	Very poor	Fair

YIELDS PER ACRE OF PASTURE AND HAYLAND
McPherson County, Kansas

Use and Explanation of Pastureland and Hayland Interpretations

This subsection provides information concerning the suitability of soils for the production of pasture and hayland. This subsection may contain pasture and hayland suitability groupings, land capability and yield estimates, yield estimates for individual grasses or legumes, or other information pertaining to the production of forage.

Pasture and Hayland Suitability Groupings

Soils are placed in pasture and hayland groups according to their suitability for the production of forage. The soils in each group are enough alike to be suited to the same grasses or legumes, to have similar limitations and hazards, to require similar management, and to have similar productivity and other responses to management. Thus, the pasture and hayland suitability group is a convenient way of grouping the soils for their management. If used, these groupings are identified and described in other reports in the subsection.

Yield Estimates

The average yields per acre that can be expected of the principal pasture or hayland crops, under a high level of management, are presented in this subsection. In any given year, yields may be higher or lower than those indicated in the tables because of variations in rainfall or other climatic factors. The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations are also considered.

Under good management, proper grazing is essential for the production of high quality forage, stand survival, and erosion control. Proper grazing helps plants maintain sufficient and generally vigorous top growth during the growing season. Brush control is essential in many areas, and weed control generally is needed. Rotation grazing and renovation are also important management practices.

The Pasture and Hayland table show yield estimates in tons per acre and animal unit months for pasture and hayland groups. An animal unit month is the amount of forage required by one animal unit (AU) for 30 days. On animal unit (AU) is one (1000 pound) mature cow and a calf up to weaning age (usually six months of age) or their equivalent. The Natural Resources Conservation Service uses 900 pounds of air dry forage as the amount needed to meet this requirement. To maintain a healthy and vigorous plant community, the degree of use should never be greater than 50 percent. Therefore only 25 percent of the total biomass grown is considered consumed by the grazing animal. Animal Unit Months can be converted to air dry pounds per acre production by multiplying the AUM by 30 days, then by 30 pounds per day, and then by four. This figure is the amount of total forage production.

Planners of management systems for individual fields or farms should consider the detailed information given in the description of each soil in the Nontechnical Description section. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay		Smooth brome grass	
	N	I	N	I	N	I
			Tons	Tons	AUM	AUM
053ED: Edalgo-----	4e	---	---	---	---	---
053MB: Mccook-----	2w	---	---	---	---	---
053MC: Mccook-----	1	---	---	---	---	---
079GE: Geary-----	3e	3e	3.00	6.50	---	---
079LB: Ladysmith-----	3e	---	3.00	---	---	---
115IC: Irwin-----	4e	---	---	---	---	---
159DP: Dillwyn-----	4w	---	---	---	---	---
Plevna-----	5w	---	---	---	---	---
159GC: Geary-----	3e	3e	2.80	6.00	---	---
Clark-----	4e	---	---	---	---	---
159LA: Lancaster-----	3e	---	---	---	---	---
159NF: Naron-----	3e	2e	3.00	6.50	---	---
159PF: Pratt-----	3e	3e	---	5.50	---	---
159PR: Pratt-----	3e	3e	---	5.50	---	---
Carwile-----	2w	---	---	---	---	---
159SO: Smolan-----	3e	---	2.00	---	---	---
159WA: Waldeck-----	3w	---	3.50	5.00	---	---
169CE: Clime-----	4e	---	---	---	---	---
169ED: Edalgo-----	4e	---	---	---	---	---
169RO: Roxbury-----	1	---	---	---	---	---
1191: Blazefork-----	2w	2s	3.50	6.00	---	---
1200: Buhler-----	2w	2w	3.00	5.00	---	---
Blazefork-----	2s	2s	3.50	6.00	---	---
1324: Carway-----	2w	---	5.00	---	7.00	---
Carbika-----	2w	---	5.00	---	7.00	---
1553: Darlow-----	4s	4s	3.00	5.00	---	---
Elmer-----	3s	3s	3.50	5.00	---	---
1555: Dillhut-----	3e	3e	---	5.50	3.00	8.00
Plev-----	5w	---	---	---	---	---

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay		Smooth brome grass	
	N	I	N	I	N	I
			Tons	Tons	AUM	AUM
1985: Hayes-----	3e	3e	3.00	6.00	4.00	9.00
2391: Kaskan-----	5w	---	---	---	---	---
2588: Longford, Moderately Eroded-----	4e	---	2.00	---	5.00	---
3181: Pratt-----	3e	3e	---	5.50	3.00	8.00
Turon-----	3e	3e	---	5.50	3.00	8.00
3641: Tivin-----	6e	---	---	---	---	---
Dillhut-----	3e	3e	---	5.50	3.00	8.00
AED: Arents, Earthen Dam-----	8	---	---	---	---	---
At: Attica-----	2e	---	3.00	6.50	---	---
Br: Bridgeport-----	1	1	3.50	6.50	---	---
Ca: Carwile-----	2w	---	---	---	---	---
Cb: Cass-----	2e	2e	3.00	6.00	---	---
Ce: Clime-----	3e	---	1.80	---	---	---
Cm: Clime-----	4e	---	1.60	---	---	---
Cr: Crete-----	2s	2s	3.20	5.50	---	---
Cs: Crete-----	2e	2e	2.90	5.30	---	---
Ct: Crete-----	2e	2e	2.90	5.30	---	---
De: Detroit-----	1	---	---	---	---	---
Dr: Drummond-----	6s	---	---	---	---	---
Ed: Edalgo-----	4e	---	---	---	---	---
Ee: Edalgo-----	4e	---	---	---	---	---
Fa: Farnum-----	2e	2e	3.00	6.50	---	---
Ge: Geary-----	2e	2e	3.40	7.00	---	---
Go: Goessel-----	2s	---	3.00	---	---	---
Ho: Hord-----	1	1	3.50	6.50	---	---
INT: Aquolls-----	5w	---	---	---	---	---
Ir: Irwin, bedrock substratum	3e	---	3.00	---	---	---

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

Map symbol and soil name	Land capability		Alfalfa hay		Smooth brome grass	
	N	I	N	I	N	I
			Tons	Tons	AUM	AUM
La: Ladysmith-----	2s	---	3.00	---	---	---
Id: Ladysmith-----	4s	---	3.00	---	---	---
Drummond-----	6s	---	---	---	---	---
Le: Lancaster-----	4e	---	---	---	---	---
Lh: Lancaster-----	6e	---	---	---	---	---
Hedville-----	7s	---	---	---	---	---
LHH: Lancaster-----	6e	---	---	---	---	---
Hedville-----	6e	---	---	---	---	---
Ln: Longford-----	3e	3e	2.50	5.50	---	---
Lo: Longford-----	4e	---	2.00	---	---	---
Mc: Mccook-----	2e	2e	2.80	6.20	---	---
Ns: Ness-----	6w	---	---	---	---	---
Nw: New Cambria-----	2s	---	---	---	---	---
Pa: Plevna-----	5w	---	---	---	---	---
Pr: Pratt-----	4e	3e	---	5.50	---	---
Ro: Roxbury-----	1	1	4.20	7.00	---	---
Sm: Smolan-----	2e	2e	3.00	6.00	---	---
To: Tobin-----	2w	---	---	---	---	---
W: Water-----	---	---	---	---	---	---
Wb: Wells-----	2e	---	---	---	---	---
Wc: Wells-----	3e	---	3.50	---	---	---

CONSERVATION TREE AND SHRUB MANAGEMENT
McPherson County, Kansas

A Conservation Tree/Shrub Suitability Group (CTSG), formerly Windbreak Suitability Group, is a physiographic unit or area having similar climatic and edaphic characteristics that control the selection and height growth of trees and shrubs.

In this table, the Conservation Tree and Shrub Grouping is expressed as a group index number. The group index for Conservation Tree and Shrub groups (CTSG) are a guide for species best suited for different kinds of soil and for prediction height, growth, and effectiveness. The groupings can be used when selection woody plants for windbreaks, wildlife plantings riparian buffers, reforestation, other environmental plantings, recreation, landscaping, wetland restoration or enhancement and critical area plantings. CTSG's are developed to assure satisfactory species selection and adaptation to specific conditions of soil, climate and physiography. CTSG's are a guide for selection species best suited for different kinds of soil and prediction height growth and effectiveness.

All soil series mapped in the state have been placed in 10 groups of similar soil characteristics. Groups 1, 2, 3, 4, 6, and 9 are further divided into subgroups. In addition, all groups provide information by Major Land Resource Areas.

Each tree or shrub species has certain climatic and physiographic limits. Within these parameters a tree or shrub may be well or poorly suited because of soil characteristics. Each tree or shrub also has definable potentials of height growth depending on the factors just mentioned. Accurate definitions of potential heights are necessary for proper windbreak planning and design.

Windbreaks protect livestock, buildings, roads and yards from wind and snow. They also protect fruit trees and gardens, and they furnish habitat for wildlife. Several rows of low-growing and high-growing broadleaf and coniferous trees and shrubs provide the most protection.

Field windbreaks are narrow plantings made at right angles to the prevailing wind and at specific intervals across the field. The interval depends on the erodibility of the soil. Field windbreaks protect cropland and crops from wind, help to keep snow on the fields, and provide food and cover for wildlife.

Environmental plantings help to beautify and screen houses and other buildings and to abate noise. The plants, mostly evergreen shrubs and trees, are closely spaced. To ensure plant survival, a healthy planting stock of suitable species should be planted properly on a well prepared site and maintained in good condition.

Windbreaks are often planted on land that did not grow trees originally. Knowledge of how trees perform on such land can be gained only by observing and recording their performance where trees have been planted and survived. The problem is compounded by the fact that many favorite windbreak species are not indigenous to the areas in which they are planted.

The Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups shows the adapted species listing for each group index number. Showing the height that locally grown trees and shrubs are expected to reach in 20 years on various soils. The estimates are based on measurements and observation of established plantings that have been given adequate care. This information should be used to determine the placement of a windbreak, the area protected and the arrangement of species.

A number of attributes are included in the CTSG species tables for each group number found in this section of the Field Office Technical Guide. These attributes were rated subjectively and assigned a relative value to further assist those unfamiliar with individual species characteristics or desirability for the intended use. Definitions and explanations can be found. Additional information on planning windbreaks and screens and planting and caring for trees and shrubs can be obtained from the local office of the Natural Resources Conservation Service or of the Cooperative Extension Service or from a commercial nursery. See part 537 of the National Forestry Manual for additional information.

In the Tree and Shrub Management table interpretive ratings are given for various aspects of forest and conservation tree and shrub management. Some rating class terms indicate the degree to which the soils are suited to a specified forest management practice. Well suited indicates that the soil has features that are favorable for the specified practice and has no limitations. Good performance can be expected, and little or no maintenance is needed. Moderately well suited indicates that the soil has features that are moderately favorable for the specified practice. One or more soil properties are less than desirable and fair performance can be expected. Some maintenance is needed. Poorly suited indicates that the soil has one or more properties that are unfavorable for the specified practice. Overcoming the unfavorable properties requires special design, extra maintenance, and costly alteration. Unsited indicates that the expected performance of the soil is unacceptable for the specified practice or that extreme measures are needed to overcome the undesirable soil properties.

The paragraphs that follow indicate the soil properties considered in rating the soils for forest and conservation tree and shrub management practices. More detailed information about the criteria used in the ratings is available in the "National Forestry Manual," which is available in local offices of the Natural Resources Conservation Service or on the Internet. Also, in the Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups.

Ratings in the columns suitability for hand planting and suitability for mechanical planting are based on slope, depth to a restrictive layer, content of sand, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, moderately well suited, poorly suited, or unsited to these methods of planting. It is assumed that necessary site preparation is completed before seedlings are planted.

Ratings in the column suitability for mechanical site preparation (surface) are based on slope, depth to a restrictive layer, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsited to this management activity. The part of the soil from the surface to a depth of about 1-foot is considered in the ratings.

Ratings in the column suitability for mechanical site preparation (deep) are based on slope, depth to a restrictive layer, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsited to this management activity. The part of the soil from the surface to a depth of about 3 feet is considered in the ratings.

Ratings in the column potential for seedling mortality are based on flooding, ponding, depth to a water table, content of lime, reaction, salinity, available water capacity, soil moisture regime, soil temperature regime, aspect, and slope. The soils are described as having a low, moderate, or high potential for seedling mortality. See the National Forestry Manual, Subpart B for criteria used in rating management concerns. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

CONSERVATION TREE AND SHRUB MANAGEMENT
McPherson County,
Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
053ED: Edalgo-----	4C	Poorly suited Stickiness	Poorly suited Stickiness Slope	Poorly suited Stickiness	Well suited	Low
053MB: Mccook-----	1	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
053MC: Mccook-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
079GE: Geary-----	3	Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
079LB: Ladysmith-----	4C	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
115IC: Irwin-----	4C	Well suited	Moderately suited Slope	Well suited	Well suited	Low
159DP: Dillwyn-----	1	Well suited	Well suited	Well suited	Well suited	Low
Plevna-----	2	Well suited	Well suited	Well suited	Unsuited Wetness	High Wetness
159GC: Geary-----	3	Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
Clark-----	8	Well suited	Moderately suited Slope	Well suited	Well suited	Moderate Lime Soil reaction
159LA: Lancaster-----	6D	Well suited	Well suited	Well suited	Well suited	Low
159NF: Naron-----	5	Well suited	Well suited	Well suited	Well suited	Low
159PF: Pratt-----	7	Well suited	Well suited	Well suited	Well suited	Low
159PR: Pratt-----	7	Well suited	Well suited	Well suited	Well suited	Low
Carwile-----	1	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	High Wetness
159SO: Smolan-----	4C	Poorly suited Stickiness	Poorly suited Stickiness Slope	Poorly suited Stickiness	Well suited	Low
159WA: Waldeck-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
169CE: Clime-----	8	Moderately suited Stickiness	Moderately suited Stickiness	Poorly suited Stickiness	Well suited	Low
169ED: Edalgo-----	4C	Moderately suited Stickiness	Moderately suited Slope Stickiness	Well suited	Well suited	Low
169RO: Roxbury-----	1	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
1191: Blazefork-----	4	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
1200: Buhler-----	9W	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Moderate Salinity
Blazefork-----	4	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
1324: Carway-----	2	Well suited	Well suited	Well suited	Well suited	High Wetness
Carbika-----	2	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	High Wetness

CONSERVATION TREE AND SHRUB MANAGEMENT
McPherson County,
Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
1553: Darlow-----	8	Well suited	Well suited	Well suited	Well suited	Moderate Available water Salinity Low
Elmer-----	8	Well suited	Well suited	Well suited	Well suited	Low
1555: Dillhut-----	7	Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	High
Plev-----	2	Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	Available water High
1985: Hayes-----	5	Well suited	Well suited	Well suited	Well suited	Wetness
2391: Kaskan-----	1	Well suited	Well suited	Well suited	Well suited	Moderate Available water
2588: Longford, Moderately Eroded-----	3	Poorly suited	Poorly suited	Poorly suited	Well suited	Low
3181: Pratt-----	7	Well suited	Well suited	Well suited	Well suited	Low
Turon-----	7	Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	Low
3641: Tivin-----	7	Moderately suited Sandiness	Moderately suited Sandiness	Well suited	Well suited	Low
Dillhut-----	7	Moderately suited Sandiness	Moderately suited Sandiness Slope	Well suited	Well suited	High
AED: Arents, Earthen Dam-		Not rated	Not rated	Not rated	Not rated	Available water
At: Attica-----	5	Well suited	Well suited	Well suited	Well suited	Not rated
Br: Bridgeport-----	1K	Well suited	Well suited	Well suited	Well suited	Low
Ca: Carwile-----	1	Well suited	Well suited	Well suited	Well suited	Low
Cb: Cass-----	1	Well suited	Well suited	Well suited	Well suited	High Wetness
Ce: Clime-----	8	Moderately suited Stickiness	Moderately suited Stickiness	Poorly suited	Well suited	Low
Cm: Clime-----	8	Moderately suited Stickiness	Moderately suited Stickiness Slope	Poorly suited	Well suited	Low
Cr: Crete-----	4C	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
Cs: Crete-----	4C	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Ct: Crete-----	4C	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
De: Detroit-----	1	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Dr: Drummond-----	9W	Well suited	Well suited	Well suited	Well suited	Low
Ed: Edalgo-----	4C	Moderately suited Stickiness	Moderately suited Stickiness Slope	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT
McPherson County,
Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Ee: Edalgo-----	4C	Poorly suited Stickiness	Stickiness Poorly suited Stickiness Slope	Poorly suited Stickiness	Well suited	Low
Fa: Farnum-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Ge: Geary-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Go: Goessel-----	4C	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
Ho: Hord-----	1	Well suited	Well suited	Well suited	Well suited	Low
INT: Aguolls-----		Well suited	Well suited	Well suited	Well suited	High Wetness Soil reaction
Ir: Irwin, bedrock substratum-----	4C	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
La: Ladysmith-----	4C	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
Ld: Ladysmith-----	4C	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
Drummond-----	9W	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Moderate Salinity
Le: Lancaster-----	6D	Well suited	Well suited	Well suited	Well suited	Low
Lh: Lancaster-----	6D	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Hedville-----	10	Well suited	Moderately suited Slope Rock fragments	Well suited	Well suited	Low
LHH: Lancaster-----	6D	Well suited	Moderately suited Slope	Well suited	Well suited	Low
Hedville-----	10	Well suited	Moderately suited Slope Rock fragments	Well suited	Well suited	Low
Ln: Longford-----	3	Moderately suited Stickiness	Moderately suited Stickiness Slope	Well suited	Well suited	Low
Lo: Longford-----	3	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
Mc: Mccook-----	1K	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Ns: Ness-----	10	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	High Wetness
Nw: New Cambria-----	1K	Poorly suited Stickiness	Poorly suited Stickiness	Poorly suited Stickiness	Well suited	Low
Pa: Plevna-----	2	Well suited	Well suited	Well suited	Unsuited Wetness	High Wetness
Pr: Pratt-----	7	Well suited	Moderately suited Slope	Well suited	Well suited	Low

CONSERVATION TREE AND SHRUB MANAGEMENT
McPherson County,
Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

Map symbol and soil name	Wind break Group	Suitability for hand planting	Suitability for mechanical planting	Suitability for mechanical site preparation (surface)	Suitability for mechanical site preparation (deep)	Potential for seedling mortality
		Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features	Rating class and limiting features
Ro: Roxbury-----	1	Well suited	Well suited	Well suited	Well suited	Moderate Soil reaction
Sm: Smolan-----	4C	Moderately suited Stickiness	Moderately suited Stickiness	Well suited	Well suited	Low
To: Tobin-----	1	Well suited	Well suited	Well suited	Well suited	Low
W: Water-----		Not rated	Not rated	Not rated	Not rated	Not rated
Wb: Wells-----	3	Well suited	Well suited	Well suited	Well suited	Low
Wc: Wells-----	3	Well suited	Moderately suited Slope	Well suited	Well suited	Low

ENGINEERING INDEX PROPERTIES
McPherson County, Kansas

Engineering Index Properties table gives the engineering classifications and the range of index properties for the layers of each soil in the survey area. Depth to the upper and lower boundaries of each layer is indicated. Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter. Loam, for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, gravelly. Textural terms are defined in the Glossary.

Classification of the soils is determined according to the Unified soil classification system (ASTM, 1998) and the system adopted by the American Association of State Highway and Transportation Officials (AASHTO, 1998). The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection. If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional refinement, the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest. The AASHTO classification for soils tested, with group index numbers in parentheses, is given in Engineering Index Properties table.

Rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in the field to weight percentage. Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field.

Liquid limit and plasticity index (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination. The estimates of particle-size distribution, liquid limit, and plasticity index are generally rounded to the nearest 5 percent. Thus, if the ranges of gradation and Atterberg limits extend a marginal amount (1 or 2 percentage points) across classification boundaries, the classification in the marginal zone is generally omitted in the table.

ENGINEERING INDEX PROPERTIES--Continued
McPherson County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
053ED: Edalgo-----	0-8	Loam	CL	A-6	0	0	95-100	85-100	75-95	50-75	30-45	10-20
	8-34	Clay	CH, CL, MH	A-7	0	0	95-100	85-100	75-100	70-90	45-70	20-45
	>34	Weathered bedrock			---	---	---	---	---	---	---	---
053MB: Mccook-----	0-12	Loam	CL, CL-ML, ML	A-4	0	0	100	100	95-100	60-100	20-35	2-10
	12-60	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	95-100	80-100	15-20	NP-10
053MC: Mccook-----	0-12	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	95-100	60-100	20-35	2-10
	12-60	Silt loam	CL, CL-ML, ML	A-4	0	0	100	100	95-100	80-100	15-20	NP-10
079GE: Geary-----	0-9	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	80-100	25-40	4-15
	9-35	Silty clay	CL	A-6, A-7	0	0	100	100	96-100	85-100	35-50	15-25
	35-60	Clay loam	CL	A-6, A-7	0	0	100	100	96-100	85-100	30-45	11-22
079LB: Ladysmith-----	0-10	Silty clay	CL	A-6, A-7	0	0	100	100	95-100	85-95	30-45	15-25
	10-45	Silty clay	CH	A-7-6	0	0	100	100	95-100	85-95	50-70	30-50
	45-60	Silty clay	CH, CL	A-7-6	0	0	100	100	95-100	85-95	40-65	25-45
115IC: Irwin-----	0-13	Silty clay	CL	A-6, A-7-6	0	0	100	95-100	90-100	80-95	35-45	15-20
	13-40	Silty clay	CH	A-7-6	0	0	100	95-100	95-100	85-95	50-60	25-30
	40-60	Silty clay	CH, CL	A-7-6	0	0	100	100	95-100	80-95	40-60	20-30
159DP: Dillwyn-----	0-9	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	70-90	5-35	0-5	NP
	9-60	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	90-100	70-90	5-35	0-5	NP
Plevna-----	0-10	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	10-20	NP
	10-33	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	30-50	15-26	NP-6
	33-60	Fine sand	SM, SP	A-2, A-3	0	0	100	90-100	50-90	4-35	0-5	NP
159GC: Geary-----	0-7	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	80-100	25-40	4-15
	7-32	Silty clay	CL	A-6, A-7	0	0	100	100	96-100	85-100	35-50	15-25
	32-60	Silty clay	CL	A-6, A-7	0	0	100	100	96-100	85-100	30-45	11-22
Clark-----	0-11	Clay loam	CL	A-6	0	0	100	95-100	90-100	50-90	30-40	10-20
	11-60	Clay loam	CL	A-6	0	0	100	95-100	90-100	55-90	25-40	10-25
159LA: Lancaster-----	0-7	Loam	CL, CL-ML	A-4, A-6	---	0-5	95-100	90-100	85-100	60-90	20-35	5-15
	7-25	Sandy clay	SC, CL	A-4, A-6, A-7-6	0	0	100	95-100	80-95	40-65	25-45	8-25
	25-35	Clay loam	CL, SC, SC-SM, CL-ML	A-4, A-6	---	0-10	95-100	90-100	80-100	36-80	20-35	5-15
	>35	Weathered bedrock			---	---	---	---	---	---	---	---
159NF: Naron-----	0-14	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	95-100	75-100	25-60	15-26	1-7
	14-40	Sandy clay	CL, SC	A-4, A-6	0	0	100	95-100	80-100	36-60	26-40	8-18
	40-60	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	75-100	20-50	15-26	NP-7
159PF: Pratt-----	0-13	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	10-20	NP
	13-38		SC-SM, SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	38-60		SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-35	0-10	NP
159PR: Pratt-----	0-13	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	10-20	NP
	13-38	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	38-60	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-35	0-10	NP
Carwile-----	0-12	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	98-100	90-100	36-60	15-26	NP-7
	12-17	Sandy clay	CL, SC	A-6, A-7	0	0	100	100	90-100	36-90	35-50	14-26
	17-60	Clay loam	CH, CL, SC	A-6, A-7	0	0	100	100	90-100	40-95	35-70	14-38
159SO: Smolan-----	0-6	Silty clay	CL	A-7	0	0	100	100	95-100	85-100	42-50	22-28
	6-60	Silty clay	CH	A-7	0	0	100	100	95-100	90-100	50-65	28-40
159WA: Waldeck-----	0-25	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	95-100	75-100	25-55	15-25	NP-5
	25-42	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	30-50	15-25	NP-5
	42-60	Sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	90-100	80-100	40-60	1-35	---	NP
169CE: Clime-----	0-8	Silty clay	CL	A-7-6	0	0-5	90-100	90-100	85-100	80-95	40-50	20-25
	8-20	Silty clay	CH, CL	A-7	0	0	95-100	95-100	95-100	85-95	45-65	20-40
	20-28	Silty clay	CH, CL	A-7	0	0	95-100	95-100	95-100	85-95	45-60	20-30
	>28	Weathered bedrock			---	---	---	---	---	---	---	---
169ED: Edalgo-----	0-9	Clay loam	CL	A-6, A-7	0	0	95-100	85-100	85-100	75-95	35-45	15-20
	9-13	Clay loam	CH, CL	A-6, A-7	0	0	95-100	85-100	75-100	65-95	35-60	15-30
	13-28	Silty clay	CH, CL	A-7	0	0	95-100	85-100	75-100	70-90	45-70	20-45
	>28	Weathered bedrock			---	---	---	---	---	---	---	---
169RO: Roxbury-----	0-26	Silt loam	CL	A-4, A-6	0	0	100	100	90-100	70-90	30-35	10-15
	26-60	Silt loam	CL, ML	A-4, A-6, A-7-6	0	0	100	100	85-100	65-95	30-45	10-20

ENGINEERING INDEX PROPERTIES--Continued
McPherson County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index	
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200			
	In				Pct	Pct					Pct		
1191: Blazefork-----	0-3	Silty clay loam	CL, CH	A-7-6	0	0	100	100	95-100	85-95	45-55	25-35	
	3-7	Silty clay loam	CL, CH	A-7-6	0	0	100	100	95-100	85-95	45-55	25-35	
	7-14	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-95	50-65	30-40	
	14-22	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-95	50-65	30-40	
	22-29	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-95	50-65	30-40	
	29-34	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-95	50-65	30-40	
	34-40	Silty clay	CL	A-7-6	0	0	100	100	85-100	75-85	40-50	20-30	
	40-48	Silty clay loam	CL	A-7-6	0	0	100	100	85-100	75-85	40-50	20-30	
	48-61	Clay loam	CL	A-7-6	0	0	100	100	85-100	75-85	40-50	20-30	
	61-80	Loam	CL	A-7-6	0	0	100	100	85-100	75-85	40-50	20-30	
1200: Buhler-----	0-3	Silty clay loam	CL, CH	A-6, A-7-6	0	0	100	100	95-100	85-95	35-55	15-30	
	3-8	Silty clay loam	CH, CL	A-6, A-7-6	0	0	100	100	95-100	85-95	35-55	15-30	
	8-12	Silt loam	CL	A-6	0	0	100	100	80-95	60-85	25-30	10-15	
	12-16	Silt loam	CL	A-6	0	0	100	100	80-95	60-85	25-30	10-15	
	16-24	Clay loam	CL	A-6, A-7-6	0	0	100	100	95-100	90-99	30-50	15-25	
	24-36	Silty clay loam	CL	A-6, A-7-6	0	0	100	100	95-100	90-99	30-50	15-25	
	36-42	Silty clay loam	CL	A-6, A-7-6	0	0	100	100	95-100	90-99	30-50	15-25	
	42-50	Clay	CL, CH	A-6, A-7-6	0	0	100	100	80-100	75-95	35-55	15-30	
	50-58	Clay loam	CH, CL	A-6, A-7-6	0	0	100	100	80-100	75-95	35-55	15-30	
	58-76	Fine sandy loam	CL-ML, CL, SC, SC-SM	A-4	0	0	100	100	55-70	40-60	20-30	5-10	
Blazefork-----	76-80	Loam	CL, CL-ML, SC, SC-SM	A-4	0	0	100	100	55-70	40-60	20-30	5-10	
	0-3	Silty clay loam	CL, CH	A-7-6	0	0	100	100	95-100	85-95	45-55	25-35	
	3-7	Silty clay loam	CH, CL	A-7-6	0	0	100	100	95-100	85-95	45-55	25-35	
	7-14	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-95	50-65	30-40	
	14-22	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-95	50-65	30-40	
	22-29	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-95	50-65	30-40	
	29-34	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-95	50-65	30-40	
	34-40	Silty clay	CL	A-7-6	0	0	100	100	85-100	75-85	40-50	20-30	
	40-48	Silty clay loam	CL	A-7-6	0	0	100	100	85-100	75-85	40-50	20-30	
	48-61	Clay loam	CL	A-7-6	0	0	100	100	85-100	75-85	40-50	20-30	
61-80	Loam	CL	A-7-6	0	0	100	100	85-100	75-85	40-50	20-30		
1324: Carway-----	0-7	Fine sandy loam	SC, CL	A-2-6, A-6	0	0	100	100	80-95	30-55	20-30	10-15	
	7-10	Sandy clay loam	SC, CL	A-6	0	0	100	100	85-100	45-60	25-35	10-15	
	10-15	Sandy clay loam	SC, CL	A-6	0	0	100	100	85-100	45-60	25-35	10-15	
	15-22	Fine sandy loam	SC, CL	A-6	0	0	100	100	85-100	45-60	25-35	10-15	
	22-35	Fine sandy loam	SC, CL	A-6	0	0	100	100	85-100	45-60	25-35	10-15	
	35-40	Clay loam	CL, CH	A-7-6	0	0	100	100	90-100	85-99	45-60	25-40	
	40-54	Clay loam	CL, CH	A-7-6	0	0	100	100	90-100	85-99	45-60	25-40	
	54-63	Clay loam	CL, CH	A-7-6	0	0	100	100	90-100	85-99	45-60	25-40	
	63-72	Clay loam	CL, CH	A-7-6	0	0	100	100	90-100	85-99	45-60	25-40	
	72-80	Clay loam	SC, CL	A-6	0	0	100	100	75-90	45-60	25-35	10-15	
Carbika-----	0-11	Silt loam	SM, SC-SM	A-2-4, A-4	0	0	100	100	90-100	30-45	20-30	1-7	
	11-15	Clay	CH, CL	A-7-6	0	0	100	100	90-100	85-99	45-55	30-35	
	15-22	Clay loam	CH, CL	A-7-6	0	0	100	100	90-100	85-99	45-55	30-35	
	22-34	Clay loam	CL, SC	A-4, A-6	0	0	100	100	85-100	45-60	30-35	10-15	
	34-41	Clay loam	CL, SC	A-4, A-6	0	0	100	100	85-100	45-60	30-35	10-15	
	41-60	Clay loam	SC, CL	A-4, A-6	0	0	100	100	85-100	45-60	30-35	10-15	
	60-80	Clay loam	SC, CL	A-4, A-6	0	0	100	100	85-100	45-60	30-35	10-15	
	1553: Darlow-----	0-5	Loam	CL-ML, CL	A-4, A-6	0	0	100	100	85-95	53-75	21-30	4-11
		5-8	Loam	CL, CL-ML	A-4, A-6	0	0	100	100	85-95	53-75	21-30	4-11
		8-14	Loam	CL	A-6	0	0	100	100	90-100	60-80	30-39	11-18
14-20		Clay loam	CL	A-6	0	0	100	100	90-100	60-80	30-39	11-18	
20-26		Loam	CL	A-6	0	0	100	100	90-100	60-80	30-39	11-18	
26-33		Loam	CL	A-6, A-7-6	0	0	100	100	90-100	55-80	30-44	11-22	
33-44		Loam	CL	A-6, A-7-6	0	0	100	100	90-100	55-80	30-44	11-22	
44-53		Loam	CL, CL-ML, SC, SC-SM	A-4, A-6	0	0	100	100	90-100	40-52	23-37	6-16	
53-68		Loam	SC-SM, CL, CL-ML, SC	A-4, A-6	0	0	100	100	90-100	40-52	23-37	6-16	
Elmer-----		68-80	Sandy loam	SM, SC-SM	A-2	0	0	100	99-100	80-90	16-32	10-18	NP-5
	0-6	Fine sandy loam	CL, CL-ML, ML	A-4	0	0	100	100	90-100	50-60	20-30	3-10	
	6-9	Fine sandy loam	CL, CL-ML, ML	A-4	0	0	100	100	90-100	50-60	20-30	3-10	
	9-19	Fine sandy loam	CL, CL-ML, ML	A-4	0	0	100	100	90-100	50-60	20-30	3-10	
	19-26	Fine sandy loam	CL, SC	A-6	0	0	100	98-100	90-100	45-60	25-35	10-20	
	26-37	Fine sandy loam	SC, CL	A-6	0	0	100	98-100	90-100	45-60	25-35	10-20	
	37-43	Loam	CL	A-6, A-7-6	0	0	99-100	98-100	90-100	65-85	30-45	15-25	
	43-51	Clay loam	CL	A-6, A-7-6	0	0	99-100	98-100	90-100	65-85	30-45	15-25	
	51-61	Fine sandy loam	CL, CL-ML, SC, SC-SM	A-2-4, A-2-6, A-4, A-6	0	0	98-100	97-100	85-95	34-55	20-30	6-16	
	61-72	Fine sandy loam	CL, CL-ML, SC, SC-SM	A-2-4, A-2-6, A-4, A-6	0	0	98-100	97-100	85-95	34-55	20-30	6-16	
72-80	Fine sandy loam	CL, CL-ML, SC, SC-SM	A-2-4, A-2-6, A-4, A-6	0	0	98-100	97-100	85-95	34-55	20-30	6-16		

ENGINEERING INDEX PROPERTIES--Continued
McPherson County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
1555: Dillhut-----	In											
	0-4	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	80-100	5-15	0-0	NP
	4-9	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	80-100	5-15	0-0	NP
	9-18	Fine sand	SM, SP-SM	A-2-4, A-3	0	0	100	100	80-100	5-15	0-0	NP
	18-26	Fine sand	SM, SP-SM	A-2-4, A-3	0	0	100	100	80-100	5-15	0-0	NP
	26-41	Fine sandy loam	CL, SC, SC-SM	A-6	0	0	100	100	80-100	36-55	30-40	10-20
	41-55	Fine sandy loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6	0	0	100	100	80-95	30-55	26-32	7-11
	55-65	Fine sandy loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6	0	0	100	100	80-95	30-55	26-32	7-11
	65-70	Fine sandy loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6	0	0	100	100	80-95	30-55	26-32	7-11
	70-80	Fine sandy loam	SP-SC, SP-SM, SC, SC-SM, SM	A-2-4	0	0	100	100	50-70	5-25	22-30	NP-10
Plev-----	0-4	Loamy fine sand	SP-SM	A-2-4, A-3	0	0	100	100	80-100	5-10	0-19	NP-3
	4-12	Fine sand	SP-SM	A-3	0	0	100	100	80-100	5-10	0-0	NP
	12-35	Fine sand	SP-SM	A-3	0	0	100	100	80-100	5-10	0-0	NP
	35-46	Fine sand	SP-SM	A-3	0	0	100	100	80-100	5-10	0-0	NP
	46-57	Fine sandy loam	SC, SC-SM	A-2-4, A-2-6, A-4, A-6	0	0	100	100	80-95	30-45	20-35	5-15
	57-75	Fine sandy loam	SC, SC-SM	A-2-4, A-2-6, A-4, A-6	0	0	100	100	80-95	30-45	20-35	5-15
1985: Hayes-----	75-80	Loamy fine sand	SP-SM	A-2-4, A-3	0	0	100	100	50-70	5-10	0-0	NP
	0-8	Fine sandy loam	SC-SM	A-2-4, A-4	0	0	100	100	80-95	30-49	20-25	4-7
	8-14	Fine sandy loam	SC, SC-SM, SM, CL-ML	A-2-4, A-4	0	0	100	100	80-95	30-55	21-28	3-10
	14-23	Fine sandy loam	SC, SC-SM, SM, CL-ML	A-4, A-2-4	0	0	100	100	80-95	30-55	21-28	3-10
	23-34	Fine sandy loam	SC, SC-SM, SM, CL-ML	A-2-4, A-4	0	0	100	100	80-95	30-55	21-28	3-10
	34-42	Fine sandy loam	SC, SC-SM, SM, CL-ML	A-2-4, A-4	0	0	100	100	80-95	30-55	21-28	3-10
	42-47	Fine sandy loam	SC, SC-SM, SM, CL-ML	A-2-4, A-4	0	0	100	100	80-95	30-55	21-28	3-10
	47-56	Sandy clay loam	CL	A-6	0	0	100	100	80-100	60-85	30-35	11-15
	56-69	Silty clay	CL, CH	A-7-6	0	0	100	100	90-100	85-99	45-55	25-35
	69-80	Clay loam	CL, CH	A-7-6	0	0	100	100	90-100	85-99	45-55	25-35
2391: Kaskan-----	0-9	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	85-100	35-45	15-20
	9-13	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	85-100	35-45	15-20
	13-17	Fine sandy loam	CL, CL-ML, SC, SC-SM	A-4	0	0	100	100	75-95	40-65	20-30	5-10
	17-21	Fine sandy loam	CL, CL-ML, SC, SC-SM	A-4	0	0	100	100	75-95	40-65	20-30	5-10
	21-27	Fine sandy loam	SC-SM, CL, CL-ML, SC	A-4	0	0	100	100	75-95	40-65	20-30	5-10
	27-43	Stratified fine sand to loamy fine sand	SM	A-2-4	0	0	100	95-100	65-85	15-30	0-0	NP
	43-57	Stratified fine sand to fine sandy loam	SC-SM, SM	A-2-4, A-4	0	0	100	95-100	70-85	30-45	0-20	NP-5
	57-80	Stratified fine sand to fine sandy loam	SC-SM, SM	A-2-4, A-4	0	0	100	95-100	70-85	30-45	0-20	NP-5
2588: Longford, Moderately Eroded-----	0-6	Silty clay loam	CL	A-7	0	0	100	95-100	90-100	85-95	45-50	25-30
	6-11	Silty clay loam	CL	A-7	0	0	100	95-100	90-100	85-95	45-50	25-30
	11-28	Silty clay	CH	A-7-6	0	0	100	95-100	85-100	75-95	50-60	30-40
	28-43	Silty clay	CH	A-7-6	0	0	100	95-100	85-100	75-95	50-60	30-40
	43-60	Silty clay loam	CL	A-7	0	0	100	95-100	85-100	75-95	45-50	25-30
	60-80	Silty clay loam	CL	A-7	0	0	100	95-100	85-100	75-95	45-50	25-30
3181: Pratt-----	0-8	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	65-100	5-35	0-14	NP
	8-24	Loamy fine sand	SM, SC-SM	A-2, A-4	0	0	100	95-100	90-100	15-40	0-20	NP-6
	24-64	Stratified fine sand to loamy fine sand	SM, SC-SM	A-2, A-4	0	0	100	95-100	90-100	15-40	0-20	NP-6
	64-80	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-35	0-14	NP
Turon-----	0-8	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	80-100	5-25	0-0	NP
	8-28	Loamy fine sand	SM, SP-SM	A-2-4	0	0	100	100	80-100	10-25	0-20	NP-3
	28-40	Stratified loamy fine sand to fine sandy loam	SP-SM, SM, SC-SM	A-2-4	0	0	100	100	80-100	10-30	0-23	NP-6
	40-58	Silty clay	CH, CL	A-6, A-7-6	0	0	100	100	80-100	80-99	36-52	16-25
	58-75	Silty clay	CL, CH	A-6, A-7-6	0	0	100	100	80-100	80-99	36-52	16-25
	75-80	Silty clay	CH, CL	A-6, A-7-6	0	0	100	100	80-100	80-99	36-52	16-25

ENGINEERING INDEX PROPERTIES--Continued
McPherson County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
3641:												
Tivin-----	0-7	Fine sand	SP-SM, SM	A-2-4, A-3	0	0	100	100	90-100	5-25	0-0	NP
	7-18	Fine sand	SP-SM, SM	A-2-4, A-3	0	0	100	100	80-100	5-25	0-0	NP
	18-80	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	80-100	5-25	0-0	NP
Dillhut-----	0-4	Fine sand	SM, SP-SM	A-2, A-3	0	0	100	100	80-100	5-15	0-0	NP
	4-9	Fine sand	SP-SM, SM	A-2, A-3	0	0	100	100	80-100	5-15	0-0	NP
	9-18	Fine sand	SM, SP-SM	A-2-4, A-3	0	0	100	100	80-100	5-15	0-0	NP
	18-26	Fine sand	SP-SM, SM	A-2-4, A-3	0	0	100	100	80-100	5-15	0-0	NP
	26-41	Fine sandy loam	CL, SC	A-6	0	0	100	100	80-100	36-55	30-40	10-20
	41-55	Fine sandy loam	SC-SM, CL, CL-ML, SC	A-2, A-4, A-6	0	0	100	100	80-95	30-55	26-32	7-11
	55-65	Fine sandy loam	SC, CL, CL- ML, SC-SM	A-2, A-4, A-6	0	0	100	100	80-95	30-55	26-32	7-11
	65-70	Fine sandy loam	SC-SM, CL, CL-ML, SC	A-2, A-4, A-6	0	0	100	100	80-95	30-55	26-32	7-11
	70-80	Fine sandy loam	SP-SM, SC, SC-SM, SM, SP-SC	A-2-4	0	0	100	100	50-70	5-25	22-30	NP-10
AED: Arents, Earthen Dam-----	---	---	---	---	---	---	---	---	---	---	---	---
At: Attica-----	0-10	Loamy fine sand	SM, SP-SM	A-2	0	0	100	95-100	70-100	10-35	10-20	NP
	10-25	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	95-100	75-100	30-55	15-26	NP-7
	25-60	Loamy fine sand	SC-SM, SM	A-2, A-4	0	0	85-100	80-100	70-100	20-50	15-26	NP-7
Br: Bridgeport----	0-14	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	65-90	20-35	4-19
	14-60	Silt loam	CL	A-4, A-6	0	0	100	100	90-100	65-100	25-40	8-20
Ca: Carwile-----	0-16	Fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0	100	98-100	90-100	36-60	15-26	NP-7
	16-21	Loam	CL, SC	A-6, A-7	0	0	100	100	90-100	36-90	35-50	14-26
	21-45	Clay loam	CH, CL, SC	A-6, A-7	0	0	100	100	90-100	40-95	35-70	14-38
	45-60	Clay loam	CH, CL, SC	A-4, A-6, A-7	0	0	100	100	90-100	36-95	25-70	7-38
Cb: Cass-----	0-7	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	85-95	20-40	15-20	NP-5
	7-51	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	85-95	20-50	15-20	NP-5
	51-60	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	95-100	95-100	50-75	5-30	0-10	NP
Ce: Clime-----	0-9	Silty clay	CH	A-7-6	0	0-5	90-100	90-100	85-100	80-95	50-60	25-35
	9-27	Silty clay	CH, CL	A-7	0	0	95-100	95-100	95-100	85-95	45-65	20-40
	>27	Unweathered bedrock			---	---	---	---	---	---	---	---
Cm: Clime-----	0-9	Silty clay	CH	A-7-6	0	0-5	90-100	90-100	85-100	80-95	50-60	25-35
	9-27	Silty clay	CH, CL	A-7	0	0	95-100	95-100	95-100	85-95	45-65	20-40
	>27	Unweathered bedrock			---	---	---	---	---	---	---	---
Cr: Crete-----	0-5	Silt loam	CL, ML	A-4, A-6	0	0	100	100	100	90-100	30-40	5-15
	5-9	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	90-100	35-50	15-30
	9-19	Silty clay loam	CH	A-7	0	0	100	100	100	90-100	50-65	25-40
	19-27	Silty clay	CH	A-7	0	0	100	100	100	90-100	50-65	25-40
	27-38	Silty clay	CH	A-7	0	0	100	100	100	90-100	50-65	25-40
	38-48	Silty clay loam	CH, CL	A-6, A-7	0	0	100	100	100	95-100	30-55	10-35
	48-80	Silty clay loam	CL, CH	A-6, A-7	0	0	100	100	100	95-100	30-55	10-35
Cs: Crete-----	0-6	Silt loam	CL, ML	A-4, A-6	0	0	100	100	100	90-100	30-40	5-15
	6-13	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	90-100	35-50	15-30
	13-43	Silty clay	CH	A-7	0	0	100	100	100	90-100	50-65	25-40
	43-60	Silty clay loam	CH, CL	A-6, A-7	0	0	100	100	100	95-100	30-55	10-35
Ct: Crete-----	0-6	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	90-100	35-50	15-30
	6-13	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	90-100	35-50	15-30
	13-35	Silty clay	CH	A-7	0	0	100	100	100	90-100	50-65	25-40
	35-60	Silty clay loam	CH, CL	A-6, A-7	0	0	100	100	100	95-100	30-55	10-35
De: Detroit-----	0-16	Silty clay loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	35-50	20-30
	16-37	Silty clay loam	CH	A-7	0	0	100	100	95-100	90-100	50-60	25-35
	37-60	Silt loam	CL, ML	A-6, A-7	0	0	100	100	95-100	85-100	25-45	10-25
Dr: Drummond-----	0-19	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	96-100	65-97	22-39	3-15
	19-60	Clay loam	CH, CL	A-6, A-7	0	0	100	100	96-100	80-98	35-60	15-35
Ed: Edalgo-----	0-6	Silt loam	CL	A-6	0	0	95-100	85-100	75-100	60-95	25-35	10-15
	6-15	Silty clay loam	CH, CL	A-6, A-7	0	0	95-100	85-100	75-100	65-95	35-60	15-30
	15-30	Silty clay loam	CH, CL	A-7	0	0	95-100	85-100	75-100	70-90	45-70	20-45
	>30	Weathered bedrock			---	---	---	---	---	---	---	---
Ee: Edalgo-----	0-8	Silty clay loam	CL	A-6, A-7	0	0	95-100	85-100	85-100	75-95	35-45	15-20
	8-23	Silty clay loam	CH, CL	A-7	0	0	95-100	85-100	75-100	70-90	45-70	20-45
	>23	Weathered bedrock			---	---	---	---	---	---	---	---

ENGINEERING INDEX PROPERTIES--Continued
McPherson County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
Fa: Farnum-----	0-12	Loam	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	60-85	20-35	5-15
	12-49	Sandy clay loam	CL, SC	A-6, A-7-6	0	0	100	100	70-100	45-80	35-50	15-30
	49-60	Fine sandy loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6	0	0	100	95-100	65-100	30-80	20-35	5-15
Ge: Geary-----	0-7	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	80-100	25-40	4-15
	7-32	Silty clay loam	CL	A-6, A-7	0	0	100	100	96-100	85-100	35-50	15-25
	32-60	Silty clay loam	CL	A-6, A-7	0	0	100	100	96-100	85-100	30-45	11-22
Go: Goessel-----	0-14	Silty clay	CH	A-7-6	0	0	100	100	95-100	85-95	50-70	30-45
	14-50	Silty clay	CH	A-7-6	0	0	100	100	95-100	85-95	50-75	30-50
	50-60	Silty clay	CH, CL	A-7-6	0	0	100	100	90-100	70-95	40-65	20-40
Ho: Hord-----	0-6	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	95-100	85-100	20-35	3-18
	6-43	Silty clay loam	CL	A-4, A-6	0	0	100	100	98-100	85-100	25-40	8-23
	43-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	100	85-100	25-40	6-21
INT: Aguolls-----	0-72	Variable			---	---	---	---	---	---	---	---
Ir: Irwin, bedrock substratum----	0-11	Silty clay loam	CL, ML	A-6, A-7-6	0	0	100	95-100	90-100	80-95	35-50	10-25
	11-42	Clay	CH	A-7-6	0	0	100	95-100	95-100	85-95	50-65	30-45
	>42	Unweathered bedrock			---	---	---	---	---	---	---	---
La: Ladysmith-----	0-8	Silty clay loam	CL	A-6, A-7	0	0	100	100	95-100	85-95	30-45	15-25
	8-48	Clay	CH	A-7-6	0	0	100	100	95-100	85-95	50-70	30-50
	48-60	Silty clay	CH, CL	A-7-6	0	0	100	100	95-100	85-95	40-65	25-45
Ld: Ladysmith-----	0-8	Silty clay loam	CL	A-6, A-7	0	0	100	100	95-100	85-95	30-45	15-25
	8-48	Clay	CH	A-7-6	0	0	100	100	95-100	85-95	50-70	30-50
	48-60	Silty clay	CH, CL	A-7-6	0	0	100	100	95-100	85-95	40-65	25-45
Drummond-----	0-7	Loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	96-100	65-97	22-39	3-15
	7-60	Clay loam	CH, CL	A-6, A-7	0	0	100	100	96-100	80-98	35-60	15-35
Le: Lancaster-----	0-10	Loam	CL, CL-ML	A-4, A-6	0	0-5	95-100	90-100	85-100	60-90	20-35	5-15
	10-32	Clay loam	CL, SC	A-4, A-6, A-7-6	0	0	100	95-100	80-95	40-65	25-45	8-25
	>32	Weathered bedrock			---	---	---	---	---	---	---	---
Lh: Lancaster-----	0-10	Loam	CL-ML, CL	A-4, A-6	0	0-5	95-100	90-100	85-100	60-90	20-35	5-15
	10-32	Clay loam	SC, CL	A-4, A-6, A-7-6	0	0	100	95-100	80-95	40-65	25-45	8-25
	>32	Weathered bedrock			---	---	---	---	---	---	---	---
Hedville-----	0-15	Loam	SM, SC, ML, CL	A-4, A-6	0	0-15	80-100	75-100	65-95	45-75	15-35	NP-13
	>15	Unweathered bedrock			---	---	---	---	---	---	---	---
LHH: Lancaster-----	0-9	Loam	CL, CL-ML	A-4, A-6	0	0-5	95-100	90-100	85-100	60-90	20-35	5-15
	9-24	Clay loam	CL, SC	A-4, A-6, A-7-6	0	0	100	95-100	80-95	40-65	25-45	8-25
	24-30	Sandy clay loam	SC, CL, CL- ML, SC-SM	A-4, A-6	0	0-10	95-100	90-100	80-100	36-80	20-35	5-15
	>30	Weathered bedrock			---	---	---	---	---	---	---	---
Hedville-----	0-8	Loam	SM, CL, ML, SC	A-4, A-6	---	0-15	80-100	75-100	65-95	45-75	15-35	NP-13
	8-17	Gravelly loam	CL, ML, SC, SM	A-1-b, A-2, A-4, A-6	---	0-15	60-90	50-85	30-80	15-60	15-35	NP-13
	>17	Unweathered bedrock			---	---	---	---	---	---	---	---
Ln: Longford-----	0-15	Silty clay loam	CL	A-7	0	0	100	95-100	90-100	85-95	45-50	25-30
	15-29	Silty clay	CL	A-6, A-7	0	0	100	95-100	90-100	70-95	30-50	15-30
	29-47	Silty clay loam	CL	A-6, A-7-6	0	0	100	95-100	85-100	70-95	35-50	15-30
	47-60	Silty clay loam	CH	A-7-6	0	0	100	95-100	90-100	75-95	50-60	30-40
Lo: Longford-----	0-6	Silty clay loam	CL	A-7	0	0	100	95-100	90-100	85-95	45-50	25-30
	6-15	Silty clay loam	CL	A-7	0	0	100	95-100	90-100	85-95	45-50	25-30
	15-47	Silty clay	CH	A-7-6	0	0	100	95-100	85-100	75-95	50-60	30-40
	47-60	Silty clay loam	CL	A-7	0	0	100	95-100	85-100	75-95	45-50	25-30
Mc: Mccook-----	0-14	Fine sandy loam	SM, SC-SM, ML, CL-ML	A-4	0	0	100	100	70-85	35-55	15-20	NP-5
	14-60	Very fine sandy loam	ML, CL-ML, CL	A-4	0	0	100	100	95-100	80-100	15-20	NP-10
Ns: Ness-----	0-60	Silty clay	CH	A-7-6	0	0	100	100	95-100	90-100	50-70	30-45
Nw: New Cambria----	0-12	Silty clay	CH, MH	A-7-6	0	0	100	100	95-100	90-100	50-75	30-45
	12-34	Silty clay	CH, MH	A-7-6	0	0	100	100	95-100	85-100	50-75	25-45
	34-60	Silty clay	CH, CL, MH	A-7-6	0	0	100	100	95-100	85-100	40-60	20-40

ENGINEERING INDEX PROPERTIES--Continued
McPherson County, Kansas

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
					Pct	Pct					Pct	
Pa: Plevna-----	In											
	0-18	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	20-50	15-26	NP-6
	18-36	Fine sandy loam	SC-SM, SM	A-2, A-4	0	0	100	95-100	70-100	30-50	15-26	NP-6
	36-60	Loamy fine sand	SM, SP	A-2, A-3	0	0	100	90-100	50-90	4-35	10-20	NP
Pr: Pratt-----												
	0-11	Loamy fine sand	SM	A-2	0	0	100	95-100	70-100	15-35	10-20	NP
	11-36	Loamy fine sand	SM, SC-SM	A-2, A-4	0	0	100	95-100	90-100	15-40	15-20	NP-6
	36-60	Loamy fine sand	SM, SP-SM	A-2, A-3	0	0	100	95-100	80-100	5-35	10-20	NP
Ro: Roxbury-----												
	0-21	Silty clay loam	CL	A-6, A-7-6	0	0	100	100	95-100	85-95	35-45	15-20
	21-50	Silty clay loam	CL	A-4, A-6, A-7-6	0	0	100	100	95-100	85-100	30-45	10-20
	50-60	Silt loam	CL, ML	A-4, A-6, A-7-6	0	0	100	100	85-100	65-95	30-45	10-20
Sm: Smolan-----												
	0-11	Silty clay loam	CL	A-7	0	0	100	100	95-100	85-100	42-50	22-28
	11-16		CL	A-6, A-7	0	0	100	100	95-100	85-100	35-50	15-28
	16-60		CH	A-7	0	0	100	100	95-100	90-100	50-65	28-40
To: Tobin-----												
	0-20	Silt loam	CL	A-6	0	0	100	100	90-100	70-90	30-35	10-15
	20-32	Silt loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	30-45	10-20
	32-60	Silt loam	CL	A-6, A-7	0	0	100	100	85-100	70-95	30-45	10-20
W: Water-----	---	---	---	---	---	---	---	---	---	---	---	---
Wb: Wells-----												
	0-15	Loam	CL	A-6	0	0	100	100	85-95	60-75	30-35	10-15
	15-36	Sandy clay loam	CL, SC	A-6, A-7	0	0	100	100	85-100	40-80	35-45	10-20
	36-60	Sandy loam	CL, ML, SC, SM	A-6, A-4	0	0	100	100	70-100	35-80	20-40	NP-15
Wc: Wells-----												
	0-11	Loam	CL	A-4, A-6	0	0	100	100	85-95	60-75	30-35	10-15
	11-15	Sandy clay loam	SC, CL	A-4, A-6, A-7	0	0	100	100	85-100	40-80	35-45	10-20
	15-49	Sandy clay loam	SC, CL	A-4, A-6, A-7	0	0	100	100	85-100	40-80	35-45	10-20
	49-60	Sandy loam	SC, ML, CL, SM	A-4, A-6	0	0	100	100	70-100	35-80	20-40	NP-15

Physical Properties table shows estimates of some physical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Particle size is the effective diameter of a soil particle as measured by sedimentation, sieving, or micrometric methods. Particle sizes are expressed as classes with specific effective diameter class limits. The broad classes are sand, silt, and clay, ranging from the larger to the smaller.

Sand as a soil separate consists of mineral soil particles that are 0.05 millimeter to 2 millimeters in diameter. In this table, the estimated sand content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Silt as a soil separate consists of mineral soil particles that are 0.002 to 0.05 millimeter in diameter. In this table, the estimated silt content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Clay as a soil separate consists of mineral soil particles that are less than 0.002 millimeter in diameter. In this table, the estimated clay content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of sand, silt, and clay affects the physical behavior of a soil. Particle size is important for engineering and agronomic interpretations, for determination of soil hydrologic qualities, and for soil classification.

The amount and kind of clay affect the fertility and physical condition of the soil and the ability of the soil to adsorb cations and to retain moisture. They influence shrink-swell potential, permeability, plasticity, the ease of soil dispersion, and other soil properties. The amount and kind of clay in a soil also affect tillage and earth moving operations.

Moist bulk density is the weight of soil (oven-dry) per unit volume. Volume is measured when the soil is at field moisture capacity, that is, the moisture content at 1/3- or 1/10-bar (33kPa or 10kPa) moisture tension. Weight is determined after the soil is dried at 105 degrees C. In the table, the estimated moist bulk density of each soil horizon is expressed in grams per cubic centimeter of soil material that is less than 2 millimeters in diameter. Bulk density data are used to compute shrink-swell potential, available water capacity, total pore space, and other soil properties. The moist bulk density of a soil indicates the pore space available for water and roots. Depending on soil texture, a bulk density of more than 1.4 can restrict water storage and root penetration. Moist bulk density is influenced by texture, kind of clay, content of organic matter, and soil structure.

Permeability ($K \rightarrow \text{sat}$) refers to the ability of a soil to transmit water or air. The term "permeability," as used in soil surveys, indicates saturated hydraulic conductivity ($K \rightarrow \text{sat}$). The estimates in the table indicate the rate of water movement, in inches per hour, when the soil is saturated. They are based on soil characteristics observed in the field, particularly structure, porosity, and texture. Permeability is considered in the design of soil drainage systems and septic tank absorption fields.

Available water capacity refers to the quantity of water that the soil is capable of storing for use by plants. The capacity for water storage is given in inches of water per inch of soil for each soil layer. The capacity varies, depending on soil properties that affect retention of water. The most important properties are the content of organic matter, soil texture, bulk density, and soil structure. Available water capacity is an important factor in the choice of plants or crops to be grown and in the design and management of irrigation systems. Available water capacity is not an estimate of the quantity of water actually available to plants at any given time.

Linear extensibility refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. It is an expression of the volume change between the water content of the clod at 1/3- or 1/10-bar tension (33kPa or 10kPa tension) and oven dryness. The volume change is reported in the table as percent change for the whole soil. Volume change is influenced by the amount and type of clay minerals in the soil.

Linear extensibility is used to determine the shrink-swell potential of soils. The shrink-swell potential is low if the soil has a linear extensibility of less than 3 percent; moderate if 3 to 6 percent; high if 6 to 9 percent; and very high if more than 9 percent. If the linear extensibility is more than 3, shrinking and swelling can cause damage to buildings, roads, and other structures and to plant roots. Special design commonly is needed.

Organic matter is the plant and animal residue in the soil at various stages of decomposition. In Physical Properties table, the estimated content of organic matter is expressed as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter. The content of organic matter in a soil can be maintained by returning crop residue to the soil. Organic matter has a positive effect on available water capacity, water infiltration, soil organism activity, and tilth. It is a source of nitrogen and other nutrients for crops and soil organisms.

Erosion factors are shown in the Physical Properties table as the K factor (K_w and K_f) and the T factor. Erosion factor K indicates the susceptibility of a soil to sheet and rill erosion by water. Factor K is one of six factors used in the Universal Soil Loss Equation (USLE) and the Revised Universal Soil Loss Equation (RUSLE) to predict the average annual rate of soil loss by sheet and rill erosion in tons per acre per year. The estimates are based primarily on percentage of silt, sand, and organic matter and on soil structure and permeability. Values of K range from 0.02 to 0.69. Other factors being equal, the higher the value, the more susceptible the soil is to sheet and rill erosion by water.

Erosion factor K_w indicates the erodibility of the whole soil. The estimates are modified by the presence of rock fragments.

Erosion factor K_f indicates the erodibility of the fine-earth fraction, or the material less than 2 millimeters in size.

Erosion factor T is an estimate of the maximum average annual rate of soil erosion by wind or water that can occur without affecting crop productivity over a sustained period. The rate is in tons per acre per year.

Wind erodibility groups are made up of soils that have similar properties affecting their susceptibility to

wind erosion in cultivated areas. The soils assigned to group 1 are the most susceptible to wind erosion, and those assigned to group 8 are the least susceptible. The groups are as follows:

1. Coarse sands, sands, fine sands, and very fine sands.
2. Loamy coarse sands, loamy sands, loamy fine sands, loamy very fine sands, ash material, and sapric soil material.
3. Coarse sandy loams, sandy loams, fine sandy loams, and very fine sandy loams.
- 4L. Calcareous loams, silt loams, clay loams, and silty clay loams.
4. Clays, silty clays, noncalcareous clay loams, and silty clay loams that are more than 35 percent clay.
5. Noncalcareous loams and silt loams that are less than 20 percent clay and sandy clay loams, sandy clays, and hemic soil material.
6. Noncalcareous loams and silt loams that are more than 20 percent clay and noncalcareous clay loams that are less than 35 percent clay.
7. Silts, noncalcareous silty clay loams that are less than 35 percent clay, and fibric soil material.
8. Soils that are not subject to wind erosion because of coarse fragments on the surface or because of surface wetness.

Wind erodibility index is a numerical value indicating the susceptibility of soil to wind erosion, or the tons per acre per year that can be expected to be lost to wind erosion. There is a close correlation between wind erosion and the texture of the surface layer, the size and durability of surface clods, rock fragments, organic matter, and a calcareous reaction. Soil moisture and frozen soil layers also influence wind erosion.

Explanation of Wind Erodibility Groups

Soil erodibility by wind is directly related to the percentage of dry non-erodible surface soil aggregates larger than 0.84 mm in diameter. From this percentage, the wind erodibility index (I-factor) is determined. The I-factor is an expression of the stability of these soil aggregates against breakdown by tillage and abrasion from wind erosion. Soils are placed in Wind Erodibility Groups (WEG) having similar percentages of dry soil aggregates larger than 0.84 mm as shown in the following table.

WEG	Properties of Soil Surface Layer	Dry Soil Aggregates >0.84mm Percent	Wind Erodibility Index T/Ac/Yr (I)
1	Very fine sand, fine sand, sand, or coarse sand	1 2 3 5 7	310 1/ 250 220 180 160
2	Loamy very fine sand, loamy fine sand, loamy sand, loamy coarse sand, organic soil materials.	10	134
3	Very fine sandy loam, fine sandy loam, sandy loam, or coarse sandy loam.	25	86
4	Clay, silty clay, non-calcareous clay loam, or silty clay loam with >35 percent clay content.	25	86
4L	Calcareous 2/ loam, silt loam, clay loam, or silty clay loam.	25	86
5	Non-calcareous loam and silt loam with <20 percent clay content, or sandy clay loam, sandy clay, and hemic 3/ organic soil materials.	40	56
6	Non-calcareous loam and silt loam with >20 percent clay content, or non-calcareous clay loam with <35 percent clay content.	45	48
7	Silt, non-calcareous silty clay loam with >35 percent clay content and fibric 3/ organic soil material.	50	38
8	Soils not suitable for cultivation due to coarse fragments or wetness; wind erosion is not a problem.	--	0

1/ The "I" values for WEG 1 vary from 160 for coarse sands to 310 for very fine sands. Use an "I" of 220 as an average figure. For coarser sand that has gravel, use a lower figure. For a soil that has no gravel and very fine sand, use a higher figure. (Modification for coarse fragments is preparation.)

2/ Calcareous is a strongly or violently effervescent reaction to cold dilute (1N) HCL.

3/ See Soil Taxonomy for definition.

PHYSICAL PROPERTIES OF THE SOILS--Continued
 McPherson County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
053ED: Edalgo-----	0-8	38	36	18-35	1.30-1.40	0.60-2.00	0.18-0.22	0.0-2.9	2.0-4.0	.32	.32	3	6	48
	8-34	22	28	35-65	1.40-1.60	0.00-0.06	0.10-0.18	6.0-8.9	1.0-2.0	.37	.37			
	>34			---	---	---	---	---	---	---	---			
053MB: Mccook-----	0-12	41	42	15-20	1.20-1.40	0.60-2.00	0.20-0.24	0.0-2.9	2.0-4.0	.32	.32	5	4L	86
	12-60	14	72	10-18	1.30-1.45	0.60-2.00	0.17-0.20	0.0-2.9	---	.43	.43			
053MC: Mccook-----	0-12	14	69	15-20	1.20-1.40	0.60-2.00	0.20-0.24	0.0-2.9	2.0-4.0	.32	.32	5	4L	86
	12-60	14	72	10-18	1.30-1.45	0.60-2.00	0.17-0.20	0.0-2.9	---	.43	.43			
079GE: Geary-----	0-9	11	68	15-27	1.30-1.40	0.60-2.00	0.22-0.24	0.0-2.9	1.0-4.0	.32	.32	5	6	48
	9-35	7	62	27-35	1.35-1.50	0.60-2.00	0.17-0.20	3.0-5.9	---	.43	.43			
	35-60	28	46	20-32	1.30-1.40	0.60-2.00	0.15-0.19	3.0-5.9	---	.43	.43			
079LB: Ladysmith----	0-10	20	48	28-35	1.35-1.45	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	5	7	38
	10-45	5	45	40-60	1.35-1.50	0.00-0.06	0.10-0.15	6.0-8.9	1.0-3.0	.37	.37			
	45-60	7	48	35-55	1.40-1.60	0.00-0.60	0.10-0.19	3.0-5.9	0.0-1.0	.37	.37			
115IC: Irwin-----	0-13	3-10	40-65	28-35	1.35-1.45	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	5	7	38
	13-40	2-8	35-55	40-60	1.40-1.50	0.00-0.06	0.10-0.13	6.0-8.9	1.0-3.0	.28	.28			
	40-60	2-8	35-55	35-55	1.40-1.50	0.06-0.20	0.09-0.19	6.0-8.9	0.5-2.0	.32	.32			
159DP: Dillwyn-----	0-9	79	16	2-8	1.50-1.60	5.95-19.98	0.08-0.12	0.0-2.9	0.0-2.0	.17	.17	5	2	134
	9-60	79	16	2-8	1.50-1.60	5.95-19.98	0.06-0.10	0.0-2.9	---	.17	.17			
	0-10	79	16	2-8	1.40-1.50	2.00-6.00	0.10-0.12	0.0-2.9	1.0-4.0	.17	.17	5	2	134
	10-33	67	20	8-18	1.40-1.50	2.00-6.00	0.12-0.16	0.0-2.9	---	.20	.20			
	33-60	95	1	1-7	1.50-1.60	2.00-6.00	0.05-0.07	0.0-2.9	---	.20	.20			
159GC: Geary-----	0-7	11	68	15-27	1.30-1.40	0.60-2.00	0.22-0.24	0.0-2.9	1.0-4.0	.32	.32	5	6	48
	7-32	7	62	27-35	1.35-1.50	0.60-2.00	0.17-0.20	3.0-5.9	---	.43	.43			
	32-60	7	67	20-32	1.30-1.40	0.60-2.00	0.15-0.19	3.0-5.9	---	.43	.43			
Clark-----	0-11	34	37	27-32	1.35-1.45	0.60-2.00	0.17-0.22	3.0-5.9	1.0-2.0	.28	.28	5	4L	86
	11-60	38	36	18-35	1.35-1.70	0.60-2.00	0.14-0.19	3.0-5.9	---	.28	.28			
159LA: Lancaster----	0-7			12-26	1.35-1.45	0.60-2.00	0.17-0.22	0.0-2.9	1.0-4.0	.28	.28	3	6	48
	7-25			18-35	1.35-1.50	0.60-2.00	0.15-0.19	3.0-5.9	1.0-3.0	.28	.32			
	25-35			12-30	1.40-1.55	0.60-2.00	0.15-0.19	0.0-2.9	0.5-2.0	.28	.32			
	>35			---	---	---	---	---	0.0-0.0	---	---			
159NF: Naron-----	0-14	63	26	8-14	1.40-1.50	2.00-6.00	0.14-0.18	0.0-2.9	1.0-3.0	.20	.20	5	3	86
	14-40	60	18	18-27	1.45-1.55	0.60-2.00	0.15-0.18	0.0-2.9	---	.32	.32			
	40-60	65	27	2-14	1.50-1.60	2.00-6.00	0.10-0.15	0.0-2.9	---	.32	.32			
159PF: Pratt-----	0-13	79	16	2-8	1.40-1.55	5.95-19.98	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	13-38	86	7	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	---	.17	.17			
	38-60	79	16	1-8	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	---	.17	.17			
159PR: Pratt-----	0-13	79	16	2-8	1.40-1.55	5.95-19.98	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	13-38	86	7	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	---	.17	.17			
	38-60	95	1	1-8	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	---	.17	.17			
Carwile-----	0-12	62	26	5-18	1.30-1.65	0.60-2.00	0.11-0.20	0.0-2.9	1.0-3.0	.24	.24	5	3	86
	12-17	54	14	25-39	1.45-1.75	0.20-2.00	0.12-0.20	3.0-5.9	---	.37	.37			
	17-60	25	27	35-60	1.35-1.75	0.06-0.20	0.12-0.20	6.0-8.9	---	.37	.37			
159SO: Smolan-----	0-6	20	49	27-35	1.30-1.40	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	5	7	38
	6-60	8	50	35-50	1.30-1.45	0.06-0.20	0.12-0.18	6.0-8.9	---	.37	.37			
159WA: Waldeck-----	0-25	68	20	8-16	1.50-1.60	2.00-6.00	0.14-0.18	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	25-42	68	20	8-16	1.50-1.60	2.00-6.00	0.12-0.17	0.0-2.9	---	.20	.20			
	42-60	96	2	1-4	1.55-1.65	5.95-19.98	0.05-0.07	0.0-2.9	---	.20	.24			
169CE: Clime-----	0-8	8	56	32-40	1.35-1.45	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	3	4	86
	8-20	6	47	35-60	1.35-1.50	0.06-0.20	0.12-0.18	3.0-5.9	1.0-4.0	.28	.28			
	20-28	8	50	35-50	1.40-1.50	0.06-0.20	0.10-0.14	3.0-5.9	1.0-3.0	.32	.32			
	>28			---	---	---	---	---	---	---	---			
169ED: Edalgo-----	0-9	35	33	28-37	1.30-1.40	0.00-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	3	7	38
	9-13	28	30	28-55	1.35-1.50	0.20-0.60	0.13-0.23	3.0-5.9	1.0-3.0	.37	.37			
	13-28	5	45	35-65	1.40-1.60	0.00-0.06	0.10-0.18	6.0-8.9	1.0-2.0	.37	.37			
	>28			---	---	---	---	---	---	---	---			
169RO: Roxbury-----	0-26	10	68	18-27	1.30-1.45	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.32	.32	5	4L	86
	26-60	9	64	18-35	1.35-1.50	0.60-2.00	0.17-0.22	3.0-5.9	0.5-0.5	.43	.43			

PHYSICAL PROPERTIES OF THE SOILS--Continued
McPherson County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
1191: Blazefork----	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
	0-3	6	58	35-50	1.20-1.45	0.20-0.60	0.21-0.23	6.0-8.9	2.0-4.0	.37	.37	5	7	38
	3-7	6	57	35-50	1.35-1.45	0.20-0.60	0.21-0.23	6.0-8.9	2.0-4.0	.37	.37			
	7-14	7	50	35-50	1.25-1.55	0.06-0.20	0.11-0.14	6.0-8.9	1.0-2.0	.43	.43			
	14-22	6	50	35-50	1.25-1.55	0.06-0.20	0.11-0.14	6.0-8.9	1.0-2.0	.43	.43			
	22-29	7	48	35-50	1.25-1.55	0.06-0.20	0.11-0.14	6.0-8.9	1.0-2.0	.43	.43			
	29-34	8	48	35-50	1.25-1.55	0.06-0.20	0.11-0.14	6.0-8.9	0.5-2.0	.43	.43			
	34-40	8	50	35-50	1.30-1.55	0.20-0.60	0.15-0.18	3.0-5.9	0.5-1.0	.32	.32			
	40-48	16	49	35-50	1.35-1.55	0.20-0.60	0.15-0.18	3.0-5.9	0.2-1.0	.32	.32			
	48-61	23	48	26-35	1.35-1.55	0.20-0.60	0.15-0.18	3.0-5.9	0.1-1.0	.32	.32			
	61-80	31	43	18-27	1.35-1.55	0.20-0.60	0.15-0.18	3.0-5.9	0.1-1.0	.32	.32			
1200: Buhler-----	0-3	10	50	28-45	1.20-1.55	0.00-0.06	0.18-0.21	6.0-8.9	2.0-6.0	.43	.43	2	7	38
	3-8	9	51	28-45	1.20-1.55	0.00-0.06	0.18-0.21	6.0-8.9	2.0-4.0	.43	.43			
	8-12	31	54	15-26	1.30-1.55	0.20-0.60	0.20-0.22	0.0-2.9	1.0-2.0	.28	.28			
	12-16	25	52	15-26	1.30-1.55	0.20-0.60	0.20-0.22	0.0-2.9	1.0-2.0	.28	.28			
	16-24	20	49	20-40	1.25-1.55	0.00-0.06	0.14-0.20	6.0-8.9	1.0-2.0	.43	.43			
	24-36	16	47	20-40	1.25-1.55	0.00-0.06	0.14-0.20	6.0-8.9	0.5-2.0	.43	.43			
	36-42	16	46	20-40	1.30-1.55	0.00-0.06	0.14-0.20	6.0-8.9	0.5-2.0	.43	.43			
	42-50	22	38	27-45	1.30-1.60	0.00-0.06	0.14-0.18	6.0-8.9	0.3-2.0	.37	.37			
	50-58	44	28	27-45	1.50-1.60	0.00-0.06	0.14-0.18	6.0-8.9	0.0-2.0	.37	.37			
	58-76	68	15	10-26	1.35-1.70	0.60-2.00	0.14-0.17	0.0-2.9	0.0-0.5	.24	.24			
	76-80	51	28	10-26	1.35-1.70	0.60-2.00	0.14-0.17	0.0-2.9	0.0-0.5	.24	.24			
Blazefork----	0-3	6	58	35-50	1.20-1.45	0.20-0.60	0.21-0.23	6.0-8.9	2.0-4.0	.37	.37	5	7	38
	3-7	6	57	35-50	1.35-1.45	0.20-0.60	0.21-0.23	6.0-8.9	2.0-4.0	.37	.37			
	7-14	7	50	35-50	1.25-1.55	0.06-0.20	0.11-0.14	6.0-8.9	1.0-2.0	.43	.43			
	14-22	6	50	35-50	1.25-1.55	0.06-0.20	0.11-0.14	6.0-8.9	1.0-2.0	.43	.43			
	22-29	7	48	35-50	1.25-1.55	0.06-0.20	0.11-0.14	6.0-8.9	1.0-2.0	.43	.43			
	29-34	8	48	35-50	1.25-1.55	0.06-0.20	0.11-0.14	6.0-8.9	0.5-2.0	.43	.43			
	34-40	8	50	35-50	1.30-1.55	0.20-0.60	0.15-0.18	3.0-5.9	0.5-1.0	.32	.32			
	40-48	16	49	35-50	1.35-1.55	0.20-0.60	0.15-0.18	3.0-5.9	0.2-1.0	.32	.32			
	48-61	23	48	26-35	1.35-1.55	0.20-0.60	0.15-0.18	3.0-5.9	0.1-1.0	.32	.32			
	61-80	31	43	18-27	1.35-1.55	0.20-0.60	0.15-0.18	3.0-5.9	0.1-1.0	.32	.32			
1324: Carway-----	0-7	67	20	10-16	1.45-1.55	2.00-6.00	0.11-0.15	0.0-2.9	0.5-1.0	.20	.20	5	3	86
	7-10	61	18	20-29	1.40-1.60	0.60-2.00	0.15-0.18	0.0-2.9	0.5-1.0	.28	.28			
	10-15	61	18	20-29	1.40-1.60	0.60-2.00	0.15-0.18	0.0-2.9	0.5-1.0	.28	.28			
	15-22	62	19	18-29	1.40-1.60	0.60-2.00	0.15-0.18	0.0-2.9	0.5-1.0	.28	.28			
	22-35	62	19	18-29	1.40-1.60	0.60-2.00	0.15-0.18	0.0-2.9	0.5-1.0	.28	.28			
	35-40	34	37	28-45	1.40-1.60	0.00-0.06	0.10-0.17	6.0-8.9	0.0-0.5	.37	.37			
	40-54	33	32	30-45	1.40-1.60	0.00-0.06	0.10-0.17	6.0-8.9	0.0-0.5	.37	.37			
	54-63	29	31	30-45	1.40-1.60	0.00-0.06	0.10-0.17	6.0-8.9	0.0-0.5	.37	.37			
	63-72	30	32	30-45	1.40-1.60	0.00-0.06	0.10-0.17	6.0-8.9	0.0-0.5	.37	.37			
	72-80	35	33	15-34	1.45-1.65	0.60-2.00	0.13-0.18	0.0-2.9	0.0-0.5	.28	.28			
Carbika-----	0-11	27	55	10-22	1.45-1.55	0.60-2.00	0.15-0.18	0.0-2.9	1.0-2.0	.24	.24	5	5	56
	11-15	30	30	35-42	1.40-1.60	0.00-0.06	0.10-0.17	6.0-8.9	0.0-0.5	.37	.37			
	15-22	30	32	35-42	1.40-1.60	0.00-0.06	0.10-0.17	6.0-8.9	0.0-0.5	.37	.37			
	22-34	34	32	21-35	1.40-1.60	0.60-2.00	0.15-0.18	0.0-2.9	0.5-1.0	.28	.28			
	34-41	34	32	21-35	1.40-1.60	0.60-2.00	0.15-0.18	0.0-2.9	0.5-1.0	.28	.28			
	41-60	35	33	21-35	1.40-1.60	0.60-2.00	0.15-0.18	0.0-2.9	0.5-1.0	.28	.28			
	60-80	34	32	21-35	1.40-1.60	0.60-2.00	0.15-0.18	0.0-2.9	0.5-1.0	.28	.28			
1553: Darlow-----	0-5	42	48	8-20	1.30-1.55	0.60-2.00	0.20-0.22	0.0-2.9	1.0-3.0	.43	.43	2	5	56
	5-8	36	53	8-20	1.30-1.70	0.60-2.00	0.20-0.22	0.0-2.9	1.0-3.0	.32	.32			
	8-14	32	47	20-30	1.30-1.45	0.20-0.60	0.09-0.13	3.0-5.9	1.0-3.0	.28	.28			
	14-20	26	46	20-30	1.30-1.45	0.20-0.60	0.09-0.13	3.0-5.9	0.8-3.0	.28	.28			
	20-26	30	44	20-30	1.30-1.45	0.20-0.60	0.09-0.13	3.0-5.9	0.2-3.0	.28	.28			
	26-33	34	42	20-35	1.30-1.50	0.00-0.06	0.09-0.13	0.0-2.9	0.0-2.0	.28	.28			
	33-44	38	38	20-35	1.30-1.50	0.00-0.06	0.09-0.13	0.0-2.9	0.0-2.0	.28	.28			
	44-53	39	36	12-27	1.30-1.60	0.20-0.60	0.10-0.16	0.0-2.9	0.0-1.0	.24	.24			
	53-68	49	35	12-27	1.30-1.80	0.20-0.60	0.10-0.16	0.0-2.9	0.0-1.0	.24	.24			
	68-80	75	14	8-12	1.50-1.80	0.60-2.00	0.02-0.10	0.0-2.9	0.0-0.5	.20	.20			
Elmer-----	0-6	53	36	10-17	1.50-1.70	2.00-6.00	0.17-0.20	0.0-2.9	1.0-2.0	.32	.32	2	3	86
	6-9	51	34	10-17	1.50-1.70	2.00-6.00	0.17-0.20	0.0-2.9	1.0-2.0	.28	.28			
	9-19	53	30	10-17	1.50-1.75	2.00-6.00	0.17-0.20	0.0-2.9	1.0-2.0	.28	.28			
	19-26	60	21	17-25	1.55-1.65	0.20-0.60	0.12-0.16	0.0-2.9	0.5-1.0	.32	.32			
	26-37	65	18	17-25	1.55-1.70	0.20-0.60	0.12-0.16	0.0-2.9	0.0-1.0	.32	.32			
	37-43	40	35	20-28	1.50-1.60	0.06-0.20	0.12-0.16	6.0-8.9	0.0-0.5	.37	.37			
	43-51	25	47	20-28	1.50-1.60	0.06-0.20	0.12-0.16	6.0-8.9	0.0-0.5	.37	.37			
	51-61	52	28	14-22	1.60-1.80	0.20-0.60	0.10-0.16	0.0-2.9	0.0-0.5	.28	.28			
	61-72	68	17	14-22	1.60-1.80	0.20-0.60	0.10-0.16	0.0-2.9	0.0-0.5	.28	.28			
	72-80	72	14	14-20	1.60-1.80	0.20-0.60	0.10-0.16	0.0-2.9	0.0-0.5	.28	.28			

PHYSICAL PROPERTIES OF THE SOILS--Continued
McPherson County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
1555: Dillhut-----	0-4	92	6	1-3	1.40-1.55	6.00-19.99	0.02-0.10	0.0-2.9	0.0-1.0	.15	.15	5	1	220
	4-9	95	4	1-3	1.40-1.55	5.95-19.98	0.02-0.10	0.0-2.9	0.0-1.0	.15	.15			
	9-18	96	3	1-3	1.40-1.55	5.95-19.98	0.02-0.10	0.0-2.9	0.0-0.5	.15	.15			
	18-26	93	5	1-3	1.40-1.55	5.95-19.98	0.02-0.10	0.0-2.9	0.0-0.5	.15	.15			
	26-41	70	16	13-30	1.40-1.55	0.60-2.00	0.12-0.15	0.0-2.9	0.0-0.0	.24	.24			
	41-55	75	15	10-22	1.55-1.65	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.0	.17	.17			
	55-65	77	13	9-17	1.55-1.65	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.0	.17	.17			
	65-70	57	25	10-20	1.55-1.65	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.0	.17	.17			
	70-80	76	15	5-15	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	0.0-0.0	.15	.15			
Plev-----	0-4	86	10	3-8	1.40-1.55	6.00-19.99	0.02-0.10	0.0-2.9	0.0-1.0	.17	.17	5	2	134
	4-12	88	10	1-5	1.40-1.55	5.95-19.98	0.02-0.10	0.0-2.9	0.0-0.5	.15	.15			
	12-35	94	5	0-2	1.40-1.55	5.95-19.98	0.02-0.10	0.0-2.9	0.0-0.0	.10	.10			
	35-46	96	4	0-2	1.40-1.55	5.95-19.98	0.02-0.10	0.0-2.9	0.0-0.0	.10	.10			
	46-57	70	14	10-27	1.55-1.65	0.60-2.00	0.08-0.10	0.0-2.9	0.0-0.0	.20	.20			
	57-75	79	9	10-27	1.55-1.65	0.60-2.00	0.08-0.10	0.0-2.9	0.0-0.0	.20	.20			
	75-80	84	11	4-6	1.45-1.60	5.95-19.98	0.04-0.10	0.0-2.9	0.0-0.0	.15	.15			
1985: Hayes-----	0-8	63	26	9-13	1.45-1.55	2.00-6.00	0.11-0.15	0.0-2.9	0.5-1.0	.20	.20	5	3	86
	8-14	65	20	8-17	1.45-1.55	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.5	.24	.24			
	14-23	65	19	8-17	1.45-1.55	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.5	.24	.24			
	23-34	65	20	8-17	1.45-1.55	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.5	.24	.24			
	34-42	67	20	8-17	1.45-1.55	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.5	.24	.24			
	42-47	66	20	8-17	1.45-1.55	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.5	.24	.24			
	47-56	61	18	19-28	1.40-1.60	0.20-0.60	0.15-0.18	0.0-2.9	0.0-0.5	.28	.28			
	56-69	8	50	28-45	1.40-1.60	0.06-0.20	0.10-0.17	6.0-8.9	0.0-0.5	.37	.37			
	69-80	34	37	28-45	1.40-1.60	0.06-0.20	0.10-0.17	6.0-8.9	0.0-0.5	.37	.37			
2391: Kaskan-----	0-9	17	52	27-35	1.35-1.45	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	5	7	38
	9-13	14	55	27-35	1.35-1.45	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37			
	13-17	60	26	10-17	1.45-1.55	2.00-6.00	0.14-0.18	0.0-2.9	0.0-1.0	.24	.24			
	17-21	62	25	10-17	1.45-1.55	2.00-6.00	0.14-0.18	0.0-2.9	0.0-1.0	.24	.24			
	21-27	65	23	10-17	1.45-1.55	2.00-6.00	0.14-0.18	0.0-2.9	0.0-1.0	.24	.24			
	27-43	84	13	0-5	1.50-1.60	5.95-19.98	0.06-0.09	0.0-2.9	0.0-0.5	.10	.10			
	43-57	82	10	1-15	1.45-1.55	1.98-19.98	0.07-0.11	0.0-2.9	0.0-1.0	.10	.10			
	57-80	83	10	1-15	1.45-1.55	1.98-19.98	0.07-0.11	0.0-2.9	0.0-1.0	.10	.10			
2588: Longford, Moderately Eroded-----	0-6			27-35	1.30-1.40	0.20-0.60	0.21-0.23	3.0-5.9	1.0-2.0	.37	.37	5	7	38
	6-11			27-35	1.30-1.40	0.20-0.60	0.21-0.23	3.0-5.9	0.5-0.9	.32	.32			
	11-28			35-45	1.35-1.50	0.00-0.60	0.11-0.20	6.0-8.9	0.5-1.0	.43	.43			
	28-43			35-45	1.35-1.50	0.00-0.60	0.11-0.20	6.0-8.9	0.5-1.0	.43	.43			
	43-60			27-35	1.30-1.40	0.20-0.60	0.14-0.20	3.0-5.9	0.0-0.5	.32	.32			
	60-80			27-35	1.30-1.40	0.20-0.60	0.14-0.20	3.0-5.9	0.0-0.5	.32	.32			
3181: Pratt-----	0-8	90	4	1-7	1.40-1.55	6.00-19.99	0.07-0.09	0.0-2.9	0.5-1.0	.15	.15	5	1	220
	8-24	87	3	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	0.0-0.5	.17	.17			
	24-64	89	3	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	0.0-0.5	.17	.17			
	64-80	89	4	1-8	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	0.0-0.5	.17	.17			
Turon-----	0-8	88	8	1-5	1.40-1.55	6.00-19.99	0.02-0.10	0.0-2.9	0.0-1.0	.15	.15	5	1	220
	8-28	88	4	3-10	1.40-1.55	5.95-19.98	0.09-0.12	0.0-2.9	0.5-1.0	.17	.17			
	28-40	88	3	2-18	1.40-1.55	1.98-19.98	0.10-0.15	0.0-2.9	0.0-0.0	.24	.24			
	40-58	11	49	27-41	1.45-1.60	0.00-0.60	0.12-0.18	3.0-5.9	0.0-0.5	.32	.32			
	58-75	6	53	27-45	1.45-1.60	0.00-0.60	0.12-0.18	3.0-5.9	0.0-0.5	.32	.32			
	75-80	4	54	27-45	1.45-1.60	0.00-0.60	0.12-0.18	3.0-5.9	0.0-0.5	.32	.32			
3641: Tivin-----	0-7	97	1	0-2	1.35-1.50	6.00-19.98	0.07-0.09	0.0-2.9	0.0-1.0	.15	.15	5	1	220
	7-18	97	1	0-2	1.35-1.50	5.95-19.98	0.02-0.08	0.0-2.9	0.0-0.5	.10	.10			
	18-80	97	0	0-3	1.50-1.70	5.95-19.98	0.02-0.08	0.0-2.9	0.0-0.0	.10	.10			
Dillhut-----	0-4	92	6	1-3	1.40-1.55	6.00-19.99	0.02-0.10	0.0-2.9	0.0-1.0	.15	.15	5	1	220
	4-9	95	4	1-3	1.40-1.55	5.95-19.98	0.02-0.10	0.0-2.9	0.0-1.0	.15	.15			
	9-18	96	3	1-3	1.40-1.55	5.95-19.98	0.02-0.10	0.0-2.9	0.0-0.5	.15	.15			
	18-26	93	5	1-3	1.40-1.55	5.95-19.98	0.02-0.10	0.0-2.9	0.0-0.5	.15	.15			
	26-41	70	16	13-30	1.40-1.55	0.60-2.00	0.12-0.15	0.0-2.9	0.0-0.0	.24	.24			
	41-55	75	15	10-21	1.55-1.65	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.0	.17	.17			
	55-65	77	13	9-17	1.55-1.65	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.0	.17	.17			
	65-70	57	25	10-20	1.55-1.65	2.00-6.00	0.11-0.15	0.0-2.9	0.0-0.0	.17	.17			
	70-80	76	15	5-15	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	0.0-0.0	.15	.15			
AED: Arents, Earthen Dam- At:	---			---	---	---	---	---	---	---	---	-	---	---
Attica-----	0-10	87	7	2-10	1.50-1.60	2.00-6.00	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	10-25	67	20	8-18	1.50-1.60	2.00-6.00	0.12-0.17	0.0-2.9	---	.24	.24			
	25-60	88	1	4-18	1.50-1.60	2.00-6.00	0.08-0.16	0.0-2.9	---	.24	.28			
Br: Bridgeport---	0-14	11	68	14-27	1.30-1.40	0.60-2.00	0.20-0.24	0.0-2.9	1.0-4.0	.32	.32	5	4L	86
	14-60	9	67	18-30	1.35-1.50	0.60-2.00	0.20-0.24	0.0-2.9	---	.43	.43			
Ca: Carwile-----	0-16	62	26	5-18	1.30-1.65	0.60-2.00	0.11-0.20	0.0-2.9	1.0-3.0	.24	.24	5	3	86
	16-21	35	33	25-39	1.45-1.75	0.20-2.00	0.12-0.20	3.0-5.9	---	.37	.37			
	21-45	25	27	35-60	1.35-1.75	0.06-0.20	0.12-0.20	6.0-8.9	---	.37	.37			
	45-60	35	33	20-45	1.35-1.75	0.20-2.00	0.12-0.20	6.0-8.9	---	.32	.32			

PHYSICAL PROPERTIES OF THE SOILS--Continued
McPherson County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Cb:														
Cass-----	0-7	68	20	7-17	1.40-1.60	2.00-6.00	0.16-0.18	0.0-2.9	1.0-2.0	.20	.20	4	3	86
	7-51	64	26	5-15	1.40-1.60	2.00-6.00	0.15-0.17	0.0-2.9	---	.20	.20			
	51-60	87	7	2-10	1.50-1.70	5.95-19.98	0.08-0.10	0.0-2.9	---	.20	.20			
Ce:														
Clime-----	0-9	7	48	40-50	1.35-1.45	0.06-0.20	0.12-0.14	3.0-5.9	1.0-4.0	.28	.28	3	4	86
	9-27	6	47	35-60	1.35-1.50	0.06-0.20	0.12-0.18	3.0-5.9	1.0-4.0	.28	.28			
	>27			---	---	---	---	---	---	---	---			
Cm:														
Clime-----	0-9	7	48	40-50	1.35-1.45	0.06-0.20	0.12-0.14	3.0-5.9	1.0-4.0	.28	.28	3	4	86
	9-27	6	47	35-60	1.35-1.50	0.06-0.20	0.12-0.18	3.0-5.9	1.0-4.0	.28	.28			
	>27			---	---	---	---	---	---	---	---			
Cr:														
Crete-----	0-5	24	52	20-27	1.20-1.40	0.60-2.00	0.22-0.24	3.0-5.9	2.0-4.0	.37	.37	5	6	48
	5-9	20	49	27-35	1.20-1.40	0.20-0.60	0.21-0.23	6.0-8.9	1.0-3.0	.37	.37			
	9-19	7	48	35-55	1.10-1.30	0.06-0.20	0.12-0.20	6.0-8.9	0.5-2.0	.37	.37			
	19-27	7	48	35-55	1.10-1.30	0.06-0.20	0.12-0.20	6.0-8.9	0.5-2.0	.37	.37			
	27-38	7	48	35-55	1.10-1.30	0.06-0.20	0.12-0.20	6.0-8.9	0.5-2.0	.37	.37			
	38-48	20	48	25-40	1.20-1.40	0.20-2.00	0.18-0.22	6.0-8.9	0.5-1.0	.43	.43			
	48-80	20	48	25-40	1.20-1.40	0.20-2.00	0.18-0.22	6.0-8.9	0.5-1.0	.43	.43			
Cs:														
Crete-----	0-6	24	52	20-27	1.20-1.40	0.60-2.00	0.22-0.24	3.0-5.9	2.0-4.0	.37	.37	5	6	48
	6-13	20	49	27-35	1.20-1.40	0.20-0.60	0.21-0.23	6.0-8.9	---	.37	.37			
	13-43	7	48	35-55	1.10-1.30	0.00-0.60	0.12-0.20	6.0-8.9	---	.37	.37			
	43-60	20	48	25-40	1.20-1.40	0.20-2.00	0.18-0.22	6.0-8.9	---	.37	.37			
Ct:														
Crete-----	0-6	20	49	27-35	1.20-1.40	0.20-0.60	0.21-0.23	6.0-8.9	2.0-4.0	.37	.37	5	7	38
	6-13	20	49	27-35	1.20-1.40	0.20-0.60	0.21-0.23	6.0-8.9	---	.37	.37			
	13-35	7	48	35-55	1.10-1.30	0.00-0.60	0.12-0.20	6.0-8.9	---	.37	.37			
	35-60	20	48	25-40	1.20-1.40	0.20-2.00	0.18-0.22	6.0-8.9	---	.37	.37			
De:														
Detroit-----	0-16	20	48	28-35	1.25-1.40	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	5	7	38
	16-37	7	53	35-45	1.35-1.50	0.06-0.20	0.12-0.18	6.0-8.9	1.0-3.0	.37	.37			
	37-60	24	50	18-35	1.30-1.50	0.20-0.60	0.18-0.22	3.0-5.9	0.5-1.0	.37	.37			
Dr:														
Drummond-----	0-19	38	36	20-30	1.35-1.55	0.60-2.00	0.11-0.18	0.0-2.9	0.5-1.0	.49	.49	2	4L	48
	19-60	25	27	35-60	1.40-1.65	0.00-0.06	0.09-0.17	6.0-8.9	---	.55	.55			
Ed:														
Edalgo-----	0-6	26	53	15-27	1.30-1.40	0.60-2.00	0.22-0.24	0.0-2.9	2.0-4.0	.37	.37	3	6	48
	6-15	7	52	28-55	1.35-1.50	0.00-0.60	0.13-0.23	3.0-5.9	1.0-3.0	.37	.37			
	15-30	6	44	35-65	1.40-1.60	0.00-0.06	0.10-0.18	6.0-8.9	1.0-2.0	.37	.37			
	>30			---	---	---	---	---	---	---	---			
Ee:														
Edalgo-----	0-8	20	48	28-37	1.30-1.40	0.00-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	3	7	38
	8-23	6	44	35-65	1.40-1.60	0.00-0.06	0.10-0.18	6.0-8.9	1.0-2.0	.37	.37			
	>23			---	---	---	---	---	---	---	---			
Fa:														
Farnum-----	0-12	42	38	14-27	1.35-1.45	0.60-2.00	0.19-0.22	0.0-2.9	1.0-3.0	.28	.28	5	6	48
	12-49	56	14	25-35	1.40-1.50	0.60-2.00	0.15-0.19	3.0-5.9	---	.28	.28			
	49-60	61	19	12-29	1.40-1.55	0.60-2.00	0.13-0.16	0.0-2.9	---	.28	.28			
Ge:														
Geary-----	0-7	11	68	15-27	1.30-1.40	0.60-2.00	0.22-0.24	0.0-2.9	1.0-4.0	.32	.32	5	6	48
	7-32	7	62	27-35	1.35-1.50	0.60-2.00	0.17-0.20	3.0-5.9	---	.43	.43			
	32-60	7	67	20-32	1.30-1.40	0.60-2.00	0.15-0.19	3.0-5.9	---	.43	.43			
Go:														
Goessel-----	0-14	6	47	40-55	1.30-1.40	0.00-0.06	0.12-0.16	6.0-8.9	1.0-4.0	.28	.28	5	4	86
	14-50	6	47	40-55	1.35-1.45	0.00-0.06	0.10-0.15	6.0-8.9	---	.28	.28			
	50-60	8	52	30-50	1.40-1.55	0.00-0.06	0.09-0.14	6.0-8.9	---	.28	.28			
Ho:														
Hord-----	0-6	11	67	17-27	1.30-1.40	0.60-2.00	0.20-0.24	0.0-2.9	2.0-4.0	.32	.32	5	6	48
	6-43	7	65	20-35	1.35-1.45	0.60-2.00	0.17-0.22	0.0-2.9	---	.32	.32			
	43-60	9	67	18-30	1.30-1.50	0.60-2.00	0.17-0.22	0.0-2.9	---	.43	.43			
INT:														
Aquolls-----	0-72			---	---	---	---	---	---	---	---	-	---	0
Ir:														
Irwin, bedrock substratum--	0-11	20	48	28-35	1.35-1.45	0.20-0.60	0.18-0.23	3.0-5.9	2.0-4.0	.37	.37	4	7	38
	11-42 >42	23	29	40-55	1.40-1.50	0.00-0.06	0.10-0.15	6.0-8.9	---	.37	.37			
La:														
Ladysmith----	0-8	20	48	28-35	1.35-1.45	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	5	7	38
	8-48	22	28	40-60	1.35-1.50	0.00-0.06	0.10-0.15	6.0-8.9	1.0-3.0	.37	.37			
	48-60	7	48	35-55	1.40-1.60	0.00-0.60	0.10-0.19	3.0-5.9	0.0-1.0	.37	.37			
Ld:														
Ladysmith----	0-8	20	48	28-35	1.35-1.45	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	5	7	38
	8-48	22	28	40-60	1.35-1.50	0.00-0.06	0.10-0.15	6.0-8.9	1.0-3.0	.37	.37			
	48-60	7	48	35-55	1.40-1.60	0.00-0.60	0.10-0.19	3.0-5.9	0.0-1.0	.37	.37			
Drummond-----	0-7	38	36	20-30	1.35-1.55	0.60-2.00	0.11-0.18	0.0-2.9	0.5-1.0	.49	.49	2	6	48
	7-60	25	27	35-60	1.40-1.65	0.00-0.06	0.09-0.17	6.0-8.9	---	.55	.55			
Le:														
Lancaster----	0-10	43	38	12-26	1.35-1.45	0.60-2.00	0.17-0.22	0.0-2.9	1.0-4.0	.28	.28	3	6	48
	10-32 >32	35	38	18-35	1.35-1.50	0.60-2.00	0.15-0.19	3.0-5.9	---	.28	.32			

PHYSICAL PROPERTIES OF THE SOILS--Continued
McPherson County, Kansas: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										K	Kf	T		
Lh:	In	Pct	Pct	Pct	g/cc	in/hr	In/in	Pct	Pct					
Lancaster----	0-10	43	38	12-26	1.35-1.45	0.60-2.00	0.17-0.22	0.0-2.9	1.0-4.0	.28	.28	3	6	48
	10-32	35	38	18-35	1.35-1.50	0.60-2.00	0.15-0.19	3.0-5.9	----	.28	.32			
	>32			----	----	----	----	----	----	----	----			
Hedville-----	0-15	44	41	8-22	1.35-1.50	0.60-2.00	0.18-0.20	0.0-2.9	1.0-4.0	.32	.32	2	5	56
	>15			----	----	----	----	----	----	----	----			
LHH:														
Lancaster----	0-9	43	38	12-26	1.35-1.45	0.60-2.00	0.17-0.22	0.0-2.9	1.0-4.0	.28	.43	3	6	48
	9-24	35	38	18-35	1.35-1.50	0.60-2.00	0.15-0.19	3.0-5.9	----	.28	.32			
	24-30	61	18	12-30	1.40-1.55	0.60-2.00	0.15-0.19	0.0-2.9	----	.28	.32			
	>30			----	----	----	----	----	----	----	----			
Hedville-----	0-8	44	41	8-22	1.35-1.50	0.60-2.00	0.18-0.20	0.0-2.9	1.0-4.0	.32	.43	1	5	56
	8-17	44	41	8-22	1.35-1.50	0.60-2.00	0.08-0.18	0.0-2.9	----	.32	.55			
	>17			----	----	----	----	----	----	----	----			
Ln:														
Longford-----	0-15	20	49	27-35	1.30-1.40	0.20-0.60	0.21-0.23	3.0-5.9	1.0-4.0	.32	.32	5	7	38
	15-29	10	65	15-35	1.30-1.40	0.60-2.00	0.18-0.22	3.0-5.9	1.0-3.0	.32	.32			
	29-47	20	53	20-35	1.30-1.40	0.20-0.60	0.15-0.20	3.0-5.9	0.5-2.0	.32	.32			
	47-60	7	53	35-45	1.35-1.50	0.06-0.20	0.14-0.20	6.0-8.9	0.5-2.0	.32	.32			
Lo:														
Longford-----	0-6	20	49	27-35	1.30-1.40	0.20-0.60	0.21-0.23	3.0-5.9	1.0-2.0	.32	.32	5	7	38
	6-15	20	49	27-35	1.30-1.40	0.20-0.60	0.21-0.23	3.0-5.9	1.0-2.0	.32	.32			
	15-47	8	52	35-45	1.35-1.50	0.00-0.60	0.11-0.20	6.0-8.9	0.5-1.0	.43	.43			
	47-60	20	49	27-35	1.30-1.40	0.20-0.60	0.14-0.20	3.0-5.9	0.5-0.5	.32	.32			
Mc:														
Mccook-----	0-14	66	20	10-17	1.40-1.60	2.00-6.00	0.16-0.18	0.0-2.9	1.0-2.0	.20	.20	5	3	86
	14-60	60	26	10-18	1.30-1.45	0.60-2.00	0.17-0.20	0.0-2.9	----	.43	.43			
Ns:														
Ness-----	0-60	5	45	40-60	1.30-1.45	0.00-0.06	0.11-0.14	6.0-8.9	1.0-3.0	.28	.28	5	4	86
Nw:														
New Cambria--	0-12	5	45	40-60	1.30-1.40	0.06-0.20	0.12-0.14	6.0-8.9	2.0-4.0	.28	.28	5	4	86
	12-34	5	46	38-60	1.35-1.45	0.06-0.20	0.13-0.18	6.0-8.9	----	.28	.28			
	34-60	8	52	30-50	1.35-1.45	0.00-0.60	0.12-0.16	6.0-8.9	----	.28	.28			
Pa:														
Plevna-----	0-18	67	20	8-18	1.40-1.50	2.00-6.00	0.14-0.16	0.0-2.9	1.0-4.0	.20	.20	5	3	86
	18-36	67	20	8-18	1.40-1.50	2.00-6.00	0.12-0.16	0.0-2.9	----	.20	.20			
	36-60	79	17	1-7	1.50-1.60	2.00-6.00	0.05-0.07	0.0-2.9	----	.20	.20			
Pr:														
Pratt-----	0-11	79	16	2-8	1.40-1.55	5.95-19.98	0.10-0.13	0.0-2.9	0.5-1.0	.17	.17	5	2	134
	11-36	86	7	4-11	1.45-1.55	5.95-19.98	0.09-0.12	0.0-2.9	----	.17	.17			
	36-60	79	16	1-8	1.45-1.60	5.95-19.98	0.08-0.12	0.0-2.9	----	.17	.17			
Ro:														
Roxbury-----	0-21	7	62	27-35	1.30-1.45	0.60-2.00	0.21-0.23	3.0-5.9	2.0-4.0	.43	.43	5	4L	86
	21-50	7	66	18-35	1.35-1.50	0.60-2.00	0.17-0.22	3.0-5.9	1.0-3.0	.43	.43			
	50-60	9	64	18-35	1.35-1.50	0.60-2.00	0.17-0.22	3.0-5.9	0.5-0.5	.43	.43			
Sm:														
Smolan-----	0-11	20	49	27-35	1.30-1.40	0.20-0.60	0.21-0.23	3.0-5.9	2.0-4.0	.37	.37	5	7	38
	11-16	24	50	18-35	1.30-1.40	0.14-0.14	0.21-0.24	3.0-5.9	----	.37	.37			
	16-60	7	51	35-50	1.30-1.45	0.06-0.20	0.12-0.18	6.0-8.9	----	.37	.37			
To:														
Tobin-----	0-20	10	68	18-27	1.30-1.40	0.60-2.00	0.20-0.24	0.0-2.9	1.0-4.0	.32	.32	5	6	48
	20-32	9	64	18-35	1.35-1.50	0.60-2.00	0.17-0.20	3.0-5.9	1.0-4.0	.32	.32			
	32-60	9	64	18-35	1.35-1.45	0.60-2.00	0.18-0.22	3.0-5.9	0.5-0.5	.43	.43			
W:														
Water-----	----			----	----	----	----	----	----	----	----	----	----	----
Wb:														
Wells-----	0-15	40-60	15-30	18-27	1.35-1.50	0.60-2.00	0.20-0.22	0.0-2.9	1.0-4.0	.28	.28	5	6	48
	15-36	40-60	15-25	27-35	1.35-1.50	0.60-2.00	0.15-0.19	3.0-5.9	1.0-3.0	.32	.32			
	36-60	40-70	15-25	10-30	1.35-1.60	0.60-2.00	0.12-0.18	0.0-2.9	0.5-2.0	.32	.32			
Wc:														
Wells-----	0-11	40	38	18-27	1.35-1.50	0.60-2.00	0.20-0.22	0.0-2.9	1.0-4.0	.28	.28	5	6	48
	11-15	55	14	27-35	1.35-1.60	0.60-2.00	0.15-0.19	3.0-5.9	----	.28	.28			
	15-49	55	14	27-35	1.35-1.50	0.60-2.00	0.15-0.19	3.0-5.9	----	.32	.32			
	49-60	65	15	10-30	1.35-1.60	0.60-2.00	0.12-0.18	0.0-2.9	----	.32	.32			

CHEMICAL PROPERTIES OF THE SOILS
McPherson County, Kansas

The Chemical Properties table shows estimates of some characteristics and features that affect soil behavior. These estimates are given for the major layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils. Depth to the upper and lower boundaries of each layer is indicated.

Cation-exchange capacity is the total amount of extractable bases that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. Soils having a high cation-exchange capacity can retain cations. The ability to retain cations helps to prevent the pollution of ground water.

Effective cation-exchange capacity refers to the sum of extractable bases plus aluminum expressed in terms of milliequivalents per 100 grams of soil. It is determined for soils that have pH of less than 5.5.

Soil reaction is a measure of acidity or alkalinity and is expressed as a range in pH values. The range in pH of each major horizon is based on many field tests. For many soils, values have been verified by laboratory analyses. Soil reaction is important in selecting crops and other plants, in evaluating soil amendments for fertility and stabilization, and in determining the risk of corrosion.

Calcium carbonate equivalent is the percent of carbonates, by weight, in the fraction of the soil less than 2 millimeters in size. The availability of plant nutrients is influenced by the amount of carbonates in the soil. Incorporating nitrogen fertilizer into calcareous soils helps to prevent nitrite accumulation and ammonium-N volatilization.

Gypsum is expressed as a percent, by weight, of hydrated calcium sulfates in the fraction of the soil less than 20 millimeters in size. Gypsum is partially soluble in water and can be dissolved and removed by water. Soils that have a high content of gypsum may collapse if the gypsum is removed by percolating water.

Salinity is a measure of soluble salts in the soil at saturation. It is expressed as the electrical conductivity of the saturation extract, in millimhos per centimeter at 25 degrees C. Estimates are based on field and laboratory measurements at representative sites of nonirrigated soils. The salinity of irrigated soils is affected by the quality of the irrigation water and by the frequency of water application. Hence, the salinity of soils in individual fields can differ greatly from the value given in the table. Salinity affects the suitability of a soil for crop production, the stability of soil if used as construction material, and the potential of the soil to corrode metal and concrete.

Sodium adsorption ratio (SAR) is a measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste. It is the ratio of the Na concentration divided by the square root of one-half of the Ca + Mg concentration. Soils that have SAR values of 13 or more may be characterized by an increased dispersion of organic matter and clay particles, reduced permeability and aeration, and a general degradation of soil structure.

CHEMICAL PROPERTIES OF THE SOILS--Continued
McPherson County, Kansas

PAGE 2 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Effective Cation Exchange Capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm		
053ED: Edalgo-----	0-8 8-34 >34	8.0-24 14-40 ---	--- --- 0.0-0.0	5.6-6.0 5.6-8.4 ---	--- --- ---	--- --- ---	--- --- ---	--- --- ---
053MB: Mccook-----	0-12 12-60	6.0-15 4.0-11	--- ---	7.4-8.4 7.4-8.4	--- ---	0 0	0 0	0 0
053MC: Mccook-----	0-12 12-60	6.0-15 4.0-11	--- ---	7.4-8.4 7.4-8.4	--- ---	0 0	0 0	0 0
079GE: Geary-----	0-9 9-35 35-60	6.0-19 10-21 8.0-19	--- --- ---	5.6-6.5 5.6-7.8 6.1-8.4	0 0 0	0 0 0	0 0 0	0 0 0
079LB: Ladysmith-----	0-10 10-45 45-60	12-24 16-36 14-33	--- --- ---	5.6-7.3 5.6-7.8 7.4-8.4	--- 0-5 0-5	--- --- ---	--- --- ---	--- --- ---
115IC: Irwin-----	0-13 13-40 40-60	12-24 16-38 14-34	--- --- ---	5.6-7.3 5.6-8.4 6.6-8.4	0 0 0	0 0 0	0 0 0	0 0 0
159DP: Dillwyn-----	0-9 9-60	0.0-6.0 0.0-5.0	--- ---	5.6-7.3 5.6-7.8	--- ---	--- ---	--- ---	--- ---
Plevna-----	0-10 10-33 33-60	1.0-7.0 3.0-11 0.0-4.0	--- --- ---	6.6-8.4 6.6-8.4 6.6-8.4	0 0 0	0 0 0	0 0 0	0 0 0
159GC: Geary-----	0-7 7-32 32-60	6.0-19 10-21 8.0-19	--- --- ---	5.6-6.5 5.6-7.8 6.1-8.4	0 0 0	0 0 0	0 0 0	0 0 0
Clark-----	0-11 11-60	11-21 7.0-21	--- ---	7.4-8.4 7.4-8.4	0-5 15-45	--- ---	--- ---	--- ---
159LA: Lancaster-----	0-7 7-25 25-35 >35	5.0-18 7.0-21 4.0-18 ---	--- --- --- 0.0-0.0	5.6-6.5 5.6-7.3 6.1-7.3 ---	0 0 0 ---	0 0 0 ---	0 0 0 ---	0 0 0 ---
159NF: Naron-----	0-14 14-40 40-60	3.0-10 7.0-16 0.0-9.0	--- --- ---	5.6-7.3 5.6-7.8 6.1-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
159PF: Pratt-----	0-13 13-38 38-60	1.0-5.0 1.0-7.0 0.0-5.0	--- --- ---	5.6-7.3 5.6-7.3 6.1-7.3	--- --- ---	--- --- ---	--- --- ---	--- --- ---
159PR: Pratt-----	0-13 13-38 38-60	1.0-5.0 1.0-7.0 0.0-5.0	--- --- ---	5.6-7.3 5.6-7.3 6.1-7.3	--- --- ---	--- --- ---	--- --- ---	--- --- ---
Carwile-----	0-12 12-17 17-60	2.0-13 10-24 14-36	--- --- ---	5.1-7.3 5.1-7.3 6.1-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
159SO: Smolan-----	0-6 6-60	11-24 14-30	--- ---	5.6-7.3 5.6-7.8	--- ---	--- ---	--- ---	--- ---
159WA: Waldeck-----	0-25 25-42 42-60	3.0-11 3.0-10 0.0-3.0	--- --- ---	7.4-8.4 7.4-8.4 7.4-8.4	--- --- ---	--- --- ---	--- --- ---	--- --- ---
169CE: Clime-----	0-8 8-20 20-28 >28	13-27 14-39 14-32 ---	--- --- --- 0.0-0.0	6.6-8.4 7.4-8.4 7.4-8.4 ---	5-10 5-10 10-15 ---	--- --- --- ---	--- --- --- ---	--- --- --- ---
169ED: Edalgo-----	0-9 9-13 13-28 >28	12-25 11-35 14-40 ---	--- --- --- 0.0-0.0	5.6-6.0 6.1-6.5 5.6-8.4 ---	--- --- --- ---	--- --- --- ---	--- --- --- ---	--- --- --- ---
169RO: Roxbury-----	0-26 26-60	8.0-19 7.0-21	--- ---	7.4-8.4 7.4-8.4	1-5 5-10	--- ---	--- ---	--- ---

CHEMICAL PROPERTIES OF THE SOILS--Continued
McPherson County, Kansas

PAGE 3 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Effective Cation Exchange Capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm		
1191: Blazefork-----	0-3	15-32	---	4.5-6.5	0	0	0	0
	3-7	15-32	---	4.5-6.5	0	0	0	0
	7-14	19-40	---	6.1-8.4	0	0	0	0
	14-22	19-40	---	6.1-8.4	0	0	0	0
	22-29	19-40	---	6.1-8.4	0	0	0	0
	29-34	19-40	---	6.1-8.4	0	0	0	0
	34-40	14-30	---	7.4-8.4	0	0	0	0
	40-48	14-26	---	7.4-8.4	0	0	0	0
	48-61	14-26	---	7.4-8.4	0	0	0	0
	61-80	14-26	---	7.4-8.4	0	0	0	0
1200: Buhler-----	0-3	20-35	---	5.6-7.3	0	0-3	0.0-6.0	0-7
	3-8	20-35	---	5.6-7.3	0	0-3	0.0-4.0	20-30
	8-12	10-15	---	6.1-7.8	0	1-5	0.0-4.0	20-30
	12-16	10-15	---	6.1-7.8	0	1-5	0.0-4.0	20-30
	16-24	12-30	---	6.6-7.8	1-5	0-5	4.0-8.0	20-30
	24-36	12-30	---	6.6-7.8	1-5	0-5	4.0-8.0	15-25
	36-42	12-30	---	6.6-7.8	1-5	0-5	4.0-8.0	10-20
	42-50	15-30	---	7.4-9.0	1-5	0-5	0.0-8.0	10-20
	50-58	15-25	---	7.4-9.0	1-5	0-5	0.0-8.0	10-20
	58-76	8.0-15	---	7.4-8.4	0	0	0.0-4.0	0-15
	76-80	8.0-15	---	7.4-8.4	0	0	0.0-4.0	0-15
Blazefork-----	0-3	15-32	---	4.5-6.5	0	0	0	0
	3-7	15-32	---	4.5-6.5	0	0	0	0
	7-14	19-40	---	6.1-8.4	0	0	0	0
	14-22	19-40	---	6.1-8.4	0	0	0	0
	22-29	19-40	---	6.1-8.4	0	0	0	0
	29-34	19-40	---	6.1-8.4	0	0	0	0
	34-40	14-30	---	7.4-8.4	0	0	0	0
	40-48	14-26	---	7.4-8.4	0	0	0	0
	48-61	14-26	---	7.4-8.4	0	0	0	0
	61-80	14-26	---	7.4-8.4	0	0	0	0
1324: Carway-----	0-7	7.0-12	---	5.6-6.5	0	0	0	0
	7-10	12-18	---	6.1-7.3	0	0	0	0
	10-15	12-18	---	6.1-7.3	0	0	0	0
	15-22	12-18	---	6.1-7.3	0	0	0	0
	22-35	12-18	---	6.1-7.3	0	0	0	0
	35-40	24-35	---	6.6-7.8	0-5	0	0	0
	40-54	24-35	---	6.6-7.8	0-5	0	0	0
	54-63	24-35	---	6.6-7.8	0-5	0	0	0
	63-72	24-35	---	6.6-7.8	0-5	0	0	0
	72-80	9.0-16	---	6.6-7.8	0-5	0	0	0
Carbika-----	0-11	7.0-12	---	6.1-7.3	0	0	0	0
	11-15	28-38	---	6.1-8.4	0-5	0	0	0
	15-22	28-38	---	6.1-8.4	0-5	0	0	0
	22-34	12-16	---	6.1-8.4	0-5	0	0	0
	34-41	12-16	---	6.1-8.4	0-5	0	0	0
	41-60	12-16	---	6.1-8.4	0-5	0	0	0
	60-80	12-16	---	7.4-8.4	0-5	0	0	0
1553: Darlow-----	0-5	5.0-15	---	4.5-7.8	0	0	0.0-2.0	0-5
	5-8	5.0-15	---	4.5-7.8	0	0	0.0-2.0	0-5
	8-14	15-25	---	6.6-9.0	0-2	0	2.0-8.0	10-40
	14-20	15-25	---	6.6-9.0	0-2	0	2.0-8.0	15-40
	20-26	15-25	---	6.6-9.0	0-2	0	2.0-8.0	15-40
	26-33	15-30	---	7.9-9.0	0-1	0	4.0-16.0	30-40
	33-44	15-30	---	7.9-9.0	0-1	0	4.0-16.0	30-40
	44-53	10-20	---	7.9-8.4	0-1	0	0.0-4.0	25-35
	53-68	10-15	---	7.9-8.4	0-1	0	0.0-4.0	25-35
	68-80	0.0-10	---	7.4-8.4	0-1	0	0.0-2.0	10-30
Elmer-----	0-6	5.0-12	---	4.5-7.3	0	0	0.0-2.0	0-1
	6-9	5.0-12	---	4.5-7.3	0	0	0.0-2.0	0-1
	9-19	5.0-12	---	4.5-7.3	0	0	0.0-2.0	0-1
	19-26	10-18	---	7.4-9.0	0	0	0.0-2.0	7-20
	26-37	10-18	---	7.4-9.0	0	0	1.0-2.0	7-20
	37-43	15-20	---	7.9-9.0	1-2	0	1.0-4.0	20-30
	43-51	15-20	---	7.9-9.0	1-2	0	1.0-4.0	20-30
	51-61	8.0-15	---	7.4-9.0	0-1	0	1.0-2.0	5-25
	61-72	8.0-15	---	7.4-9.0	0-1	0	1.0-2.0	5-20
	72-80	8.0-15	---	7.4-9.0	0-1	0	0.0-2.0	5-20

CHEMICAL PROPERTIES OF THE SOILS--Continued
McPherson County, Kansas

PAGE 4 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Effective Cation Exchange Capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm		
1555: Dillhut-----	0-4	1.0-3.0	---	5.6-6.5	0	0	0	0
	4-9	1.0-3.0	---	5.6-6.5	0	0	0	0
	9-18	0.0-2.0	---	5.6-6.5	0	0	0	0
	18-26	0.0-2.0	---	5.6-6.5	0	0	0	0
	26-41	10-18	---	6.6-7.3	0	0	0	0
	41-55	7.0-10	---	6.6-7.3	0	0	0	0
	55-65	7.0-10	---	6.6-7.3	0	0	0	0
	65-70	7.0-10	---	6.6-7.3	0	0	0	0
	70-80	3.0-9.0	---	6.1-7.3	0	0	0	0
Plev-----	0-4	3.0-6.0	---	5.1-6.5	0	0	0	0
	4-12	0.0-3.0	---	5.1-6.5	0	0	0	0
	12-35	0.0-1.0	---	5.6-6.5	0	0	0	0
	35-46	0.0-1.0	---	5.6-6.5	0	0	0	0
	46-57	7.0-15	---	6.1-7.3	0	0	0	0
	57-75	7.0-15	---	6.1-7.3	0	0	0	0
	75-80	3.0-9.0	---	6.1-7.3	0	0	0	0
1985: Hayes-----	0-8	5.0-8.0	---	5.1-7.3	0	0	0	0
	8-14	6.0-10	---	6.1-7.3	0	0	0	0
	14-23	6.0-10	---	6.1-7.3	0	0	0	0
	23-34	6.0-10	---	6.1-7.3	0	0	0	0
	34-42	6.0-10	---	6.1-7.3	0	0	0	0
	42-47	6.0-10	---	6.1-7.3	0	0	0	0
	47-56	10-15	---	6.6-7.8	0	0	0	0
	56-69	24-35	---	6.6-7.8	0-5	0	0	0
	69-80	24-35	---	6.6-7.8	0-5	0	0	0
2391: Kaskan-----	0-9	20-30	---	6.1-7.3	0	0	0	0
	9-13	20-30	---	6.1-7.3	0	0	0	0
	13-17	6.0-10	---	6.1-7.8	0-1	0	0	0
	17-21	6.0-10	---	6.1-7.8	0-1	0	0	0
	21-27	6.0-10	---	6.1-7.8	0-1	0	0	0
	27-43	0.0-3.0	---	6.1-7.8	0	0	0	0
	43-57	0.0-8.0	---	6.1-7.8	0-1	0	0	0
	57-80	0.0-8.0	---	6.1-7.8	0-1	0	0	0
2588: Longford, Moderately Eroded-----	0-6	15-25	---	5.6-7.3	0	0	0	0
	6-11	15-25	---	6.1-7.3	0	0	0	0
	11-28	15-30	---	6.1-7.3	0	0	0	0
	28-43	15-30	---	6.1-7.3	0	0	0	0
	43-60	10-20	---	6.1-7.8	0	0	0	0
	60-80	10-20	---	6.1-7.8	0	0	0	0
3181: Pratt-----	0-8	0.0-3.0	0.0-0.0	5.6-7.3	0	0	0	0
	8-24	2.0-5.0	0.0-0.0	5.6-7.3	0	0	0	0
	24-64	3.0-7.0	0.0-0.0	5.6-7.3	0	0	0	0
	64-80	1.0-3.0	0.0-0.0	6.1-7.3	0	0	0	0
Turon-----	0-8	1.0-3.0	---	5.1-7.3	0	0	0	0
	8-28	2.0-5.0	---	5.1-7.3	0	0	0	0
	28-40	3.0-7.0	---	5.1-7.3	0	0	0	0
	40-58	24-33	---	6.6-7.8	0	0	0	0
	58-75	24-33	---	6.6-7.8	0	0	0	0
	75-80	24-33	---	6.6-7.8	0	0	0	0
3641: Tivin-----	0-7	0.0-1.0	---	5.6-6.5	0	0	0.0-1.0	0
	7-18	0.0-1.0	---	6.1-7.3	0	0	0.0-1.0	0
	18-80	0.0-1.0	---	6.1-7.3	0	0	0.0-1.0	0
Dillhut-----	0-4	1.0-3.0	---	5.6-6.5	0	0	0	0
	4-9	1.0-3.0	---	5.6-6.5	0	0	0	0
	9-18	0.0-2.0	---	5.6-6.5	0	0	0	0
	18-26	0.0-2.0	---	5.6-6.5	0	0	0	0
	26-41	10-18	---	6.6-7.3	0	0	0	0
	41-55	7.0-10	---	6.6-7.3	0	0	0	0
	55-65	7.0-10	---	6.6-7.3	0	0	0	0
	65-70	7.0-10	---	6.6-7.3	0	0	0	0
	70-80	3.0-9.0	---	6.1-7.3	0	0	0	0
AED: Arents, Earthen Dam-----	---	---	---	---	---	---	---	---
At: Attica-----	0-10	1.0-7.0	---	5.6-7.3	---	---	---	---
	10-25	3.0-11	---	5.6-6.5	---	---	---	---
	25-60	1.0-11	---	6.1-7.8	---	---	---	---
Br: Bridgeport-----	0-14	6.0-19	---	6.6-8.4	---	0	0	0
	14-60	7.0-18	---	7.4-8.4	---	0	0	0
Ca: Carwile-----	0-16	2.0-13	---	5.1-7.3	---	---	---	---
	16-21	10-24	---	5.1-7.3	---	---	---	---
	21-45	14-36	---	6.1-8.4	---	---	---	---
	45-60	8.0-27	---	6.6-8.4	---	---	---	---

CHEMICAL PROPERTIES OF THE SOILS--Continued
McPherson County, Kansas

PAGE 5 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Effective Cation Exchange Capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm		
Cb:								
Cass-----	0-7	3.0-11	---	5.6-7.3	0	0	0	0
	7-51	2.0-9.0	---	6.1-8.4	0	0	0	0
	51-60	0.0-6.0	---	6.1-8.4	0	0	0	0
Ce:								
Cline-----	0-9	16-33	---	6.6-8.4	5-10	---	---	---
	9-27	14-39	---	7.4-8.4	5-10	---	---	---
	>27	---	0.0-0.0	---	---	---	---	---
Cm:								
Cline-----	0-9	16-33	---	6.6-8.4	5-10	---	---	---
	9-27	14-39	---	7.4-8.4	5-10	---	---	---
	>27	---	0.0-0.0	---	---	---	---	---
Cr:								
Crete-----	0-5	16-23	---	5.6-6.0	0	0	0	0
	5-9	20-28	---	5.6-6.0	0	0	0	0
	9-19	25-41	---	6.1-7.3	0	0	0	0
	19-27	25-41	---	6.1-7.3	0	0	0	0
	27-38	25-41	---	6.1-7.3	0	0	0	0
	38-48	18-29	---	7.4-8.4	0-5	0	0	0
	48-80	18-29	---	7.4-8.4	0-5	0	0	0
Cs:								
Crete-----	0-6	8.0-19	---	5.6-6.0	0	0	0	0
	6-13	10-21	---	5.6-6.0	0	0	0	0
	13-43	14-33	---	6.1-7.3	0	0	0	0
	43-60	10-24	---	7.4-8.4	0-5	0	0	0
Ct:								
Crete-----	0-6	11-24	---	5.6-6.0	0	0	0	0
	6-13	10-21	---	5.6-6.0	0	0	0	0
	13-35	14-33	---	6.1-7.3	0	0	0	0
	35-60	10-24	---	7.4-8.4	0-5	0	0	0
De:								
Detroit-----	0-16	12-24	---	6.1-7.3	0	0	0	0
	16-37	14-27	---	6.6-7.8	0	0	0	0
	37-60	7.0-21	---	6.6-8.4	---	0	0	---
Dr:								
Drummond-----	0-19	8.0-19	---	6.1-8.4	---	---	0.0-4.0	---
	19-60	14-36	---	7.4-9.0	---	---	2.0-8.0	---
Ed:								
Edalgo-----	0-6	6.0-19	---	5.6-6.0	---	---	---	---
	6-15	11-35	---	6.1-6.5	---	---	---	---
	15-30	14-40	---	5.6-8.4	---	---	---	---
	>30	---	0.0-0.0	---	---	---	---	---
Ee:								
Edalgo-----	0-8	12-25	---	5.6-6.0	---	---	---	---
	8-23	14-40	---	5.6-8.4	---	---	---	---
	>23	---	0.0-0.0	---	---	---	---	---
Fa:								
Farnum-----	0-12	6.0-18	---	5.6-7.3	---	---	---	---
	12-49	10-21	---	6.1-8.4	---	---	---	---
	49-60	4.0-18	---	6.6-8.4	---	---	---	---
Ge:								
Geary-----	0-7	6.0-19	---	5.6-6.5	0	0	0	0
	7-32	10-21	---	5.6-7.8	0	0	0	0
	32-60	8.0-19	---	6.1-8.4	0	0	0	0
Go:								
Goessel-----	0-14	16-36	---	6.1-7.3	---	---	---	---
	14-50	16-33	---	7.4-8.4	---	---	---	---
	50-60	12-30	---	7.9-8.4	---	---	---	---
Ho:								
Hord-----	0-6	7.0-19	---	5.6-7.3	0	0	0	0
	6-43	8.0-21	---	6.1-7.8	0	0	0	0
	43-60	7.0-18	---	7.4-8.4	0-5	0	0	0
INT:								
Aquolls-----	0-72	---	---	---	---	---	---	---
Ir:								
Irwin, bedrock substratum-----	0-11	12-24	---	5.6-7.3	---	---	---	---
	11-42	16-33	---	5.6-8.4	---	---	---	---
	>42	---	0.0-0.0	---	---	---	---	---
La:								
Ladysmith-----	0-8	12-24	---	5.6-7.3	---	---	---	---
	8-48	16-36	---	5.6-7.8	---	---	---	---
	48-60	14-33	---	7.4-8.4	---	---	---	---
Ld:								
Ladysmith-----	0-8	12-24	---	5.6-7.3	---	---	---	---
	8-48	16-36	---	5.6-7.8	---	---	---	---
	48-60	14-33	---	7.4-8.4	---	---	---	---
Drummond-----	0-7	8.0-19	---	6.1-8.4	---	---	0.0-4.0	---
	7-60	14-36	---	7.4-9.0	---	---	2.0-8.0	---
Le:								
Lancaster-----	0-10	5.0-18	---	5.6-6.5	0	0	0	0
	10-32	7.0-21	---	5.6-7.3	0	0	0	0
	>32	---	0.0-0.0	---	---	---	---	---

CHEMICAL PROPERTIES OF THE SOILS--Continued
McPherson County, Kansas

PAGE 6 of 6

Map symbol and soil name	Depth	Cation- exchange capacity	Effective Cation Exchange Capacity	Soil reaction	Calcium carbonate	Gypsum	Salinity	Sodium adsorption ratio
	In	meq/100g	pH	Pct	Pct	mmhos/cm		
Lh:								
Lancaster-----	0-10	5.0-18	---	5.6-6.5	0	0	0	0
	10-32	7.0-21	---	5.6-7.3	0	0	0	0
	>32	---	0.0-0.0	---	---	---	---	---
Hedville-----	0-15	3.0-16	---	5.6-7.3	0	0	0	0
	>15	---	0.0-0.0	---	---	---	---	---
LHH:								
Lancaster-----	0-9	5.0-18	---	5.6-6.5	0	0	0	0
	9-24	7.0-21	---	5.6-7.3	0	0	0	0
	24-30	4.0-18	---	6.1-7.3	0	0	0	0
	>30	---	0.0-0.0	---	---	---	---	---
Hedville-----	0-8	3.0-16	---	5.6-7.3	0	0	0	0
	8-17	3.0-13	---	5.6-7.3	0	0	0	0
	>17	---	0.0-0.0	---	---	---	---	---
Ln:								
Longford-----	0-15	11-24	---	5.6-7.3	0	0	0	0
	15-29	6.0-23	---	6.1-7.3	0	0	0	0
	29-47	8.0-22	---	6.1-7.8	---	0	0	0
	47-60	14-28	---	5.1-7.3	0	0	0	0
Lo:								
Longford-----	0-6	11-22	---	5.6-7.3	---	---	---	---
	6-15	11-22	---	6.1-7.3	---	---	---	---
	15-47	14-28	---	6.1-7.3	---	---	---	---
	47-60	11-21	---	6.1-7.8	---	---	---	---
Mc:								
Mccook-----	0-14	4.0-11	---	7.4-8.4	---	0	0	0
	14-60	4.0-11	---	7.4-8.4	---	0	0	0
Ns:								
Ness-----	0-60	16-38	---	6.1-8.4	---	---	---	---
Nw:								
New Cambria-----	0-12	16-39	---	6.6-8.4	---	---	---	---
	12-34	15-36	---	7.9-8.4	---	---	---	---
	34-60	12-30	---	7.9-8.4	---	---	---	---
Pa:								
Plevna-----	0-18	3.0-13	---	6.6-8.4	0	0	0	0
	18-36	3.0-11	---	6.6-8.4	0	0	0	0
	36-60	0.0-4.0	---	6.6-8.4	0	0	0	0
Pr:								
Pratt-----	0-11	1.0-5.0	---	5.6-7.3	---	---	---	---
	11-36	1.0-7.0	---	5.6-7.3	---	---	---	---
	36-60	0.0-5.0	---	6.1-7.3	---	---	---	---
Ro:								
Roxbury-----	0-21	11-24	---	7.4-8.4	1-5	---	---	---
	21-50	7.0-23	---	7.4-8.4	1-5	---	---	---
	50-60	7.0-21	---	7.4-8.4	5-10	---	---	---
Sm:								
Smolan-----	0-11	11-24	---	5.6-7.3	---	---	---	---
	11-16	7.0-21	---	5.6-7.3	---	---	---	---
	16-60	14-30	---	5.6-7.8	---	---	---	---
To:								
Tobin-----	0-20	7.0-19	---	5.6-7.8	---	---	---	---
	20-32	7.0-24	---	7.4-8.4	---	---	---	---
	32-60	7.0-21	---	7.4-8.4	---	---	---	---
W:								
Water-----	---	---	---	---	---	---	---	---
Wb:								
Wells-----	0-15	7.0-19	---	5.6-6.5	0	0	0	0
	15-36	0.0-21	---	5.6-7.3	0	0	0	0
	36-60	4.0-18	---	6.1-7.8	0	0	0	0
Wc:								
Wells-----	0-11	7.0-19	---	5.6-6.5	---	---	---	---
	11-15	10-21	---	5.6-7.3	---	---	---	---
	15-49	10-21	---	5.6-7.3	---	---	---	---
	49-60	4.0-18	---	6.1-7.8	---	0	0	0

WATER FEATURES
McPherson County, Kansas

The Water Features table gives estimates of various water features. The estimates are used in land use planning that involves engineering considerations. Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The four hydrologic soil groups are:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

The months in the table indicate the portion of the year in which the feature is most likely to be a concern.

Water table refers to a saturated zone in the soil. The Water Features table indicates, by month, depth to the top (upper limit) and base (lower limit) of the saturated zone in most years. Estimates of the upper and lower limits are based mainly on observations of the water table at selected sites and on evidence of a saturated zone, namely grayish colors or mottles (redoximorphic features) in the soil. A saturated zone that lasts for less than a month is not considered a water table. Ponding is standing water in a closed depression. Unless a drainage system is installed, the water is removed only by percolation, transpiration, or evaporation. The Water Features table indicates surface water depth and the duration and frequency of ponding. Duration is expressed as very brief if less than 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, rare, occasional, and frequent. None means that ponding is not probable; rare that it is unlikely but possible under unusual weather conditions (the chance of ponding is nearly 0 percent to 5 percent in any year); occasional that it occurs, on the average, once or less in 2 years (the chance of ponding is 5 to 50 percent in any year); and frequent that it occurs, on the average, more than once in 2 years (the chance of ponding is more than 50 percent in any year).

Flooding, the temporary inundation of an area, is caused by overflowing streams, by runoff from adjacent slopes, or by tides. Water standing for short periods after rainfall or snowmelt is not considered flooding, and water standing in swamps and marshes is considered ponding rather than flooding.

Duration and frequency are estimated. Duration is expressed as extremely brief if 0.1 hour to 4 hours, very brief if 4 hours to 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, very rare, rare, occasional, frequent, and very frequent. None means that flooding is not probable; very rare that it is very unlikely but possible under extremely unusual weather conditions (the chance of flooding is less than 1 percent in any year); rare that it is unlikely but possible under unusual weather conditions (the chance of flooding is 1 to 5 percent in any year); occasional that it occurs infrequently under normal weather conditions (the chance of flooding is 5 to 50 percent in any year); frequent that it is likely to occur often under normal weather conditions (the chance of flooding is more than 50 percent in any year but is less than 50 percent in all months in any year); and very frequent that it is likely to occur very often under normal weather conditions (the chance of flooding is more than 50 percent in all months of any year).

The information is based on evidence in the soil profile, namely thin strata of gravel, sand, silt, or clay deposited by floodwater; irregular decrease in organic matter content with increasing depth; and little or no horizon development.

Also considered are local information about the extent and levels of flooding and the relation of each soil on the landscape to historic floods. Information on the extent of flooding based on soil data is less specific than that provided by detailed engineering surveys that delineate flood-prone areas at specific flood frequency levels.

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
053ED: Edalgo-----	C		Ft	Ft	Ft				
053MB: Mccook-----	B		---	---	---	---	---	---	---
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		October	---	---	---	---	---	Very brief	Occasional
053MC: Mccook-----	B								
		January	---	---	---	---	---	---	Rare
		February	---	---	---	---	---	---	Rare
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	---	---	---	---	---	---	Rare
		December	---	---	---	---	---	---	Rare
079GE: Geary-----	B								
079LB: Ladysmith-----	D								
115IC: Irwin-----	D								
159DP: Dillwyn-----	A								
		January	1.0-3.0	>6.0	---	---	---	---	None
		February	1.0-3.0	>6.0	---	---	---	---	None
		March	1.0-3.0	>6.0	---	---	---	---	None
		April	1.0-3.0	>6.0	---	---	---	---	None
		May	1.0-3.0	>6.0	---	---	---	---	None
		June	1.0-3.0	>6.0	---	---	---	---	None
		July	1.0-3.0	>6.0	---	---	---	---	None
		August	1.0-3.0	>6.0	---	---	---	---	None
		September	1.0-3.0	>6.0	---	---	---	---	None
		October	1.0-3.0	>6.0	---	---	---	---	None
		November	1.0-3.0	>6.0	---	---	---	---	None
		December	1.0-3.0	>6.0	---	---	---	---	None
Plevna-----	D								
		January	0.0-2.0	>6.0	---	---	---	---	None
		February	0.0-2.0	>6.0	---	---	---	---	None
		March	0.0-2.0	>6.0	---	---	---	Long	Frequent
		April	0.0-2.0	>6.0	---	---	---	Long	Frequent
		May	0.0-2.0	>6.0	---	---	---	Long	Frequent
		June	0.0-2.0	>6.0	---	---	---	Long	Frequent
		July	0.0-2.0	>6.0	---	---	---	Long	Frequent
		August	0.0-2.0	>6.0	---	---	---	Long	Frequent
		September	0.0-2.0	>6.0	---	---	---	Long	Frequent
		October	0.0-2.0	>6.0	---	---	---	Long	Frequent
		November	0.0-2.0	>6.0	---	---	---	---	None
		December	0.0-2.0	>6.0	---	---	---	---	None
159GC: Geary-----	B								
Clark-----	B								
159LA: Lancaster-----	B								
159NF: Naron-----	B								
159PF: Pratt-----	A								
159PR: Pratt-----	A								

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Carwile-----	D		Ft	Ft	Ft				
		January	0.0	>6.0	0.0-1.0	Brief	---	---	None
		February	0.0	>6.0	0.0-1.0	Brief	---	---	None
		March	0.0	>6.0	0.0-1.0	Brief	---	---	None
		April	0.0	>6.0	0.0-1.0	Brief	---	---	None
		October	0.0	>6.0	0.0-1.0	Brief	---	---	None
		November	0.0	>6.0	0.0-1.0	Brief	---	---	None
		December	0.0	>6.0	0.0-1.0	Brief	---	---	None
159SO: Smolan-----	C		---	---	---	---	---	---	---
159WA: Waldeck-----	C								
		January	2.0-4.0	>6.0	---	---	---	---	None
		February	2.0-4.0	>6.0	---	---	---	---	None
		March	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		April	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		May	---	---	---	---	---	Brief	Occasional
		June	---	---	---	---	---	Brief	Occasional
		July	---	---	---	---	---	Brief	Occasional
		August	---	---	---	---	---	Brief	Occasional
		September	---	---	---	---	---	Brief	Occasional
		October	2.0-4.0	>6.0	---	---	---	Brief	Occasional
		November	2.0-4.0	>6.0	---	---	---	---	None
		December	2.0-4.0	>6.0	---	---	---	---	None
169CE: Clime-----	C		---	---	---	---	---	---	---
169ED: Edalgo-----	C		---	---	---	---	---	---	---
169RO: Roxbury-----	B								
		January	---	---	---	---	---	---	Rare
		February	---	---	---	---	---	---	Rare
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	---	---	---	---	---	---	Rare
		December	---	---	---	---	---	---	Rare
1191: Blazefork-----	D								
		January	4.0	>6.0	---	---	---	---	Rare
		February	4.0	>6.0	---	---	---	---	Rare
		March	4.0	>6.0	---	---	---	---	Rare
		April	4.0	>6.0	---	---	---	---	Rare
		May	4.0	>6.0	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	4.0	>6.0	---	---	---	---	Rare
		December	4.0	>6.0	---	---	---	---	Rare
1200: Buhler-----	D								
		January	---	---	---	---	---	---	Rare
		February	5.0	>6.0	---	---	---	---	Rare
		March	5.0	>6.0	---	---	---	---	Rare
		April	5.0	>6.0	---	---	---	---	Rare
		May	5.0	>6.0	---	---	---	---	Rare
		June	5.0	>6.0	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	---	---	---	---	---	---	Rare
		December	---	---	---	---	---	---	Rare

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Blazefork-----	D		Ft	Ft	Ft				
		January	4.0	>6.0	---	---	---	---	Rare
		February	4.0	>6.0	---	---	---	---	Rare
		March	4.0	>6.0	---	---	---	---	Rare
		April	4.0	>6.0	---	---	---	---	Rare
		May	4.0	>6.0	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	4.0	>6.0	---	---	---	---	Rare
		December	4.0	>6.0	---	---	---	---	Rare
1324: Carway-----	D								
		January	0.0	2.0	0.3-1.0	Long	Occasional	---	None
		February	0.0	2.0	0.3-1.0	Long	Occasional	---	None
		March	0.0	2.0	0.3-1.0	Long	Frequent	---	None
		April	0.0	2.0	0.3-1.0	Long	Frequent	---	None
		May	0.0	2.0	0.3-1.0	Long	Frequent	---	None
		June	0.0	2.0	0.3-1.0	Long	Frequent	---	None
		July	---	---	0.3-1.0	Long	Occasional	---	None
		August	---	---	0.3-1.0	Brief	Rare	---	None
		September	---	---	0.3-1.0	Brief	Rare	---	None
		October	---	---	0.3-1.0	Long	Occasional	---	None
		November	---	---	0.3-1.0	Long	Occasional	---	None
		December	0.0	2.0	0.3-1.0	Long	Occasional	---	None
Carbika-----	D								
		January	0.0	2.0	0.3-1.0	Long	Occasional	---	None
		February	0.0	2.0	0.3-1.0	Long	Occasional	---	None
		March	0.0	2.0	0.3-1.0	Long	Frequent	---	None
		April	0.0	2.0	0.3-1.0	Long	Frequent	---	None
		May	0.0	2.0	0.3-1.0	Long	Frequent	---	None
		June	0.0	2.0	0.3-1.0	Long	Frequent	---	None
		July	---	---	0.3-1.0	Long	Occasional	---	None
		August	---	---	0.0-0.5	Brief	Rare	---	None
		September	---	---	0.0-0.5	Brief	Rare	---	None
		October	---	---	0.3-1.0	Long	Occasional	---	None
		November	---	---	0.3-1.0	Long	Occasional	---	None
		December	0.0	2.0	0.3-1.0	Long	Occasional	---	None
1553: Darlow-----	C								
			---	---	---	---	---	---	---
Elmer-----	C								
			---	---	---	---	---	---	---
1555: Dillhut-----	B								
			---	---	---	---	---	---	---
Plev-----	B								
			---	---	---	---	---	---	---
		February	0.5	4.0	---	---	---	---	None
		March	0.5	4.0	---	---	---	---	None
		April	0.5	4.0	---	---	---	---	None
		May	0.5	4.0	---	---	---	---	None
1985: Hayes-----	B								
			---	---	---	---	---	---	---
2391: Kaskan-----	B								
			---	---	---	---	---	---	---
		January	---	---	---	---	---	Brief	Frequent
		February	5.0	>6.0	---	---	---	Brief	Frequent
		March	5.0	>6.0	---	---	---	Brief	Frequent
		April	5.0	>6.0	---	---	---	Brief	Frequent
		May	5.0	>6.0	---	---	---	Brief	Frequent
		June	5.0	>6.0	---	---	---	Brief	Frequent
		July	---	---	---	---	---	Brief	Frequent
		August	---	---	---	---	---	Brief	Frequent
		September	---	---	---	---	---	Brief	Frequent
		October	---	---	---	---	---	Brief	Frequent
		November	---	---	---	---	---	Brief	Frequent
		December	---	---	---	---	---	Brief	Frequent
2588: Longford, Moderately Eroded-----	C								
			---	---	---	---	---	---	---
3181: Pratt-----	A								
			---	---	---	---	---	---	---
Turon-----	A								
			---	---	---	---	---	---	---
3641: Tivin-----	A								
			---	---	---	---	---	---	---
Dillhut-----	B								
			---	---	---	---	---	---	---

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
At: Attica-----	B		Ft	Ft	Ft				
Br: Bridgeport-----	B		---	---	---	---	---	---	---
		January	---	---	---	---	---	Brief	Rare
		February	---	---	---	---	---	Brief	Rare
		March	---	---	---	---	---	Brief	Rare
		April	---	---	---	---	---	Brief	Rare
		May	---	---	---	---	---	Brief	Rare
		June	---	---	---	---	---	Brief	Rare
		July	---	---	---	---	---	Brief	Rare
		August	---	---	---	---	---	Brief	Rare
		September	---	---	---	---	---	Brief	Rare
		October	---	---	---	---	---	Brief	Rare
		November	---	---	---	---	---	Brief	Rare
		December	---	---	---	---	---	Brief	Rare
Ca: Carwile-----	D								
		January	0.0	>6.0	0.0-1.0	Brief	---	---	None
		February	0.0	>6.0	0.0-1.0	Brief	---	---	None
		March	0.0	>6.0	0.0-1.0	Brief	---	---	None
		April	0.0	>6.0	0.0-1.0	Brief	---	---	None
		May	---	---	0.0-	---	---	---	None
		June	---	---	0.0-	---	---	---	None
		July	---	---	0.0-	---	---	---	None
		August	---	---	0.0-	---	---	---	None
		September	---	---	0.0-	---	---	---	None
		October	0.0	>6.0	0.0-1.0	Brief	---	---	None
		November	0.0	>6.0	0.0-1.0	Brief	---	---	None
		December	0.0	>6.0	0.0-1.0	Brief	---	---	None
Cb: Cass-----	B								
		March	---	---	---	---	---	Brief	Rare
		April	---	---	---	---	---	Brief	Rare
		May	---	---	---	---	---	Brief	Rare
		June	---	---	---	---	---	Brief	Rare
Ce: Clime-----	C								
Cm: Clime-----	C								
Cr: Crete-----	C								
Cs: Crete-----	C								
Ct: Crete-----	C								
De: Detroit-----	C								
		January	---	---	---	---	---	---	Rare
		February	---	---	---	---	---	---	Rare
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	---	---	---	---	---	---	Rare
		December	---	---	---	---	---	---	Rare
Dr: Drummond-----	D								
		January	2.0-6.0	>6.0	---	---	---	---	None
		February	2.0-6.0	>6.0	---	---	---	---	None
		March	2.0-6.0	>6.0	---	---	---	---	None
		April	2.0-6.0	>6.0	---	---	---	---	None
		November	2.0-6.0	>6.0	---	---	---	---	None
		December	2.0-6.0	>6.0	---	---	---	---	None
Ed: Edalgo-----	C								
Ee: Edalgo-----	C								
Fa: Farnum-----	B								

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Ge: Geary-----	B		Ft	Ft	Ft				
Go: Goessel-----	D		---	---	---	---	---	---	---
		April	2.0-3.0	>6.0	---	---	---	---	None
		May	2.0-3.0	>6.0	---	---	---	---	None
		June	2.0-3.0	>6.0	---	---	---	---	None
Ho: Hord-----	B								
		January	---	---	---	---	---	---	Rare
		February	---	---	---	---	---	---	Rare
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	---	---	---	---	---	---	Rare
		December	---	---	---	---	---	---	Rare
INT: Aquolls-----	C								
		March	0.0	>6.0	0.0-0.8	Brief	Occasional	---	None
		April	0.0	>6.0	0.0-0.8	Brief	Occasional	---	None
		May	0.0	>6.0	0.0-0.8	Brief	Occasional	---	None
		June	0.0	>6.0	0.0-0.8	Brief	Occasional	---	None
Ir: Irwin, bedrock substratum-	D								
			---	---	---	---	---	---	---
La: Ladysmith-----	D								
			---	---	---	---	---	---	---
Ld: Ladysmith-----	D								
			---	---	---	---	---	---	---
Drummond-----	D								
		January	2.0-6.0	>6.0	---	---	---	---	None
		February	2.0-6.0	>6.0	---	---	---	---	None
		March	2.0-6.0	>6.0	---	---	---	---	None
		April	2.0-6.0	>6.0	---	---	---	---	None
		November	2.0-6.0	>6.0	---	---	---	---	None
		December	2.0-6.0	>6.0	---	---	---	---	None
Le: Lancaster-----	B								
			---	---	---	---	---	---	---
Lh: Lancaster-----	B								
			---	---	---	---	---	---	---
Hedville-----	D								
			---	---	---	---	---	---	---
LHH: Lancaster-----	B								
			---	---	---	---	---	---	---
Hedville-----	D								
			---	---	---	---	---	---	---
Ln: Longford-----	C								
			---	---	---	---	---	---	---
Lo: Longford-----	C								
			---	---	---	---	---	---	---
Mc: Mccook-----	B								
		January	---	---	---	---	---	---	Rare
		February	---	---	---	---	---	---	Rare
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	---	---	---	---	---	---	Rare
		December	---	---	---	---	---	---	Rare
Ns:									

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Soil Saturation		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
Ness-----	D		Ft	Ft	Ft				
		January	---	---	0.0-	---	---	---	None
		February	---	---	0.0-	---	---	---	None
		March	0.0	>6.0	0.0-1.0	Long	---	---	None
		April	0.0	>6.0	0.0-1.0	Long	---	---	None
		May	0.0	>6.0	0.0-1.0	Long	---	---	None
		June	0.0	>6.0	0.0-1.0	Long	---	---	None
		July	---	---	0.0-	---	---	---	None
		August	---	---	0.0-	---	---	---	None
		September	---	---	0.0-	---	---	---	None
		October	---	---	0.0-	---	---	---	None
		November	---	---	0.0-	---	---	---	None
		December	---	---	0.0-	---	---	---	None
Nw: New Cambria-----	C								
		January	---	---	---	---	---	---	Rare
		February	---	---	---	---	---	---	Rare
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	---	---	---	---	---	---	Rare
		December	---	---	---	---	---	---	Rare
Pa: Plevna-----	D								
		January	0.0-2.0	>6.0	---	---	---	---	None
		February	0.0-2.0	>6.0	---	---	---	---	None
		March	0.0-2.0	>6.0	---	---	---	Long	Frequent
		April	0.0-2.0	>6.0	---	---	---	Long	Frequent
		May	0.0-2.0	>6.0	---	---	---	Long	Frequent
		June	0.0-2.0	>6.0	---	---	---	Long	Frequent
		July	0.0-2.0	>6.0	---	---	---	Long	Frequent
		August	0.0-2.0	>6.0	---	---	---	Long	Frequent
		September	0.0-2.0	>6.0	---	---	---	Long	Frequent
		October	0.0-2.0	>6.0	---	---	---	Long	Frequent
		November	0.0-2.0	>6.0	---	---	---	---	None
		December	0.0-2.0	>6.0	---	---	---	---	None
Pr: Pratt-----	A								
			---	---	---	---	---	---	---
Ro: Roxbury-----	B								
		January	---	---	---	---	---	---	Rare
		February	---	---	---	---	---	---	Rare
		March	---	---	---	---	---	---	Rare
		April	---	---	---	---	---	---	Rare
		May	---	---	---	---	---	---	Rare
		June	---	---	---	---	---	---	Rare
		July	---	---	---	---	---	---	Rare
		August	---	---	---	---	---	---	Rare
		September	---	---	---	---	---	---	Rare
		October	---	---	---	---	---	---	Rare
		November	---	---	---	---	---	---	Rare
		December	---	---	---	---	---	---	Rare
Sm: Smolan-----	C								
			---	---	---	---	---	---	---
To: Tobin-----	B								
		April	---	---	---	---	---	Very brief	Occasional
		May	---	---	---	---	---	Very brief	Occasional
		June	---	---	---	---	---	Very brief	Occasional
		July	---	---	---	---	---	Very brief	Occasional
		August	---	---	---	---	---	Very brief	Occasional
		September	---	---	---	---	---	Very brief	Occasional
		October	---	---	---	---	---	Very brief	Occasional
W: Water-----	---								
			---	---	---	---	---	---	---
Wb: Wells-----	B								
			---	---	---	---	---	---	---
Wc: Wells-----	B								
			---	---	---	---	---	---	---

The following table gives estimates of various soil features. The estimates are used in land use planning that involves engineering considerations.

A restrictive layer is a nearly continuous layer that has one or more physical, chemical, or thermal properties that significantly impede the movement of water and air through the soil or that restrict roots or otherwise provide an unfavorable root environment. Examples are bedrock, cemented layers, dense layers, and frozen layers. The table indicates the hardness and thickness of the restrictive layer, both of which significantly affect the ease of excavation. Depth to top is the vertical distance from the soil surface to the upper boundary of the restrictive layer.

Potential for frost action is the likelihood of upward or lateral expansion of the soil caused by the formation of segregated ice lenses (frost heave) and the subsequent collapse of the soil and loss of strength on thawing. Frost action occurs when moisture moves into the freezing zone of the soil. Temperature, texture, density, permeability, content of organic matter, and depth to the water table are the most important factors considered in evaluating the potential for frost action. It is assumed that the soil is not insulated by vegetation or snow and is not artificially drained. Silty and highly structured, clayey soils that have a high water table in winter are the most susceptible to frost action. Well drained, very gravelly, or very sandy soils are the least susceptible. Frost heave and low soil strength during thawing cause damage to pavements and other rigid structures.

Risk of corrosion pertains to potential soil-induced electrochemical or chemical action that corrodes or weakens uncoated steel or concrete. The rate of corrosion of uncoated steel is related to such factors as soil moisture, particle-size distribution, acidity, and electrical conductivity of the soil. The rate of corrosion of concrete is based mainly on the sulfate and sodium content, texture, moisture content, and acidity of the soil. Special site examination and design may be needed if the combination of factors results in a severe hazard of corrosion. The steel or concrete in installations that intersect soil boundaries or soil layers is more susceptible to corrosion than the steel or concrete in installations that are entirely within one kind of soil or within one soil layer.

For uncoated steel, the risk of corrosion, expressed as low, moderate, or high, is based on soil drainage class, total acidity, electrical resistivity near field capacity, and electrical conductivity of the saturation extract.

For concrete, the risk of corrosion also is expressed as low, moderate, or high. It is based on soil texture, acidity, and amount of sulfates in the saturation extract.

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness		Uncoated Steel	Concrete
053ED: Edalgo-----	20-40	In Bedrock (paralithic)	---	---	Low	Moderate	Low
053MB: Mccook-----	---	---	---	---	Low	Low	Low
053MC: Mccook-----	---	---	---	---	Low	Low	Low
079GE: Geary-----	---	---	---	---	Low	Low	Low
079LB: Ladysmith-----	---	---	---	---	Low	High	Low
115IC: Irwin-----	---	---	---	---	Moderate	High	Low
159DP: Dillwyn-----	---	---	---	---	Low	Low	Low
159GC: Plevna-----	---	---	---	---	Low	High	Low
159GC: Geary-----	---	---	---	---	Low	Low	Low
159LA: Clark-----	---	---	---	---	Low	Moderate	Low
159LA: Lancaster-----	20-40	Bedrock (paralithic)	---	Moderately cemented	Low	Low	Moderate
159NF: Naron-----	---	---	---	---	Low	Low	Low
159PF: Pratt-----	---	---	---	---	Low	Low	Moderate
159PR: Pratt-----	---	---	---	---	Low	Low	Moderate
159SO: Carwile-----	---	---	---	---	Low	High	Moderate
159SO: Smolan-----	---	---	---	---	Low	Moderate	Low
159WA: Waldeck-----	---	---	---	---	Low	Moderate	Low
169CE: Clime-----	20-40	Bedrock (paralithic)	---	Moderately cemented	Low	High	Low
169ED: Edalgo-----	20-40	Bedrock (paralithic)	---	---	Low	Moderate	Low
169RO: Roxbury-----	---	---	---	---	Low	Low	Low
1191: Blazefork-----	---	---	---	---	Low	High	Low
1200: Buhler-----	---	---	---	---	Low	High	Low
1324: Blazefork-----	---	---	---	---	Low	High	Low
1553: Carway-----	---	---	---	---	Low	High	Moderate
1553: Carbika-----	---	---	---	---	Low	Moderate	Low
1553: Darlow-----	---	---	---	---	Low	High	Low
1555: Elmer-----	---	---	---	---	Low	High	Low
1555: Dillhut-----	---	---	---	---	Low	Low	Moderate
1985: Plev-----	---	---	---	---	Low	High	Moderate
2391: Hayes-----	---	---	---	---	Low	Moderate	Low
2588: Kaskan-----	---	---	---	---	Low	High	Moderate
3181: Longford, Moderately Eroded-----	---	---	---	---	Moderate	High	Low
3641: Pratt-----	---	---	---	---	Low	Low	Moderate
3641: Turon-----	---	---	---	---	Low	Low	Moderate
3641: Tivin-----	---	---	---	---	Low	Low	Low
AED: Dillhut-----	---	---	---	---	Low	Low	Moderate
AED: Arents, Earthen Dam-----	---	---	---	---	---	---	---
At: Attica-----	---	---	---	---	Low	Low	Low
Br: Bridgeport-----	---	---	---	---	Low	Low	Low
Ca: Carwile-----	---	---	---	---	Low	High	Moderate
Cb: Cass-----	---	---	---	---	Low	Moderate	Low
Ce: Clime-----	20-40	Bedrock (paralithic)	---	Moderately cemented	Low	High	Low
Cm: Clime-----	20-40	Bedrock (paralithic)	---	Moderately cemented	Low	High	Low

Map symbol and soil name	Restrictive layer				Potential for Frost action	Risk of corrosion	
	Kind	Depth to top In	Thickness In	Hardness		Uncoated Steel	Concrete
Cr: Crete-----	---	---	---	---	Low	Moderate	Low
Cs: Crete-----	---	---	---	---	Low	Moderate	Low
Ct: Crete-----	---	---	---	---	Low	Moderate	Low
De: Detroit-----	---	---	---	---	Low	High	Low
Dr: Drummond-----	---	---	---	---	Low	High	High
Ed: Edalgo-----	20-40	Bedrock (paralithic)	---	---	Low	Moderate	Low
Ee: Edalgo-----	20-40	Bedrock (paralithic)	---	Moderately cemented	Low	Moderate	Low
Fa: Farnum-----	---	---	---	---	Low	Moderate	Low
Ge: Geary-----	---	---	---	---	Low	Low	Low
Go: Goessel-----	---	---	---	---	Low	High	Low
Ho: Hord-----	---	---	---	---	Low	High	Low
INT: Aguolls-----	---	---	---	---	Low	---	---
Ir: Irwin, bedrock substratum-----	40-60	Bedrock (paralithic)	---	---	Low	High	Low
La: Ladysmith-----	---	---	---	---	Low	High	Low
Ld: Ladysmith-----	---	---	---	---	Low	High	Low
Le: Lancaster-----	20-40	Bedrock (paralithic)	---	Moderately cemented	Low	Low	Moderate
Lh: Lancaster-----	20-40	Bedrock (paralithic)	---	Moderately cemented	Low	Low	Moderate
Hedville-----	4-20	Bedrock (lithic)	---	Strongly cemented	Low	Low	Moderate
LHH: Lancaster-----	20-40	Bedrock (paralithic)	---	Moderately cemented	Low	Low	Moderate
Hedville-----	4-20	Bedrock (lithic)	---	Strongly cemented	Low	Low	Moderate
Ln: Longford-----	---	---	---	---	Low	High	Low
Lo: Longford-----	---	---	---	---	Low	High	Low
Mc: Mccook-----	---	---	---	---	Low	Low	Low
Ns: Ness-----	---	---	---	---	Low	High	Low
Nw: New Cambria-----	---	---	---	---	Low	High	Low
Pa: Plevna-----	---	---	---	---	Low	High	Low
Pr: Pratt-----	---	---	---	---	Low	Low	Moderate
Ro: Roxbury-----	---	---	---	---	Low	Low	Low
Sm: Smolan-----	---	---	---	---	Low	Moderate	Low
To: Tobin-----	---	---	---	---	Low	Low	Low
W: Water-----	---	---	---	---	Low	---	---
Wb: Wells-----	---	---	---	---	Moderate	Low	Moderate
Wc: Wells-----	---	---	---	---	Low	Low	Moderate

WATER MANAGEMENT
McPherson County, Kansas

The soils of the survey area are rated in the Water Management table according to limitations that affect their suitability for water management. Soils are rated for pond reservoir areas, drainage, irrigation, terraces and diversions, and grassed waterways. Restrictive features that affect each soil for the specified use is also provided in the table.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the specified use. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Moderately limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Limited indicates that the soil has one or more features that are significant limitations for the specified use. The limitations can be overcome, but generally require special design, soil reclamation, or installation procedures that may result in additional expense. Fair performance and moderate to high maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Limitation class terms, such as very limited or limited, etc., limitation ratings, and numerical ratings are shown for each soil feature listed. As many as three soil features may be listed for each soil component if applicable. The overall limitation rating for the soil component is based on the most severe limitation.

Pond reservoir areas hold water behind a dam or embankment. Soils best suited to this use have low seepage potential in the upper 60 inches. The seepage potential is determined by the permeability of the soil and the depth to fractured bedrock or other permeable material. Excessive slope can affect the storage capacity of the reservoir area.

Embankments, dikes, and levees are raised structures of soil material, generally less than 20 feet high, constructed to impound water or to protect land against overflow. In this table, the soils are rated as a source of material for embankment fill. The ratings apply to the soil material below the surface layer to a depth of about 5 feet. It is assumed that soil layers will be uniformly mixed and compacted during construction.

The ratings do not indicate the ability of the natural soil to support an embankment. Soil properties to a depth even greater than the height of the embankment can affect performance and safety of the embankment. Generally, deeper onsite investigation is needed to determine these properties.

Soil material in embankments must be resistant to seepage, piping, and erosion and have favorable compaction characteristics. Unfavorable features include less than 5 feet of suitable material and a high content of stones or boulders, organic matter, or salts or sodium. A high water table affects the amount of usable material. It also affects traffic ability.

Aquifer-fed excavated ponds are pits or dugouts that extend to a ground-water aquifer or to a depth below a permanent water table. Excluded are ponds that are fed only by surface runoff and embankment ponds that impound water 3 feet or more above the original surface. Excavated ponds are affected by depth to a permanent water table, permeability of the aquifer, and quality of the water as inferred from the salinity of the soil. Depth to bedrock and the content of large stones affect the ease of excavation.

Drainage is the removal of excess surface and subsurface water from the soil. How easily and effectively the soil is drained depends on the depth to bedrock, to a cemented pan, or to other layers that affect the rate of water movement; permeability; depth to a high water table or depth of standing water if the soil is subject to ponding; slope; susceptibility to flooding; subsidence of organic layers; and the potential for frost action. Excavating and grading and the stability of ditch banks are affected by depth to bedrock or to a cemented pan, large stones, slope, and the hazard of cutbanks caving. The productivity of the soil after drainage is adversely affected by extreme acidity or by toxic substances in the root zone, such as salts, sodium, and sulfur. Availability of drainage outlets is not considered in the ratings.

Irrigation is the controlled application of water to supplement rainfall and support plant growth. The design and management of an irrigation system are affected by depth to the water table, the need for drainage, flooding, available water capacity, intake rate, permeability, erosion hazard, and slope. The construction of a system is affected by large stones and depth to bedrock or to a cemented pan. The performance of a system is affected by the depth of the root zone, the amount of salts or sodium, and soil reaction.

Terraces and diversions are embankments or a combination of channels and ridges constructed across a slope to control erosion and conserve moisture by intercepting runoff. Slope, wetness, large stones, and depth to bedrock or to a cemented pan affect the construction of terraces and diversions. A restricted rooting depth, a very limited hazard of wind erosion or water erosion, an excessively coarse texture, and restricted permeability adversely affect maintenance.

Grassed waterways are natural or constructed channels, generally broad and shallow, which conduct surface water to outlets at a non-erosive velocity. Large stones, wetness, slope, and depth to bedrock or to a cemented pan affect the construction of grassed waterways. A hazard of wind erosion, low available water capacity, restricted rooting depth, toxic substances such as salts and sodium, and restricted permeability adversely affect the growth and maintenance of the grass after construction.

WATER MANAGEMENT--Continued
McPherson County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
053ED: Edalgo-----	Limitation: deep to water	Limitation: percs slowly slope thin layer	Limitation: area reclaim erodes easily	Limitation: area reclaim erodes easily
053MB: Mccook-----	Limitation: deep to water	Limitation: flooding	Limitation: erodes easily	Limitation: erodes easily
053MC: Mccook-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
079GE: Geary-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
079LB: Ladysmith-----	Limitation: percs slowly	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily percs slowly wetness
115IC: Irwin-----	Limitation: deep to water	Limitation: erodes easily percs slowly slope	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly
159DP: Dillwyn-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: wetness droughty
Plevna-----	Limitation: flooding cutbanks cave	Limitation: fast intake wetness soil blowing	Limitation: too sandy wetness soil blowing	Limitation: wetness
159GC: Geary-----	Limitation: deep to water	Limitation: slope	Limitation: erodes easily	Limitation: erodes easily
Clark-----	Limitation: deep to water	Limitation: slope	Favorable	Favorable
159LA: Lancaster-----	Limitation: deep to water	Limitation: thin layer	Limitation: area reclaim	Limitation: area reclaim
159NF: Naron-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable
159PF: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
159PR: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
Carwile-----	Limitation: percs slowly	Limitation: percs slowly wetness soil blowing	Limitation: erodes easily wetness soil blowing	Limitation: erodes easily percs slowly rooting depth
159SO: Smolan-----	Limitation: deep to water	Limitation: erodes easily percs slowly slope	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly
159WA: Waldeck-----	Limitation: flooding cutbanks cave	Limitation: flooding wetness soil blowing	Limitation: too sandy wetness soil blowing	Favorable
169CE: Clime-----	Limitation: deep to water	Limitation: percs slowly slope depth to rock	Limitation: erodes easily percs slowly depth to rock	Limitation: erodes easily percs slowly depth to rock
169ED: Edalgo-----	Limitation: deep to water	Limitation: percs slowly slope depth to rock	Limitation: erodes easily percs slowly depth to rock	Limitation: erodes easily percs slowly depth to rock
169RO: Roxbury-----	Limitation: deep to water	---	Limitation: erodes easily	Limitation: erodes easily
1191: Blazefork-----	Limitation: deep to water	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly

WATER MANAGEMENT--Continued
McPherson County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
1200: Buhler-----	Limitation: excess sodium percs slowly	Limitation: percs slowly wetness	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily excess sodium percs slowly
Blazefork-----	Limitation: deep to water	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly
1324: Carway-----	Limitation: percs slowly	Limitation: wetness soil blowing	Limitation: erodes easily wetness soil blowing	Limitation: erodes easily percs slowly wetness
Carbika-----	Limitation: percs slowly ponding	Limitation: percs slowly soil blowing ponding	Limitation: erodes easily soil blowing ponding	Limitation: erodes easily percs slowly wetness
1553: Darlow-----	Limitation: deep to water	Limitation: excess sodium excess salt percs slowly	Limitation: percs slowly	Limitation: excess sodium percs slowly
Elmer-----	Limitation: deep to water	Limitation: excess sodium soil blowing	Limitation: erodes easily soil blowing	Limitation: erodes easily excess sodium
1555: Dillhut-----	Favorable	Limitation: wetness droughty	Limitation: wetness soil blowing	Limitation: droughty
Plev-----	Limitation: cutbanks cave	Limitation: fast intake wetness droughty	Limitation: too sandy wetness soil blowing	Limitation: wetness droughty
1985: Hayes-----	Limitation: deep to water	Limitation: fast intake soil blowing	Limitation: soil blowing	Favorable
2391: Kaskan-----	Limitation: deep to water	Limitation: erodes easily flooding	Limitation: erodes easily	Limitation: erodes easily
2588: Longford, Moderately Eroded-----	Limitation: deep to water	Limitation: percs slowly slope	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly
3181: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
Turon-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: too sandy soil blowing	Limitation: droughty
3641: Tivin-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Dillhut-----	Favorable	Limitation: wetness droughty	Limitation: wetness soil blowing	Limitation: droughty
AED: Arents, Earthen Dam-----	---	---	---	---
At: Attica-----	Limitation: deep to water	Limitation: fast intake soil blowing	Limitation: soil blowing	Favorable
Br: Bridgeport-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
Ca: Carwile-----	Limitation: percs slowly	Limitation: percs slowly wetness soil blowing	Limitation: erodes easily wetness soil blowing	Limitation: erodes easily percs slowly rooting depth
Cb: Cass-----	Limitation: deep to water	Limitation: soil blowing	Limitation: soil blowing	Favorable

WATER MANAGEMENT--Continued
McPherson County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Ce: Clime-----	Limitation: deep to water	Limitation: percs slowly slow intake depth to rock	Limitation: percs slowly depth to rock	Limitation: percs slowly depth to rock
Cm: Clime-----	Limitation: deep to water	Limitation: percs slowly slope slow intake	Limitation: percs slowly depth to rock	Limitation: percs slowly depth to rock
Cr: Crete-----	Limitation: deep to water	Limitation: erodes easily percs slowly	Limitation: erodes easily	Limitation: erodes easily percs slowly
Cs: Crete-----	Limitation: deep to water	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly
Ct: Crete-----	Limitation: deep to water	Limitation: erodes easily percs slowly	Limitation: erodes easily	Limitation: erodes easily percs slowly
De: Detroit-----	Limitation: deep to water	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly
Dr: Drummond-----	Limitation: excess sodium percs slowly	Limitation: percs slowly wetness droughty	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily excess sodium droughty
Ed: Edalgo-----	Limitation: deep to water	Limitation: percs slowly slope depth to rock	Limitation: erodes easily slope depth to rock	Limitation: erodes easily slope depth to rock
Ee: Edalgo-----	Limitation: deep to water	Limitation: percs slowly slope depth to rock	Limitation: erodes easily percs slowly depth to rock	Limitation: erodes easily percs slowly depth to rock
Fa: Farnum-----	Limitation: deep to water	Favorable	Favorable	Favorable
Ge: Geary-----	Limitation: deep to water	Favorable	Limitation: erodes easily	Limitation: erodes easily
Go: Goessel-----	Limitation: percs slowly	Limitation: percs slowly slow intake wetness	Limitation: percs slowly wetness	Limitation: percs slowly
Ho: Hord-----	Limitation: deep to water	Favorable	Favorable	Favorable
INT: Aquolls-----	---	---	---	---
Ir: Irwin, bedrock substratum-----	Limitation: deep to water	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly
La: Ladysmith-----	Limitation: percs slowly	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily percs slowly wetness
Ld: Ladysmith-----	Limitation: percs slowly	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily percs slowly wetness
Drummond-----	Limitation: excess sodium percs slowly	Limitation: percs slowly wetness droughty	Limitation: erodes easily percs slowly wetness	Limitation: erodes easily excess sodium droughty
Le: Lancaster-----	Limitation: deep to water	Limitation: slope depth to rock	Limitation: depth to rock	Limitation: depth to rock
Lh: Lancaster-----	Limitation: deep to water	Limitation: slope depth to rock	Limitation: slope depth to rock	Limitation: slope depth to rock

WATER MANAGEMENT--Continued
McPherson County, Kansas

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

Map symbol and soil name	Features affecting--			
	Drainage	Irrigation	Terraces and diversions	Grassed waterways
Hedville-----	Limitation: deep to water	Limitation: slope depth to rock	Limitation: slope depth to rock	Limitation: slope depth to rock
LHH: Lancaster-----	Limitation: deep to water	Limitation: slope depth to rock	Limitation: depth to rock	Limitation: depth to rock
Hedville-----	Limitation: deep to water	Limitation: slope depth to rock	Limitation: slope depth to rock	Limitation: slope depth to rock
Ln: Longford-----	Limitation: deep to water	Limitation: percs slowly slope	Favorable	Favorable
Lo: Longford-----	Limitation: deep to water	Limitation: percs slowly slope	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly
Mc: Mccook-----	Limitation: deep to water	Limitation: soil blowing	Limitation: erodes easily soil blowing	Limitation: erodes easily
Ns: Ness-----	Limitation: percs slowly	Limitation: percs slowly slow intake wetness	Limitation: percs slowly wetness	Limitation: percs slowly wetness
Nw: New Cambria-----	Limitation: deep to water	Limitation: percs slowly slow intake	Limitation: percs slowly	Limitation: percs slowly
Pa: Plevna-----	Limitation: flooding cutbanks cave	Limitation: flooding wetness soil blowing	Limitation: too sandy wetness soil blowing	Limitation: wetness
Pr: Pratt-----	Limitation: deep to water	Limitation: fast intake slope droughty	Limitation: slope too sandy soil blowing	Limitation: slope droughty
Ro: Roxbury-----	Limitation: deep to water	Limitation: erodes easily	Limitation: erodes easily	Limitation: erodes easily
Sm: Smolan-----	Limitation: deep to water	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly	Limitation: erodes easily percs slowly
To: Tobin-----	Limitation: deep to water	Limitation: flooding	Favorable	Favorable
W: Water-----	---	---	---	---
Wb: Wells-----	Limitation: deep to water	Favorable	Favorable	Favorable
Wc: Wells-----	Limitation: deep to water	Limitation: slope	Favorable	Favorable

WATER MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
053ED: Edalgo-----	100	Somewhat limited Depth to bedrock	0.05	Somewhat limited Thin layer Hard to pack	0.74 0.72	Very limited Deep to water	1.00
053MB: Mccook-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
053MC: Mccook-----	100	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Deep to water	1.00
079GE: Geary-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.06	Very limited Deep to water	1.00
079LB: Ladysmith-----	100	Somewhat limited Seepage	0.01	Somewhat limited Hard to pack	0.99	Very limited Deep to water	1.00
115IC: Irwin-----	90	Not limited		Somewhat limited Hard to pack	0.17	Very limited Deep to water	1.00
159DP: Dillwyn-----	60	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.18	Very limited Cutbanks cave Deep to water	1.00 0.00
Plevna-----	40	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 1.00	Very limited Cutbanks cave	1.00
159GC: Geary-----	75	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.08	Very limited Deep to water	1.00
Clark-----	25	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.13	Very limited Deep to water	1.00
159LA: Lancaster-----	90	Somewhat limited Seepage Depth to bedrock	0.70 0.04	Somewhat limited Thin layer Piping	0.70 0.69	Very limited Deep to water	1.00
159NF: Naron-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.08	Very limited Deep to water	1.00
159PF: Pratt-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
159PR: Pratt-----	65	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Carwile-----	35	Not limited		Very limited Depth to saturated zone Hard to pack Seepage	1.00 0.10 0.07	Somewhat limited Slow refill Cutbanks cave	0.30 0.10
159SO: Smolan-----	100	Not limited		Somewhat limited Hard to pack	0.82	Very limited Deep to water	1.00
159WA: Waldeck-----	100	Very limited Seepage	1.00	Very limited Seepage Depth to saturated zone	1.00 0.43	Very limited Cutbanks cave Deep to water	1.00 0.25
169CE: Clime-----	100	Somewhat limited Depth to bedrock	0.17	Somewhat limited Thin layer	0.91	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
169ED: Edalgo-----	100	Somewhat limited Depth to bedrock	0.11	Hard to pack Somewhat limited Thin layer Hard to pack	0.31 0.85 0.57	Very limited Deep to water	1.00
169RO: Roxbury-----	99	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.65	Very limited Deep to water	1.00
1191: Blazefork-----	90	Somewhat limited Seepage	0.05	Not limited		Somewhat limited Slow refill Deep to water Cutbanks cave	0.95 0.81 0.10
1200: Buhler-----	65	Somewhat limited Seepage	0.70	Very limited Piping Seepage	1.00 0.09	Very limited Deep to water	1.00
Blazefork-----	30	Somewhat limited Seepage	0.05	Not limited		Somewhat limited Slow refill Deep to water Cutbanks cave	0.95 0.81 0.10
1324: Carway-----	50	Somewhat limited Seepage	0.70	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Deep to water	1.00
Carbika-----	30	Somewhat limited Seepage	0.70	Very limited Ponding Depth to saturated zone Piping	1.00 1.00 0.40	Very limited Deep to water	1.00
1553: Darlow-----	70	Somewhat limited Seepage	0.70	Very limited Piping Salinity Seepage	1.00 0.12 0.12	Very limited Deep to water	1.00
Elmer-----	20	Very limited Seepage	1.00	Very limited Piping Seepage	1.00 0.10	Very limited Deep to water	1.00
1555: Dillhut-----	35	Very limited Seepage	1.00	Somewhat limited Seepage	0.13	Very limited Deep to water	1.00
Plev-----	35	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Deep to water	1.00
1985: Hayes-----	60	Very limited Seepage	1.00	Somewhat limited Piping	0.42	Very limited Deep to water	1.00
2391: Kaskan-----	75	Very limited Seepage	1.00	Somewhat limited Seepage	0.43	Very limited Deep to water	1.00
2588: Longford, Moderately Eroded-----	90	Somewhat limited Seepage	 0.05	Somewhat limited Hard to pack	 0.59	Very limited Deep to water	 1.00
3181: Pratt-----	45	Very limited Seepage	1.00	Somewhat limited Seepage	0.86	Very limited Deep to water	1.00
Turon-----	30	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.87 0.77	Very limited Deep to water	1.00

WATER MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
3641: Tivin-----	45	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Deep to water	1.00
Dillhut-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.13	Very limited Deep to water	1.00
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
At: Attica-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.72	Very limited Deep to water	1.00
Br: Bridgeport-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.71	Very limited Deep to water	1.00
Ca: Carwile-----	100	Somewhat limited Seepage	0.57	Very limited Depth to saturated zone	1.00	Somewhat limited Slow refill Cutbanks cave	0.30 0.10
Cb: Cass-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.70	Very limited Deep to water	1.00
Ce: Clime-----	100	Somewhat limited Depth to bedrock	0.19	Somewhat limited Thin layer Hard to pack	0.93 0.50	Very limited Deep to water	1.00
Cm: Clime-----	100	Somewhat limited Depth to bedrock	0.19	Somewhat limited Thin layer Hard to pack	0.93 0.50	Very limited Deep to water	1.00
Cr: Crete-----	100	Somewhat limited Seepage	0.57	Not limited		Very limited Deep to water	1.00
Cs: Crete-----	100	Somewhat limited Seepage	0.57	Somewhat limited Hard to pack	0.36	Very limited Deep to water	1.00
Ct: Crete-----	100	Somewhat limited Seepage	0.57	Not limited		Very limited Deep to water	1.00
De: Detroit-----	100	Somewhat limited Seepage	0.05	Not limited		Very limited Deep to water	1.00
Dr: Drummond-----	100	Not limited		Not limited		Very limited Slow refill Deep to water Cutbanks cave Salty water	1.00 0.81 0.10 0.01
Ed: Edalgo-----	100	Somewhat limited Depth to bedrock	0.11	Somewhat limited Thin layer Hard to pack	0.85 0.50	Very limited Deep to water	1.00
Ee: Edalgo-----	100	Somewhat limited Depth to bedrock	0.34	Somewhat limited Thin layer Hard to pack	0.99 0.72	Very limited Deep to water	1.00
Fa: Farnum-----	100	Somewhat limited Seepage	0.70	Somewhat limited Seepage Piping	0.06 0.01	Very limited Deep to water	1.00
Ge: Geary-----	100	Somewhat limited		Somewhat limited		Very limited	

WATER MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Go: Goessel-----	100	Seepage	0.70	Piping	0.08	Deep to water	1.00
		Not limited		Somewhat limited Hard to pack Depth to saturated zone	0.98 0.86	Very limited Deep to water	1.00
Ho: Hord-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.53	Very limited Deep to water	1.00
INT: Aquolls-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Somewhat limited Cutbanks cave	0.10
Ir: Irwin, bedrock substratum-----	100	Somewhat limited Depth to bedrock	0.00	Somewhat limited Hard to pack Thin layer	0.95 0.11	Very limited Deep to water	1.00
La: Ladysmith-----	100	Somewhat limited Seepage	0.01	Somewhat limited Hard to pack	0.99	Very limited Deep to water	1.00
Ld: Ladysmith-----	70	Somewhat limited Seepage	0.01	Somewhat limited Hard to pack	0.99	Very limited Deep to water	1.00
Drummond-----	30	Not limited		Not limited		Very limited Slow refill Deep to water Cutbanks cave Salty water	1.00 0.81 0.10 0.01
Le: Lancaster-----	100	Somewhat limited Seepage Depth to bedrock	0.70 0.08	Somewhat limited Thin layer Piping	0.81 0.24	Very limited Deep to water	1.00
Lh: Lancaster-----	60	Somewhat limited Seepage Depth to bedrock	0.70 0.08	Somewhat limited Thin layer Piping	0.81 0.24	Very limited Deep to water	1.00
Hedville-----	40	Very limited Seepage Depth to bedrock	1.00 1.00	Very limited Thin layer Piping	1.00 1.00	Very limited Deep to water	1.00
LHH: Lancaster-----	60	Somewhat limited Seepage Depth to bedrock	0.70 0.11	Somewhat limited Thin layer Piping	0.86 0.58	Very limited Deep to water	1.00
Hedville-----	40	Very limited Seepage Depth to bedrock Slope	1.00 1.00 0.00	Very limited Thin layer	1.00	Very limited Deep to water	1.00
Ln: Longford-----	100	Somewhat limited Seepage	0.70	Not limited		Very limited Deep to water	1.00
Lo: Longford-----	100	Somewhat limited Seepage	0.05	Somewhat limited Hard to pack	0.70	Very limited Deep to water	1.00
Mc: McCook-----	100	Somewhat limited Seepage	0.70	Very limited Piping Seepage	1.00 0.09	Very limited Deep to water	1.00
Ns: Ness-----	100	Not limited		Very limited Depth to saturated zone	1.00	Very limited Slow refill	1.00

WATER MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Pond Reservoir Area		Embankments, Dikes, and Levees		Excavated Ponds (Aquifer- fed)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Nw: New Cambria-----	100	Somewhat limited Seepage	0.01	Hard to pack Somewhat limited Hard to pack	0.97 0.72	Cutbanks cave Very limited Deep to water	0.10 1.00
Pa: Plevna-----	100	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.20	Very limited Cutbanks cave	1.00
Pr: Pratt-----	100	Very limited Seepage	1.00	Somewhat limited Seepage	0.57	Very limited Deep to water	1.00
Ro: Roxbury-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.40	Very limited Deep to water	1.00
Sm: Smolan-----	100	Not limited		Somewhat limited Hard to pack	0.72	Very limited Deep to water	1.00
To: Tobin-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping	0.60	Very limited Deep to water	1.00
W: Water-----	100	Not rated		Not rated		Not rated	
Wb: Wells-----	90	Somewhat limited Seepage	0.70	Somewhat limited Piping Seepage	0.97 0.06	Very limited Deep to water	1.00
Wc: Wells-----	100	Somewhat limited Seepage	0.70	Somewhat limited Piping Seepage	0.78 0.08	Very limited Deep to water	1.00

SANITARY FACILITIES
McPherson County, Kansas

Sanitary Facilities

The following tables show the degree and kind of soil limitations that affect septic tank absorption fields, sewage lagoons, sanitary landfills, and daily cover for landfill. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Septic tank absorption fields are areas in which effluent from a septic tank is distributed into the soil through subsurface tiles or perforated pipe. Only that part of the soil between depths of 24 and 60 inches is evaluated. The ratings are based on the soil properties that affect absorption of the effluent, construction and maintenance of the system, and public health. Permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, and flooding affect absorption of the effluent. Stones and boulders, ice, and bedrock or a cemented pan interfere with installation. Subsidence interferes with installation and maintenance. Excessive slope may cause lateral seepage and surfacing of the effluent in downslope areas.

Some soils are underlain by loose sand and gravel or fractured bedrock at a depth of less than 4 feet below the distribution lines. In these soils the absorption field may not adequately filter the effluent, particularly when the system is new. As a result, the ground water may become contaminated.

Sewage lagoons are shallow ponds constructed to hold sewage while aerobic bacteria decompose the solid and liquid wastes. Lagoons should have a nearly level floor surrounded by cut slopes or embankments of compacted soil. Nearly impervious soil material for the lagoon floor and sides is required to minimize seepage and contamination of ground water. Considered in the ratings are slope, permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, flooding, large stones, and content of organic matter.

Soil permeability is a critical property affecting the suitability for sewage lagoons. Most porous soils eventually become sealed when they are used as sites for sewage lagoons. Until sealing occurs, however, the hazard of pollution is severe. Soils that have a permeability rate of more than 2 inches per hour are too porous for the proper functioning of sewage lagoons. In these soils, seepage of the effluent can result in contamination of the ground water. Ground-water contamination is also a hazard if fractured bedrock is within a depth of 40 inches, if the water table is high enough to raise the level of sewage in the lagoon, or if floodwater overtops the lagoon.

A high content of organic matter is detrimental to proper functioning of the lagoon because it inhibits aerobic activity. Slope, bedrock, and cemented pans can cause construction problems, and large stones can hinder compaction of the lagoon floor. If the lagoon is to be uniformly deep throughout, the slope must be gentle enough and the soil material must be thick enough over bedrock or a cemented pan to make land smoothing practical.

A trench sanitary landfill is an area where solid waste is placed in successive layers in an excavated trench. The waste is spread, compacted, and covered daily with a thin layer of soil excavated at the site. When the trench is full, a final cover of soil material at least 2 feet thick is placed over the landfill. The ratings in the table are based on the soil properties that affect the risk of pollution, the ease of excavation, trafficability, and revegetation. These properties include permeability, depth to bedrock or a cemented pan, depth to a water table, ponding, slope, flooding, texture, stones and boulders, highly organic layers, soil reaction, and content of salts and sodium. Unless otherwise stated, the ratings apply only to that part of the soil within a depth of about 6 feet. For deeper trenches, onsite investigation may be needed.

Hard, nonrippable bedrock, creviced bedrock, or highly permeable strata in or directly below the proposed trench bottom can affect the ease of excavation and the hazard of ground-water pollution. Slope affects construction of the trenches and the movement of surface water around the landfill. It also affects the construction and performance of roads in areas of the landfill.

Soil texture and consistence affect the ease with which the trench is dug and the ease with which the soil can be used as daily or final cover. They determine the workability of the soil when dry and when wet. Soils that are plastic and sticky when wet are difficult to excavate, grade, or compact and are difficult to place as a uniformly thick cover over a layer of refuse.

The soil material used as the final cover for a trench landfill should be suitable for plants. It should not have excess sodium or salts and should not be too acid. The surface layer generally has the best workability, the highest content of organic matter, and the best potential for plants. Material from the surface layer should be stockpiled for use as the final cover.

SANITARY FACILITIES
McPherson County, Kansas

In an area sanitary landfill, solid waste is placed in successive layers on the surface of the soil. The waste is spread, compacted, and covered daily with a thin layer of soil from a source away from the site. A final cover of soil material at least 2 feet thick is placed over the completed landfill. The ratings in the table are based on the soil properties that affect trafficability and the risk of pollution. These properties include flooding, permeability, depth to a water table, ponding, slope, and depth to bedrock or a cemented pan.

Flooding is a serious problem because it can result in pollution in areas downstream from the landfill. If permeability is too rapid or if fractured bedrock, a fractured cemented pan, or the water table is close to the surface, the leachate can contaminate the water supply. Slope is a consideration because of the extra grading required to maintain roads in the steeper areas of the landfill. Also, leachate may flow along the surface of the soils in the steeper areas and cause difficult seepage problems.

Daily cover for landfill is the soil material that is used to cover compacted solid waste in an area sanitary landfill. The soil material is obtained offsite, transported to the landfill, and spread over the waste. The ratings in the table also apply to the final cover for a landfill. They are based on the soil properties that affect workability, the ease of digging, and the ease of moving and spreading the material over the refuse daily during wet and dry periods. These properties include soil texture, depth to a water table, ponding, rock fragments, slope, depth to bedrock or a cemented pan, reaction, and content of salts, sodium, or lime.

Loamy or silty soils that are free of large stones and excess gravel are the best cover for a landfill. Clayey soils may be sticky and difficult to spread; sandy soils are subject to wind erosion.

Slope affects the ease of excavation and of moving the cover material. Also, it can influence runoff, erosion, and reclamation of the borrow area.

After soil material has been removed, the soil material remaining in the borrow area must be thick enough over bedrock, a cemented pan, or the water table to permit revegetation. The soil material used as the final cover for a landfill should be suitable for plants. It should not have excess sodium, salts, or lime and should not be too acid.

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
053ED: Edalgo-----	100	Very limited Restricted permeability Depth to bedrock	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 0.67
053MB: Mccook-----	100	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
053MC: Mccook-----	100	Somewhat limited Restricted permeability Flooding	0.50 0.40	Somewhat limited Seepage Flooding	0.50 0.40
079GE: Geary-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope Seepage	0.67 0.50
079LB: Ladysmith-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.00
115IC: Irwin-----	90	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.67
159DP: Dillwyn-----	60	Very limited Depth to saturated zone Filtering capacity	1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00
Plevna-----	40	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
159GC: Geary-----	75	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope Seepage	0.67 0.50
Clark-----	25	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope Seepage	0.67 0.50
159LA: Lancaster-----	90	Very limited Depth to bedrock Restricted permeability	1.00 0.50	Very limited Depth to soft bedrock Seepage Slope	1.00 0.50 0.00
159NF: Naron-----	100	Somewhat limited Restricted permeability	0.50	Very limited Seepage Slope	1.00 0.00
159PF: Pratt-----	100	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.09
159PR: Pratt-----	65	Very limited Filtering capacity	1.00	Very limited Seepage Slope	1.00 0.09
Carwile-----	35	Very limited Restricted permeability Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 0.32
159SO: Smolan-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.67

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
159WA: Waldeck-----	100	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
169CE: Clime-----	100	Very limited Restricted permeability Depth to bedrock	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 0.33
169ED: Edalgo-----	100	Very limited Restricted permeability Depth to bedrock	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 0.67
169RO: Roxbury-----	99	Somewhat limited Restricted permeability Flooding	0.50 0.40	Somewhat limited Seepage Flooding	0.50 0.40
1191: Blazefork-----	90	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 1.00 0.40	Somewhat limited Depth to saturated zone Flooding	0.71 0.40
1200: Buhler-----	65	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 0.43 0.40	Somewhat limited Seepage Flooding	0.50 0.40
Blazefork-----	30	Very limited Restricted permeability Depth to saturated zone Flooding	1.00 1.00 0.40	Somewhat limited Depth to saturated zone Flooding	0.71 0.40
1324: Carway-----	50	Very limited Restricted permeability Ponding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Seepage	1.00 0.50
Carbika-----	30	Very limited Ponding Depth to saturated zone Restricted permeability	1.00 1.00 0.50	Very limited Ponding Seepage	1.00 0.50
1553: Darlow-----	70	Very limited Restricted permeability	1.00	Not limited	
Elmer-----	20	Very limited Restricted permeability	1.00	Very limited Seepage	1.00
1555: Dillhut-----	35	Very limited Filtering capacity Restricted permeability	1.00 0.50	Very limited Seepage Slope	1.00 0.00
Plev-----	35	Very limited Depth to saturated zone Filtering capacity Restricted permeability	1.00 1.00 0.50	Very limited Seepage	1.00
1985: Hayes-----	60	Very limited Restricted permeability	1.00	Very limited Seepage Slope	1.00 0.09

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
2391: Kaskan-----	75	Very limited Flooding Filtering capacity Depth to saturated zone	1.00 1.00 0.43	Very limited Flooding Seepage	1.00 1.00
2588: Longford, Moderately Eroded-----	90	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.33
3181: Pratt-----	45	Very limited Filtering capacity	1.00	Very limited Seepage	1.00
Turon-----	30	Very limited Restricted permeability Filtering capacity	1.00 1.00	Slope Very limited Seepage	0.09 1.00
3641: Tivin-----	45	Very limited Filtering capacity Slope	1.00 0.16	Very limited Seepage Slope	1.00 1.00
Dillhut-----	40	Very limited Filtering capacity Restricted permeability	1.00 0.50	Very limited Seepage Slope	1.00 0.09
AED: Arents, Earthen Dam-	100	Not rated		Not rated	
At: Attica-----	100	Not limited		Very limited Seepage Slope	1.00 0.09
Br: Bridgeport-----	100	Somewhat limited Restricted permeability Flooding	0.50 0.40	Somewhat limited Seepage Flooding	0.50 0.40
Ca: Carwile-----	100	Very limited Restricted permeability Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 0.32
Cb: Cass-----	100	Very limited Filtering capacity Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40
Ce: Clime-----	100	Very limited Restricted permeability Depth to bedrock	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 0.00
Cm: Clime-----	100	Very limited Restricted permeability Depth to bedrock	1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 0.67
Cr: Crete-----	100	Very limited Restricted permeability	1.00	Somewhat limited Seepage	0.32
Cs: Crete-----	100	Very limited Restricted permeability	1.00	Somewhat limited Seepage Slope	0.32 0.00
Ct: Crete-----	100	Very limited Restricted permeability	1.00	Somewhat limited Seepage Slope	0.32 0.00

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
De: Detroit-----	100	Very limited Restricted permeability Flooding	1.00 0.40	Somewhat limited Flooding	0.40
Dr: Drummond-----	100	Very limited Restricted permeability Depth to saturated zone	1.00 1.00	Somewhat limited Depth to saturated zone	0.71
Ed: Edalgo-----	100	Very limited Restricted permeability Depth to bedrock Slope	1.00 1.00 0.04	Very limited Depth to soft bedrock Slope	1.00 1.00
Ee: Edalgo-----	100	Very limited Depth to bedrock	1.00	Very limited Depth to soft bedrock Slope	1.00 0.91
Fa: Farnum-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.00
Ge: Geary-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage Slope	0.50 0.00
Go: Goessel-----	100	Very limited Restricted permeability Depth to saturated zone	1.00 1.00	Somewhat limited Depth to saturated zone	0.81
Ho: Hord-----	100	Somewhat limited Restricted permeability Flooding	0.50 0.40	Somewhat limited Seepage Flooding	0.50 0.40
INT: Aquolls-----	100	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
Ir: Irwin, bedrock substratum-----	100	Very limited Restricted permeability Depth to bedrock	1.00 0.78	Somewhat limited Depth to soft bedrock Slope	0.42 0.00
La: Ladysmith-----	100	Very limited Restricted permeability	1.00	Not limited	
Ld: Ladysmith-----	70	Very limited Restricted permeability	1.00	Not limited	
Drummond-----	30	Very limited Restricted permeability Depth to saturated zone	1.00 1.00	Somewhat limited Depth to saturated zone	0.71
Le: Lancaster-----	100	Very limited Depth to bedrock Restricted permeability	1.00 0.50	Very limited Depth to soft bedrock Seepage Slope	1.00 0.50 0.33
Lh: Lancaster-----	60	Very limited Depth to bedrock Restricted permeability	1.00 0.50	Very limited Depth to soft bedrock Slope	1.00 1.00

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
Hedville-----	40	Slope Very limited Depth to bedrock	0.04 1.00	Seepage Very limited Depth to hard bedrock	0.50 1.00
LHH: Lancaster-----	60	Slope Very limited Depth to bedrock	0.04 1.00	Slope Very limited Depth to soft bedrock	1.00 1.00
		Restricted permeability Slope	0.50 0.00	Slope Seepage	1.00 0.50
Hedville-----	40	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	0.63	Slope Seepage	1.00 0.50
Ln: Longford-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.67
				Seepage	0.50
Lo: Longford-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.33
Mc: Mccook-----	100	Somewhat limited Restricted permeability Flooding	0.50 0.40	Somewhat limited Seepage	0.50
				Flooding	0.40
Ns: Ness-----	100	Very limited Restricted permeability Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone	1.00
Nw: New Cambria-----	100	Very limited Restricted permeability Flooding	1.00 0.40	Somewhat limited Flooding	0.40
Pa: Plevna-----	100	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Seepage	1.00 1.00
				Depth to saturated zone	1.00
Pr: Pratt-----	100	Very limited Filtering capacity Slope	1.00 0.04	Very limited Seepage	1.00
				Slope	1.00
Ro: Roxbury-----	100	Somewhat limited Restricted permeability Flooding	0.50 0.40	Somewhat limited Seepage	0.50
				Flooding	0.40
Sm: Smolan-----	100	Very limited Restricted permeability	1.00	Somewhat limited Slope	0.00
To: Tobin-----	100	Very limited Flooding Restricted permeability	1.00 0.50	Very limited Flooding Seepage	1.00 0.50
W: Water-----	100	Not rated		Not rated	
Wb: Wells-----	90	Somewhat limited Restricted permeability	0.50	Somewhat limited Seepage	0.50
				Slope	0.00
Wc: Wells-----	100	Somewhat limited Restricted permeability	0.50	Somewhat limited Slope	0.67

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
				Seepage	0.50

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
053ED: Edalgo-----	100	Very limited Depth to bedrock Too clayey Seepage	1.00 1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Too clayey Hard to compact	1.00 1.00 1.00
053MB: Mccook-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Not limited	
053MC: Mccook-----	100	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Not limited	
079GE: Geary-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
079LB: Ladysmith-----	100	Very limited Too clayey	1.00	Not limited		Very limited Too clayey Hard to compact	1.00 1.00
115IC: Irwin-----	90	Very limited Too clayey	1.00	Not limited		Very limited Too clayey Hard to compact	1.00 1.00
159DP: Dillwyn-----	60	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Seepage	1.00
Plevna-----	40	Too Sandy	1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Depth to saturated zone Too Sandy	0.86 0.50
		Very limited Flooding	1.00			Very limited Too Sandy	1.00
		Depth to saturated zone Seepage	1.00 1.00			Depth to saturated zone Seepage	1.00 0.50
		Too Sandy	1.00				
159GC: Geary-----	75	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Clark-----	25	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
159LA: Lancaster-----	90	Very limited Depth to bedrock Seepage	1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock	1.00
159NF: Naron-----	100	Very limited Seepage	1.00	Not limited		Not limited	
159PF: Pratt-----	100	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Seepage Too Sandy	1.00 0.50
159PR: Pratt-----	65	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Seepage Too Sandy	1.00 0.50
Carwile-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Too clayey	0.50			Hard to compact Too clayey	1.00 0.50
159SO: Smolan-----	100	Very limited Too clayey	1.00	Not limited		Very limited Too clayey Hard to compact	1.00 1.00
159WA: Waldeck-----	100	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Too Sandy	1.00
		Depth to saturated zone	1.00			Seepage	1.00
		Seepage	1.00			Depth to saturated zone	0.09
		Too Sandy	1.00				
169CE: Clime-----	100	Very limited Depth to bedrock Too clayey Seepage	1.00 1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Too clayey Hard to compact	1.00 1.00 1.00
169ED: Edalgo-----	100	Very limited Depth to bedrock Too clayey Seepage	1.00 1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Too clayey Hard to compact	1.00 1.00 1.00

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
169RO: Roxbury-----	99	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Not limited	
1191: Blazefork-----	90	Very limited Depth to saturated zone Too clayey Flooding	1.00 0.50 0.40	Very limited Depth to saturated zone Flooding	1.00 0.40	Somewhat limited Too clayey	0.50
1200: Buhler-----	65	Very limited Depth to saturated zone Sodium content Flooding	1.00 1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Sodium content Too clayey	1.00 0.50
Blazefork-----	30	Very limited Depth to saturated zone Too clayey Flooding	1.00 0.50 0.40	Very limited Depth to saturated zone Flooding	1.00 0.40	Somewhat limited Too clayey	0.50
1324: Carway-----	50	Very limited Depth to saturated zone Ponding Too clayey	1.00 1.00 0.50	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Hard to compact Too clayey	1.00 1.00 1.00 0.50
Carbika-----	30	Very limited Depth to saturated zone Ponding Too clayey	1.00 1.00 0.50	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Too clayey	1.00 1.00 0.50
1553: Darlow-----	70	Very limited Sodium content	1.00	Not limited		Very limited Sodium content	1.00
Elmer-----	20	Very limited Sodium content Seepage	1.00 1.00	Not limited		Very limited Sodium content Seepage	1.00 0.16
1555: Dillhut-----	35	Very limited Seepage	1.00	Very limited Seepage	1.00	Not limited	
Plev-----	35	Very limited Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Depth to saturated zone Too Sandy Seepage	1.00 1.00 1.00
1985: Hayes-----	60	Very limited Too clayey	1.00	Very limited Seepage	1.00	Very limited Hard to compact	1.00
2391: Kaskan-----	75	Very limited Flooding Depth to saturated zone Too Sandy Seepage	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Too Sandy Seepage	1.00 1.00
2588: Longford, Moderately Eroded-----	90	Very limited Too clayey	1.00	Not limited		Very limited Too clayey	1.00
3181: Pratt-----	45	Very limited Seepage Too Sandy	1.00 1.00	Very limited Seepage	1.00	Very limited Too Sandy Seepage	1.00 1.00
Turon-----	30	Very limited Too Sandy	1.00	Very limited Seepage	1.00	Very limited Seepage Too Sandy	1.00 0.50
3641: Tivin-----	45	Very limited Seepage Too Sandy Slope	1.00 1.00 0.16	Very limited Seepage Slope	1.00 0.16	Very limited Too Sandy Seepage Slope	1.00 1.00 0.16
Dillhut-----	40	Very limited Seepage	1.00	Very limited Seepage	1.00	Not limited	
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
At: Attica-----	100	Very limited Seepage	1.00	Very limited Seepage	1.00	Somewhat limited Seepage	0.50

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Br: Bridgeport-----	100	Too Sandy	1.00			Too Sandy	0.50
Ca: Carwile-----	100	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Not limited	
Cb: Cass-----	100	Very limited Depth to saturated zone Too clayey	1.00 0.50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Hard to compact Too clayey	1.00 1.00 0.50
Ce: Clime-----	100	Very limited Seepage Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40	Somewhat limited Seepage	0.50
Cm: Clime-----	100	Very limited Depth to bedrock Too clayey Seepage	1.00 1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Too clayey Hard to compact	1.00 1.00 1.00
Cr: Crete-----	100	Very limited Depth to bedrock Too clayey Seepage	1.00 1.00 1.00	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Too clayey Hard to compact	1.00 1.00 1.00
Cs: Crete-----	100	Somewhat limited Too clayey	0.50	Not limited		Very limited Too clayey Hard to compact	1.00 1.00
Ct: Crete-----	100	Very limited Too clayey	1.00	Not limited		Very limited Too clayey Hard to compact	1.00 1.00
De: Detroit-----	100	Somewhat limited Too clayey	0.50	Not limited		Very limited Hard to compact Too clayey	1.00 0.50
Dr: Drummond-----	100	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Not limited	
Ed: Edalgo-----	100	Very limited Depth to saturated zone Too clayey	1.00 0.50	Very limited Depth to saturated zone	1.00	Very limited Hard to compact	1.00
Ee: Edalgo-----	100	Very limited Depth to bedrock Seepage Too clayey Slope	1.00 1.00 0.50 0.04	Very limited Depth to bedrock Slope	1.00 0.04	Too clayey	0.50
Fa: Farnum-----	100	Very limited Depth to bedrock Seepage Too clayey	1.00 1.00 0.50	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Hard to compact Too clayey	1.00 1.00 0.50
Ge: Geary-----	100	Not limited		Not limited		Not limited	
Go: Goessel-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Ho: Hord-----	100	Very limited Too clayey Depth to saturated zone	1.00 0.86	Somewhat limited Depth to saturated zone	0.19	Very limited Too clayey	1.00
INT: Aquolls-----	100	Depth to saturated zone	0.47			Hard to compact	1.00
Ir: Irwin, bedrock substratum-----	100	Somewhat limited Too clayey Flooding	0.50 0.40	Somewhat limited Flooding	0.40	Depth to saturated zone	0.47
		Somewhat limited Too clayey	0.50			Somewhat limited Too clayey	0.50
		Very limited Depth to saturated zone Ponding Seepage	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Somewhat limited Too clayey	0.50
		Very limited Depth to bedrock	1.00	Somewhat limited Depth to bedrock	0.42	Very limited Too clayey	1.00

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
La: Ladysmith-----	100	Too clayey Seepage	1.00 1.00	Not limited		Hard to compact Depth to bedrock	1.00 0.42
		Very limited Too clayey	1.00			Very limited Too clayey Hard to compact	1.00 1.00
Ld: Ladysmith-----	70	Very limited Too clayey	1.00	Not limited		Very limited Too clayey Hard to compact	1.00 1.00
		Very limited Depth to saturated zone Too clayey	1.00 1.00 0.50			Very limited Too clayey Hard to compact	1.00 1.00 1.00
Drummond-----	30	Very limited Depth to saturated zone Too clayey	1.00 1.00 0.50	Very limited Depth to saturated zone	1.00	Very limited Hard to compact	1.00
Le: Lancaster-----	100	Very limited Depth to bedrock Seepage Too clayey	1.00 1.00 1.00 0.50	Very limited Depth to bedrock	1.00	Very limited Depth to bedrock Too clayey	1.00 0.50
Lh: Lancaster-----	60	Very limited Depth to bedrock Seepage Too clayey Slope	1.00 1.00 1.00 0.50 0.04	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Too clayey Slope	1.00 0.50 0.04
Hedville-----	40	Very limited Depth to bedrock Seepage Slope	1.00 1.00 1.00 0.04	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Slope	1.00 0.04
LHH: Lancaster-----	60	Very limited Depth to bedrock Seepage Too clayey Slope	1.00 1.00 1.00 0.50 0.00	Very limited Depth to bedrock Slope	1.00 0.00	Very limited Depth to bedrock Too clayey Slope	1.00 0.50 0.00
Hedville-----	40	Very limited Depth to bedrock Seepage Slope	1.00 1.00 1.00 0.63	Very limited Depth to bedrock Slope	1.00 0.63	Very limited Depth to bedrock Slope	1.00 0.63
Ln: Longford-----	100	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
Lo: Longford-----	100	Very limited Too clayey	1.00	Not limited		Very limited Too clayey Hard to compact	1.00 1.00
Mc: Mccook-----	100	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Not limited	
Ns: Ness-----	100	Very limited Depth to saturated zone Too clayey	1.00 1.00 1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Too clayey Hard to compact	1.00 1.00 1.00 1.00
Nw: New Cambria-----	100	Very limited Too clayey Flooding	1.00 1.00 0.40	Somewhat limited Flooding	0.40	Very limited Too clayey Hard to compact	1.00 1.00 1.00
Pa: Plevna-----	100	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Depth to saturated zone Seepage	1.00 0.50
Pr: Pratt-----	100	Depth to saturated zone Seepage Too Sandy	1.00 1.00 1.00 1.00	Depth to saturated zone Seepage	1.00 1.00	Too Sandy	0.50
Pr: Pratt-----	100	Very limited Seepage Too Sandy Slope	1.00 1.00 1.00 0.04	Very limited Seepage Slope	1.00 0.04	Very limited Seepage Too Sandy Slope	1.00 0.50 0.04
Ro: Roxbury-----	100	Somewhat limited Too clayey Flooding	0.50 0.40	Somewhat limited Flooding	0.40	Somewhat limited Too clayey	0.50
Sm: Smolan-----	100	Somewhat limited Too clayey	0.50	Not limited		Very limited Hard to compact Too clayey	1.00 0.50
To: Tobin-----	100	Very limited		Very limited		Not limited	

SANITARY FACILITIES--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
		Flooding	1.00	Flooding	1.00		
W: Water-----	100	Not rated		Not rated		Not rated	
Wb: Wells-----	90	Not limited		Not limited		Not limited	
Wc: Wells-----	100	Not limited		Not limited		Not limited	

AGRICULTURAL WASTE MANAGEMENT
McPherson County, Kansas

The nature of the soil is also important in the application of organic wastes and wastewater to land as fertilizers and irrigation; it is also important when the soil is used as a medium for treatment and disposal of these wastes. Favorable soil properties are required to prevent environmental damage.

The use of organic wastes and wastewater as production resources will result in energy conservation, prevent the waste of these important resources, and prevent problems associated with their disposal. Where disposal is the goal, and a maximum amount is disposed in a minimum area to hold costs to a minimum, risk of environmental damage is the principal constraint. Where the reuse goal is pursued, and a minimum amount is applied to a maximum area to obtain the greatest benefit, environmental damage is unlikely.

Interpretations developed for waste management may include ratings for (1) manure and food processing wastes; (2) municipal sewage sludge; (3) irrigation use of wastewater; or (4) treatment of wastewater by the slow rate process, overland flow process, or rapid infiltration process. If available, these should be located in this subsection.

Soil properties are important considerations in areas where soils are used as sites for the treatment and disposal of organic waste and wastewater. Selection of soils with properties that favor waste management can help to prevent environmental damage.

The Ag-Waste tables show the degree and kind of soil limitations affecting the treatment of agricultural waste, including municipal and food-processing wastewater and effluent from lagoons or storage ponds. Municipal wastewater is the waste stream from a municipality. It contains domestic waste and may contain industrial waste. It may have received primary or secondary treatment. It is rarely untreated sewage. Food-processing wastewater results from the preparation of fruits, vegetables, milk, cheese, and meats for public consumption. In places it is high in content of sodium and chloride. In the context of these tables, the effluent in lagoons and storage ponds is from facilities used to treat or store food-processing wastewater or domestic or animal waste. Domestic and food-processing wastewater is very dilute, and the effluent from the facilities that treat or store it commonly is very low in content of carbonaceous and nitrogenous material; the content of nitrogen commonly ranges from 10 to 30 milligrams per liter. The wastewater from animal waste treatment lagoons or storage ponds, however, has much higher concentrations of these materials, mainly because the manure has not been diluted as much as the domestic waste. The content of nitrogen in this wastewater generally ranges from 50 to 2,000 milligrams per liter. When wastewater is applied, checks should be made to ensure that nitrogen, phosphorus, heavy metals, and salts are not added in excessive amounts.

The ratings in the tables are for waste management systems that not only dispose of and treat organic waste or wastewater but also are beneficial to crops (application of manure and food-processing waste, application of sewage sludge, and disposal of wastewater by irrigation) and for waste management systems that are designed only for the purpose of wastewater disposal and treatment (overland flow of wastewater, rapid infiltration of wastewater, and slow rate treatment of wastewater).

The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect agricultural waste management. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are generally favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Application of manure and food-processing waste not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. Manure is the excrement of livestock and poultry, and food-processing waste is damaged fruit and vegetables and the peelings, stems, leaves, pits, and soil particles removed in food preparation. The manure and food-processing waste are either solid, slurry, or liquid. Their nitrogen content varies. A high content of nitrogen limits the application rate. Toxic or otherwise dangerous wastes, such as those mixed with the lye used in food processing, are not considered in the ratings.

The ratings are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the waste is applied, and the method by which the waste is applied. The properties that affect absorption include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, and available water capacity. The properties that affect plant growth and microbial activity include reaction, the sodium adsorption ratio, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of waste. Permanently frozen soils are unsuitable for waste treatment.

Application of sewage sludge not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. In the context of this table, sewage sludge is the residual product of the treatment of municipal sewage. The solid component consists mainly of cell mass, primarily bacteria cells that developed during secondary treatment and have incorporated soluble organics into their own bodies. The sludge has small amounts of sand, silt, and other solid debris. The content of nitrogen varies. Some sludge has constituents that are toxic to plants or hazardous to the food chain, such as heavy metals and exotic organic compounds, and should be analyzed chemically prior to use.

AGRICULTURAL WASTE MANAGEMENT
McPherson County, Kansas

The content of water in the sludge ranges from about 98 percent to less than 40 percent. The sludge is considered liquid if it is more than about 90 percent water, slurry if it is about 50 to 90 percent water, and solid if it is less than about 50 percent water.

The ratings in the table are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the sludge is applied, and the method by which the sludge is applied. The properties that affect absorption, plant growth, and microbial activity include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, available water capacity, reaction, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of sludge. Permanently frozen soils are unsuitable for waste treatment.

Disposal of wastewater by irrigation not only disposes of municipal wastewater and wastewater from food-processing plants, lagoons, and storage ponds but also can improve crop production by increasing the amount of water available to crops. The ratings in the table are based on the soil properties that affect the design, construction, management, and performance of the irrigation system. The properties that affect design and management include the sodium adsorption ratio, depth to a water table, ponding, available water capacity, permeability, slope, and flooding. The properties that affect construction include stones, cobbles, depth to bedrock or a cemented pan, depth to a water table, and ponding.

The properties that affect performance include depth to bedrock or a cemented pan, bulk density, the sodium adsorption ratio, salinity, reaction, and the cation-exchange capacity, which is used to estimate the capacity of a soil to adsorb heavy metals. Permanently frozen soils are not suitable for disposal of wastewater by irrigation.

See the National Soil Handbook, September 1992, Part 620, for criteria used in rating soils for sanitary facilities and waste management.

AGRICULTURAL WASTE MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
053ED: Edalgo-----	100	Very limited Restricted permeability Depth to bedrock Droughty Too acid	1.00 0.16 0.11 0.11	Very limited Restricted permeability Too acid Depth to bedrock Droughty	1.00 0.42 0.16 0.11	Very limited Restricted permeability Too acid Too steep for surface application Depth to bedrock Droughty	1.00 0.42 0.31 0.16 0.11
053MB: Mccook-----	100	Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Somewhat limited Flooding	0.60
053MC: Mccook-----	100	Not limited		Somewhat limited Flooding	0.40	Not limited	
079GE: Geary-----	100	Somewhat limited Too acid	0.03	Somewhat limited Too acid	0.14	Somewhat limited Too steep for surface application Too acid	0.31 0.14
079LB: Ladysmith-----	100	Very limited Restricted permeability Runoff limitation	1.00 0.40	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
115IC: Irwin-----	90	Very limited Restricted permeability Runoff limitation	1.00 0.40	Very limited Restricted permeability	1.00	Very limited Restricted permeability Too steep for surface application	1.00 0.31
159DP: Dillwyn-----	60	Very limited Filtering capacity Depth to saturated zone Leaching limitation Droughty	1.00 1.00 0.45 0.20	Very limited Filtering capacity Depth to saturated zone Droughty	1.00 1.00 0.20	Very limited Filtering capacity Depth to saturated zone Droughty	1.00 1.00 0.20
Plevna-----	40	Very limited Flooding Depth to saturated zone Runoff limitation Filtering capacity	1.00 1.00 0.40 0.00	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 0.00	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 0.00
159GC: Geary-----	75	Somewhat limited Too acid	0.03	Somewhat limited Too acid	0.14	Somewhat limited Too steep for surface application Too acid	0.31 0.14
Clark-----	25	Not limited		Not limited		Somewhat limited Too steep for surface application	0.31
159LA: Lancaster-----	90	Somewhat limited Depth to bedrock Too acid	0.10 0.03	Somewhat limited Too acid Depth to bedrock	0.14 0.10	Somewhat limited Too acid Depth to bedrock	0.14 0.10
159NF: Naron-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00
159PF: Pratt-----	100	Very limited Filtering capacity Leaching limitation	1.00 0.45	Very limited Filtering capacity	1.00	Very limited Filtering capacity Too steep for surface application	1.00 0.00
159PR: Pratt-----	65	Very limited		Very limited		Very limited	

AGRICULTURAL WASTE MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Carwile-----	35	Filtering capacity	1.00	Filtering capacity	1.00	Filtering capacity	1.00
		Leaching limitation	0.45			Too steep for surface application	0.00
		Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Restricted permeability	1.00	Restricted permeability	1.00	Restricted permeability	1.00
159SO: Smolan-----	100	Runoff limitation	0.40	Too acid	0.07	Too acid	0.07
		Too acid	0.02				
159WA: Waldeck-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability Too steep for surface application	1.00
							0.31
169CE: Clime-----	100	Very limited Filtering capacity Flooding	1.00	Very limited Flooding	1.00	Very limited Filtering capacity Flooding	1.00
		Depth to saturated zone	0.60	Filtering capacity	1.00	Depth to saturated zone	0.60
			0.43	Depth to saturated zone	0.43		0.43
169ED: Edalgo-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Depth to bedrock	0.65	Depth to bedrock	0.65	Depth to bedrock	0.65
		Droughty	0.45	Droughty	0.45	Droughty	0.45
						Too steep for surface application	0.08
169RO: Roxbury-----	99	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Depth to bedrock	0.42	Depth to bedrock	0.42	Depth to bedrock	0.42
		Droughty	0.28	Too acid	0.42	Too acid	0.42
		Too acid	0.11	Droughty	0.28	Too steep for surface application Droughty	0.31
1191: Blazefork-----	90	Not limited		Somewhat limited Flooding	0.40	Not limited	
1200: Buhler-----	65	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Sodium content	1.00	Sodium content	1.00	Sodium content	1.00
		Runoff limitation	0.40	Flooding	0.40	Too acid	0.31
		Too acid	0.08	Too acid	0.31	Salinity	0.13
Blazefork-----	30	Salinity	0.06	Salinity	0.13		
		Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Too acid	0.50	Too acid	1.00	Too acid	1.00
1324: Carway-----	50	Runoff limitation	0.40	Flooding	0.40		
		Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Ponding	1.00	Ponding	1.00	Ponding	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
		Runoff limitation	0.40	Too acid	0.14	Too acid	0.14
		Too acid	0.03	Filtering capacity	0.00	Filtering capacity	0.00

AGRICULTURAL WASTE MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Carbika-----	30	Very limited Restricted permeability Ponding Depth to saturated zone Runoff limitation Too acid	1.00 1.00 1.00 0.40 0.03	Very limited Restricted permeability Ponding Depth to saturated zone Too acid	1.00 1.00 1.00 0.14	Very limited Restricted permeability Ponding Depth to saturated zone Too acid	1.00 1.00 1.00 0.14
1553: Darlow-----	70	Very limited Restricted permeability Sodium content Too acid Salinity	1.00 1.00 0.62 0.01	Very limited Restricted permeability Sodium content Too acid	1.00 1.00 1.00	Very limited Restricted permeability Sodium content Too acid	1.00 1.00 1.00
Elmer-----	20	Very limited Restricted permeability Depth to dense layer Too acid Sodium content Filtering capacity	1.00 1.00 0.68 0.32 0.00	Very limited Restricted permeability Too acid Sodium content Filtering capacity	1.00 1.00 0.32 0.00	Very limited Restricted permeability Too acid Sodium content Filtering capacity	1.00 1.00 0.32 0.00
1555: Dillhut-----	35	Very limited Filtering capacity Too acid	1.00 0.00	Very limited Filtering capacity Too acid	1.00 0.01	Very limited Filtering capacity Too acid	1.00 0.01
Plev-----	35	Very limited Filtering capacity Depth to saturated zone Too acid Droughty	1.00 1.00 0.03 0.01	Very limited Filtering capacity Depth to saturated zone Too acid Droughty	1.00 1.00 0.14 0.01	Very limited Filtering capacity Depth to saturated zone Too acid Droughty	1.00 1.00 0.14 0.01
1985: Hayes-----	60	Very limited Restricted permeability Too acid Filtering capacity	1.00 0.02 0.00	Very limited Restricted permeability Too acid Filtering capacity	1.00 0.07 0.00	Very limited Restricted permeability Too acid Filtering capacity Too steep for surface application	1.00 0.07 0.00 0.00
2391: Kaskan-----	75	Very limited Flooding Filtering capacity Restricted permeability	1.00 1.00 0.30	Very limited Flooding Filtering capacity Restricted permeability	1.00 1.00 0.22	Very limited Flooding Filtering capacity Restricted permeability	1.00 1.00 0.22
2588: Longford, Moderately Eroded-----	90	Somewhat limited Restricted permeability	0.89	Somewhat limited Restricted permeability	0.78	Somewhat limited Restricted permeability Too steep for surface application	0.78 0.08
3181: Pratt-----	45	Very limited Filtering capacity Low adsorption Leaching limitation Too acid	1.00 1.00 0.45 0.11	Very limited Filtering capacity Too acid	1.00 0.42	Very limited Filtering capacity Low adsorption Too acid Too steep for surface application	1.00 1.00 0.42 0.00
Turon-----	30	Very limited Filtering capacity Restricted permeability	1.00 0.89	Very limited Filtering capacity Restricted permeability	1.00 0.78	Very limited Filtering capacity Restricted permeability	1.00 0.78

AGRICULTURAL WASTE MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
3641: Tivin-----	45	Leaching limitation	0.45	Too acid	0.21	Too acid	0.21
		Too acid	0.05			Too steep for surface application	0.00
		Very limited Filtering capacity	1.00			Very limited Filtering capacity	1.00
		Droughty	0.64			Too steep for surface application	1.00
		Leaching limitation	0.45			Droughty	0.64
Dillhut-----	40	Slope	0.16	Slope Too acid	0.16 0.01	Too steep for sprinkler application	0.39
		Too acid	0.00			Too acid	0.01
		Very limited Filtering capacity	1.00			Very limited Filtering capacity	1.00
		Too acid	0.00			Too acid	0.01
						Too steep for surface application	0.00
AED: Arents, Earthen Dam-	100	Not rated		Not rated		Not rated	
At: Attica-----	100	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity	0.00	Somewhat limited Filtering capacity Too steep for surface application	0.00 0.00
Br: Bridgeport-----	100	Not limited		Somewhat limited Flooding	0.40	Not limited	
Ca: Carwile-----	100	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Cb: Cass-----	100	Restricted permeability	1.00	Restricted permeability	1.00	Restricted permeability	1.00
		Runoff limitation	0.40	Too acid	0.07	Too acid	0.07
		Too acid	0.02				
Ce: Clime-----	100	Very limited Filtering capacity	1.00	Very limited Filtering capacity Flooding	1.00 0.40	Very limited Filtering capacity	1.00
Cm: Clime-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Droughty	0.80	Droughty	0.80	Droughty	0.80
		Depth to bedrock	0.71	Depth to bedrock	0.71	Depth to bedrock	0.71
Cr: Crete-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
		Droughty	0.80	Droughty	0.80	Droughty	0.80
		Depth to bedrock	0.71	Depth to bedrock	0.71	Depth to bedrock	0.71
Cs: Crete-----	100	Too steep for surface application	0.31	Too steep for surface application	0.31	Too steep for surface application	0.31
		Too acid	0.11	Too acid	0.42	Too acid	0.42
Cs: Crete-----	100	Somewhat limited Restricted permeability	0.89	Somewhat limited Restricted permeability	0.78	Somewhat limited Restricted permeability	0.78
		Too acid	0.11	Too acid	0.42	Too acid	0.42

AGRICULTURAL WASTE MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Ct: Crete-----	100	Somewhat limited Restricted permeability Too acid	0.89 0.11	Somewhat limited Restricted permeability Too acid	0.78 0.42	Somewhat limited Restricted permeability Too acid	0.78 0.42
De: Detroit-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability Flooding	1.00 0.40	Very limited Restricted permeability	1.00
Dr: Drummond-----	100	Very limited Restricted permeability Runoff limitation Salinity	1.00 0.40 0.01	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
Ed: Edalgo-----	100	Very limited Restricted permeability Depth to bedrock	1.00 0.42	Very limited Restricted permeability Depth to bedrock	1.00 0.42	Very limited Restricted permeability Too steep for surface application Depth to bedrock Too acid Too steep for sprinkler application	1.00 1.00 0.42 0.42 0.22
Ee: Edalgo-----	100	Droughty Too acid Slope	0.15 0.11 0.04	Too acid Droughty Slope	0.42 0.15 0.04		
Ee: Edalgo-----	100	Very limited Restricted permeability Depth to bedrock Droughty Too acid	1.00 0.95 0.83 0.11	Very limited Restricted permeability Depth to bedrock Droughty Too acid	1.00 0.95 0.83 0.42	Very limited Restricted permeability Depth to bedrock Droughty Too steep for surface application Too acid	1.00 0.95 0.83 0.66 0.42
Fa: Farnum-----	100	Not limited		Not limited		Not limited	
Ge: Geary-----	100	Somewhat limited Too acid	0.03	Somewhat limited Too acid	0.14	Somewhat limited Too acid	0.14
Go: Goessel-----	100	Very limited Restricted permeability Depth to saturated zone Runoff limitation	1.00 0.86 0.40	Very limited Restricted permeability Depth to saturated zone	1.00 0.86	Very limited Restricted permeability Depth to saturated zone	1.00 0.86
Ho: Hord-----	100	Not limited		Somewhat limited Flooding	0.40	Not limited	
INT: Aquolls-----	100	Very limited Depth to saturated zone Low adsorption Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Low adsorption Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Low adsorption Ponding	1.00 1.00 1.00
Ir: Irwin, bedrock substratum-----	100	Very limited Restricted permeability Runoff limitation	 1.00 0.40	Very limited Restricted permeability	 1.00	Very limited Restricted permeability	 1.00
La: Ladysmith-----	100	Very limited Restricted permeability Runoff limitation	1.00 0.40	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
Ld: Ladysmith-----	70	Very limited Restricted permeability Runoff limitation	1.00 0.40	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
Drummond-----	30	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00

AGRICULTURAL WASTE MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Le: Lancaster-----	100	Runoff limitation	0.40				
		Salinity	0.01				
		Somewhat limited		Somewhat limited		Somewhat limited	
		Depth to bedrock	0.29	Depth to bedrock	0.29	Depth to bedrock	0.29
Lh: Lancaster-----	60	Too acid	0.03	Too acid	0.14	Too acid	0.14
		Droughty	0.01	Droughty	0.01	Too steep for surface application	0.08
						Droughty	0.01
Hedville-----	40	Somewhat limited	0.29	Somewhat limited	0.29	Very limited	1.00
		Depth to bedrock		Depth to bedrock		Too steep for surface application	
		Slope	0.04	Too acid	0.14	Depth to bedrock	0.29
		Too acid	0.03	Slope	0.04	Too steep for sprinkler application	0.22
LHH: Lancaster-----	60	Droughty	0.01	Droughty	0.01	Too acid	0.14
		Very limited		Very limited		Droughty	0.01
		Depth to bedrock	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Droughty	1.00	Droughty	1.00	Droughty	1.00
Hedville-----	40	Runoff limitation	0.40	Slope	0.04	Too steep for surface application	1.00
						Too steep for sprinkler application	0.22
		Slope	0.04				
Ln: Longford-----	100	Somewhat limited	0.46	Somewhat limited	0.46	Very limited	1.00
		Depth to bedrock		Depth to bedrock		Too steep for surface application	
		Droughty	0.07	Too acid	0.14	Depth to bedrock	0.46
		Too acid	0.03	Droughty	0.07	Too acid	0.14
Hedville-----	40	Slope	0.00	Slope	0.00	Too steep for sprinkler application	0.10
		Very limited		Very limited		Droughty	0.07
		Depth to bedrock	1.00	Depth to bedrock	1.00	Depth to bedrock	1.00
		Droughty	1.00	Droughty	1.00	Droughty	1.00
Lo: Longford-----	100	Slope	0.63	Slope	0.63	Too steep for surface application	1.00
		Runoff limitation	0.40			Too steep for sprinkler application	0.77
Mc: Mccook-----	100	Very limited	1.00	Very limited	1.00	Very limited	1.00
		Restricted		Restricted		Restricted	
		permeability		permeability		permeability	
						Too steep for surface application	0.31
Ns: Ness-----	100	Somewhat limited	0.89	Somewhat limited	0.78	Somewhat limited	0.78
		Restricted		Restricted		Restricted	
		permeability		permeability		permeability	
		Too steep for surface application		Too steep for surface application		Too steep for surface application	0.08
Mc: Mccook-----	100	Somewhat limited	0.00	Somewhat limited	0.40	Somewhat limited	0.00
		Filtering		Flooding		Filtering	
		capacity		capacity	0.00	capacity	
Ns: Ness-----	100	Very limited	1.00	Very limited	1.00	Very limited	1.00
		Restricted		Restricted		Restricted	
		permeability		permeability		permeability	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
Ns: Ness-----	100	Runoff limitation	0.40				

AGRICULTURAL WASTE MANAGEMENT--Continued
McPherson County, Kansas

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct of map unit	Application of manure and food- processing waste		Application of sewage sludge		Disposal of wastewater by irrigation	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
Nw: New Cambria-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability Flooding	1.00 0.40	Very limited Restricted permeability	1.00
Pa: Plevna-----	100	Very limited Flooding Depth to saturated zone Runoff limitation Filtering capacity	1.00 1.00 0.40 0.00	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 0.00	Very limited Flooding Depth to saturated zone Filtering capacity	1.00 1.00 0.00
Pr: Pratt-----	100	Very limited Filtering capacity Leaching limitation Slope	1.00 0.45 0.04	Very limited Filtering capacity Slope	1.00 0.04	Very limited Too steep for surface application Filtering capacity Too steep for sprinkler application	1.00 1.00 0.22
Ro: Roxbury-----	100	Not limited		Somewhat limited Flooding	0.40	Not limited	
Sm: Smolan-----	100	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00	Very limited Restricted permeability	1.00
To: Tobin-----	100	Somewhat limited Flooding	0.60	Very limited Flooding	1.00	Somewhat limited Flooding	0.60
W: Water-----	100	Not rated		Not rated		Not rated	
Wb: Wells-----	90	Somewhat limited Too acid	0.03	Somewhat limited Too acid	0.14	Somewhat limited Too acid	0.14
Wc: Wells-----	100	Somewhat limited Too acid	0.03	Somewhat limited Too acid	0.14	Somewhat limited Too steep for surface application Too acid	0.31 0.14

WIN-PST SPISP II
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL_KS Sort Order: MUSYM

McPherson County, Kansas: KS113

MUSYM/SEQ#	COMPONENT/TEXTURE/MU%	HYD	KFACT	SURFACE DEPTH	% OM	SPISP II Ratings		
						Leaching (SLP)	Solution Runoff (SSRP)	Adsorbed Runoff (SARP)
053ED 1	EDALGO L 100%	C	0.32	8"	3.0%	L	H	H
053MB 1	MCCOOK L 100%	B	0.32	12"	3.0%	I	I	I
053MC 1	MCCOOK SIL 100%	B	0.32	12"	3.0%	I	I	I
079GE 1	GEARY SIL 100%	B	0.32	9"	2.5%	I	I	I
079LB 1	LADYSMITH SICL 100%	D	0.37	10"	3.0%	V	H	H
115IC 1	IRWIN SICL 90%	D	0.37	13"	3.0%	V	H	H
1191 1	BLAZEFORK SICL 90%	D	0.37	3"	3.0%	V	H	H
1200 1	BUHLER SICL 65%	D	0.43	3"	4.0%	V	H	H
1200 2	BLAZEFORK SICL 30%	D	0.37	3"	3.0%	V	H	H
1324 1	CARWAY FSL 50%	D	0.20	7"	0.8%	V	H	H
1324 2	CARBIKA SIL 30%	D	0.24	11"	1.5%	V	H	H
1553 1	DARLOW L 70%	C	0.43	5"	2.0%	L	H	H
1553 2	ELMER FSL 20%	C	0.32	6"	1.5%	L	H	H
1555 1	DILLHUT FS 35%	B	0.15	4"	0.5%	H	I	I
1555 2	PLEV LFS 35%	B	0.17	4"	0.5%	H (w)	I	I
159DP 1	DILLWYN LFS 60%	A	0.17	9"	1.0%	H (w)	L	L
159DP 2	PLEVNA LFS 40%	D	0.17	10"	2.5%	H (w)	H	H
159GC 1	GEARY SIL 75%	B	0.32	7"	2.5%	I	I	I
159GC 2	CLARK CL 25%	B	0.28	11"	1.5%	I	I	I
159LA 1	LANCASTER L 90%	B	0.28	7"	2.5%	I	I	I
159NF 1	NARON FSL 100%	B	0.20	14"	2.0%	I	I	I
159PF 1	PRATT LFS 100%	A	0.17	13"	0.8%	H	L	L
159PR 1	PRATT LFS 65%	A	0.17	13"	0.8%	H	L	L
159PR 2	CARWILE FSL 35%	D	0.24	12"	2.0%	H (w)	H	H
159SO 1	SMOLAN SICL 100%	C	0.37	6"	3.0%	L	H	H
159WA 1	WALDECK FSL 100%	C	0.20	25"	1.5%	H (w)	H	I
169CE 1	CLIME SICL 100%	C	0.37	8"	3.0%	L	H	H
169ED 1	EDALGO CL 100%	C	0.37	9"	3.0%	L	H	H

WIN-PST SPISP II
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL_KS Sort Order: MUSYM

McPherson County, Kansas: KS113

169RO 1	ROXBURY SIL 99%	B	0.32	26"	3.0% L	I	I
1985 1	HAYES FSL 60%	B	0.20	8"	0.8% H	I	I
2391 1	KASKAN SICL 75%	B	0.37	9"	3.0% I	I	I
2588 1	LONGFORD SICL 90%	C	0.37	6"	1.5% L	H	H
3181 1	PRATT FS 45%	A	0.15	8"	0.8% H	L	L
3181 2	TURON FS 30%	A	0.15	8"	0.5% H	L	L
3641 1	TIVIN FS 45%	A	0.15	7"	0.5% H	L	L
3641 2	DILLHUT FS 40%	B	0.15	4"	0.5% H	I	I
AED 1	ARENTS, EARTHEN DAM 100%		0.00	0"	0.0% ?	?	?
At 1	ATTICA LFS 100%	B	0.17	10"	0.8% H	I	I
Br 1	BRIDGEPORT SIL 100%	B	0.32	14"	2.5% I	I	I
Ca 1	CARWILE FSL 100%	D	0.24	16"	2.0% H (w)	H	H
Cb 1	CASS FSL 100%	B	0.20	7"	1.5% H	I	I
Ce 1	CLIME SIC 100%	C	0.28	9"	2.5% L	H	H
Cm 1	CLIME SIC 100%	C	0.28	9"	2.5% L	H	H
Cr 1	CRETE SIL 100%	C	0.37	5"	3.0% L	H	H
Cs 1	CRETE SIL 100%	C	0.37	6"	3.0% L	H	H
Ct 1	CRETE SICL 100%	C	0.37	6"	3.0% L	H	H
De 1	DETROIT SICL 100%	C	0.37	16"	3.0% L	H	H
Dr 1	DRUMMOND L 100%	D	0.49	19"	0.8% H (w)	H	H
Ed 1	EDALGO SIL 100%	C	0.37	6"	3.0% L	H	H
Ee 1	EDALGO SICL 100%	C	0.37	8"	3.0% L	H	H
Fa 1	FARNUM L 100%	B	0.28	12"	2.0% I	I	I
Ge 1	GEARY SIL 100%	B	0.32	7"	2.5% I	I	I
Go 1	GOESSEL SIC 100%	D	0.28	14"	2.5% H (w)	H	H
Ho 1	HORD SIL 100%	B	0.32	6"	3.0% I	I	I
INT 1	AQUOLLS VAR 100%	C	0.00	72"	0.0% ?	H	?
Ir 1	IRWIN SICL 100%	D	0.37	11"	3.0% V	H	H
La 1	LADYSMITH SICL 100%	D	0.37	8"	3.0% V	H	H
Ld 1	LADYSMITH SICL 70%	D	0.37	8"	3.0% V	H	H

WIN-PST SPISP II
SOIL SENSITIVITY TO PESTICIDE LOSS RATING REPORT

Soils Data Table: SOIL_KS Sort Order: MUSYM

McPherson County, Kansas: KS113

Ld 2	DRUMMOND L 30%	D	0.49	7"	0.8% H (w)	H	H
Le 1	LANCASTER L 100%	B	0.28	10"	2.5% I	I	I
Lh 1	LANCASTER L 60%	B	0.28	10"	2.5% I	I	I
Lh 2	HEDVILLE L 40%	D	0.32	15"	2.5% V	H	H
LHH 1	LANCASTER L 60%	B	0.28	9"	2.5% I	I	I
LHH 2	HEDVILLE L 40%	D	0.32	8"	2.5% V	H	H (s)
Ln 1	LONGFORD SICL 100%	C	0.32	15"	2.5% L	H	H
Lo 1	LONGFORD SICL 100%	C	0.32	6"	1.5% L	H	H
Mc 1	MCCOOK FSL 100%	B	0.20	14"	1.5% I	I	I
Ns 1	NESS SIC 100%	D	0.28	60"	2.0% H (w)	H	H
Nw 1	NEW CAMBRIA SIC 100%	C	0.28	12"	3.0% L	H	H
Pa 1	PLEVNA FSL 100%	D	0.20	18"	2.5% H (w)	H	H
Pr 1	PRATT LFS 100%	A	0.17	11"	0.8% H	L	L
Ro 1	ROXBURY SICL 100%	B	0.43	21"	3.0% L	I	I
Sm 1	SMOLAN SICL 100%	C	0.37	11"	3.0% L	H	H
To 1	TOBIN SIL 100%	B	0.32	20"	2.5% L	I	I
W 1	WATER 100%		0.00	0"	0.0% ?	?	?
Wb 1	WELLS L 90%	B	0.28	15"	2.5% I	I	I
Wc 1	WELLS L 100%	B	0.28	11"	2.5% I	I	I

(.\REPORTS\SOILS.TXT generated on 12/12/01 at 12:11:15)

H -- High
I -- Intermediate
L -- Low
V -- Very Low

Conditions that affect ratings:

- m -- There are macropores in the surface horizon deeper than 24"
- w -- The high water table comes within 24" of the surface during the growing season
- s -- The field slope is greater than 15%

SPISP II S-Ratings:

- SLP -- Soil Leaching Potential
- SSRP -- Soil Solution Runoff Potential
- SARP -- Soil Adsorbed Runoff Potential

In this section, hydric soils are defined and described and the hydric soils in the survey area are listed. The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin and others, 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for each of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part (Federal Register, 1994). These soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 1995). These criteria are used to identify a phase of a soil series that normally is associated with wetlands. The criteria used are selected estimated soil properties that are described in "Soil Taxonomy" (USDA, 1999) and "Keys to Soil Taxonomy" (USDA, 1998) and in the "Soil Survey Manual" (USDA, 1993).

If soils are wet enough for a long enough period to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils in this survey area are specified in "Field Indicators of Hydric Soils in the United States" (Hurt and others, 1996).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

Map units in the Hydric Soil Interpretations table meet the definition of hydric soils and, in addition, have at least one of the hydric soil indicators. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; Hurt and others, 1996).

Map units that are made up of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units made up of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform.

These map units, in general, do not meet the definition of hydric soils because they do not have one of the hydric soil indicators. A portion of these map units, however, may include hydric soils. Onsite investigation is recommended to determine whether hydric soils occur and the location of the included hydric soils.

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
McPherson County, Kansas

PAGE 2 of 6

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
053ED: EDALGO LOAM, 3 TO 7 PERCENT SLOPES	EDALGO	No	hillslope	---	---	---	---
053MB: MCCOOK LOAM, OCCASIONALLY FLOODED	MCCOOK	No	flood plain	---	---	---	---
053MC: MCCOOK SILT LOAM, RARELY FLOODED	MCCOOK	No	flood plain	---	---	---	---
079GE: GEARY SILT LOAM, 3 TO 6 PERCENT SLOPES	GEARY	No	hillslope	---	---	---	---
079LB: LADYSMITH SILTY CLAY LOAM, 1 TO 2 PERCENT SLOPES	LADYSMITH	No	paleoterrace	---	---	---	---
	UNNAMED HYDRIC SOILS	Yes	depression	2B3,3,2A	YES	NO	YES
115IC: IRWIN SILTY CLAY LOAM, 3 TO 6 PERCENT SLOPES	IRWIN	No	hillslope	---	---	---	---
	CLIME	No	hillslope	---	---	---	---
	LABETTE	No	hillslope	---	---	---	---
159DP: DILLWYN-PLEVNA LOAMY FINE SANDS, 0 TO 1 PERCENT SLOPES	DILLWYN	No	interdune, dune, paleoterrace	---	---	---	---
	PLEVNA	Yes	flood plain	4,2B3	YES	YES	NO
159GC: GEARY-CLARK COMPLEX, 3 TO 7 PERCENT SLOPES, ERODED	GEARY	No	hillslope	---	---	---	---
	CLARK	No	paleoterrace	---	---	---	---
159LA: LANCASTER LOAM, 1 TO 3 PERCENT SLOPES	LANCASTER	No	hillslope	---	---	---	---
	HEDVILLE	No	hillslope	---	---	---	---
	IRWIN	No	hillslope	---	---	---	---
159NF: NARON FINE SANDY LOAM, 1 TO 3 PERCENT SLOPES	NARON	No	dune, paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
	Unnamed wet soils	Yes	depression	2A,3,2B3	YES	NO	YES
159PF: PRATT LOAMY FINE SAND, 1 TO 5 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
159PR: PRATT-CARWILE COMPLEX, 0 TO 5 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
	Unnamed wet soils	Yes	depression	2B3,2A,3	YES	NO	YES
159SO: SMOLAN SOILS, 2 TO 7 PERCENT SLOPES, ERODED	SMOLAN	No	---	---	---	---	---
159WA: WALDECK FINE SANDY LOAM, OCCASIONALLY FLOODED	WALDECK	No	flood plain	---	---	---	---
	Unnamed wet soils	Yes	depression	2A,3,2B3	YES	NO	YES
169CE: CLIME SILTY CLAY LOAM, 2 TO 6 PERCENT SLOPES	CLIME	No	hillslope	---	---	---	---
169ED: EDALGO CLAY LOAM, 3 TO 7 PERCENT SLOPES	EDALGO	No	hillslope	---	---	---	---
	SIDEHILL SEEP	Yes	---	2B2	YES	NO	NO
169RO: ROXBURY SILT LOAM, RARELY FLOODED	ROXBURY	No	flood plain	---	---	---	---
	UNNAMED HYDRIC SOIL	Yes	depression, terrace	3	NO	NO	YES

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
McPherson County, Kansas

PAGE 3 of 6

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
1191: BLAZEFOK SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES, RARELY FLOODED	BLAZEFOK	No	flood plain	---	---	---	---
	TOBIN Unnamed Wet Soils	No Yes	flood plain drainageway	--- 2A,3,4	--- YES	--- YES	--- YES
1200: BUHLER-BLAZEFOK SILTY CLAY LOAMS, 0 TO 1 PERCENT SLOPES, RARELY FLOODED	BUHLER	No	flood plain	---	---	---	---
	BLAZEFOK TOBIN Unamed Wet Soils	No No Yes	flood plain flood plain drainageway	--- --- 2A,3,4	--- --- YES	--- --- YES	--- --- YES
1324: CARWAY AND CARBIKA SOILS, 0 TO 1 PERCENT SLOPES	CARWAY	Yes	depression, interdune, paleoterrace	3,2B3	YES	NO	YES
	CARBIKA	Yes	depression, interdune, paleoterrace	2B3,3	YES	NO	YES
	SOLVAY	No	interdune, paleoterrace	---	---	---	---
1553: DARLOW-ELMER COMPLEX, 0 TO 2 PERCENT SLOPES	DARLOW	No	terrace	---	---	---	---
	ELMER	No	terrace	---	---	---	---
	PUNKIN	No	paleoterrace	---	---	---	---
	CARBIKA	Yes	depression, interdune, paleoterrace	3,2B3	YES	NO	YES
	CARWAY	Yes	depression, interdune, paleoterrace	2B3,3	YES	NO	YES
1555: DILLHUT-PLEV COMPLEX, 0 TO 2 PERCENT SLOPES	DILLHUT	No	dune, paleoterrace	---	---	---	---
	PLEV	Yes	depression, interdune, paleoterrace	2B2	YES	NO	NO
	DILLWYN	No	interdune, dune, paleoterrace	---	---	---	---
	WARNUT	Yes	interdune, depression, paleoterrace	2B3,3	YES	NO	YES
1985: HAYES FINE SANDY LOAM, 1 TO 5 PERCENT SLOPES	HAYES	No	dune, paleoterrace	---	---	---	---
	ATTICA	No	dune, paleoterrace	---	---	---	---
	SALTCREEK	No	dune, paleoterrace	---	---	---	---
2391: KASKAN SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES, FREQUENTLY FLOODED, CHANNELED	KASKAN	No	flood plain	---	---	---	---
	TOBIN Unnamed Wet Soils	No Yes	flood plain depression, drainageway	--- 2B1,2B2,2B3,4	--- YES	--- YES	--- NO
2588: LONGFORD SILTY CLAY LOAM, 3 TO 7 PERCENT SLOPES, MODERATELY ERODED	LONGFORD	No	hillslope	---	---	---	---
	GEARY	No	hillslope	---	---	---	---
3181: PRATT-TURON FINE SANDS, 1 TO 5 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
	TURON	No	dune, paleoterrace	---	---	---	---
	HAYES	No	dune, paleoterrace	---	---	---	---
	CARWAY	Yes	depression, interdune, paleoterrace	2B3,3	YES	NO	YES
	WARNUT	Yes	interdune, depression, paleoterrace	3,2B3	YES	NO	YES

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
McPherson County, Kansas

PAGE 4 of 6

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
3641: TIVIN-DILLHUT FINE SANDS, 0 TO 15 PERCENT SLOPES	TIVIN	No	dune, paleoterrace	---	---	---	---
	DILLHUT	No	dune, paleoterrace	---	---	---	---
	SOLVAY	No	interdune, paleoterrace	---	---	---	---
	CARWAY	Yes	depression, interdune, paleoterrace	3,2B3	YES	NO	YES
	WARNUT	Yes	interdune, depression, paleoterrace	3,2B3	YES	NO	YES
	PLEV	Yes	depression, interdune, paleoterrace	2B2	YES	NO	NO
AED: ARENTS, EARTHEN DAM	ARENTS, EARTHEN DAM	Unranked	---	---	---	---	---
At: ATTICA LOAMY FINE SAND, 1 TO 4 PERCENT SLOPES	ATTICA	No	dune, paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
	PLEVNA	Yes	flood plain	3,2B3	YES	NO	YES
	Unnamed wet soils	Yes	depression	2A,2B3,3	YES	NO	YES
Br: BRIDGEPORT SILT LOAM, RARELY FLOODED	BRIDGEPORT	No	flood plain	---	---	---	---
Ca: CARWILE FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
Cb: CASS FINE SANDY LOAM, RARELY FLOODED	CASS	No	flood plain	---	---	---	---
	BRIDGEPORT	No	---	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
Ce: CLIME SILTY CLAY, 1 TO 3 PERCENT SLOPES	CLIME	No	hillslope	---	---	---	---
Cm: CLIME SILTY CLAY, 3 TO 6 PERCENT SLOPES	CLIME	No	hillslope	---	---	---	---
Cr: CRETE SILT LOAM, 0 TO 1 PERCENT SLOPES	Unnamed wet soils	Yes	drainageway	3,2B3,2A,4	YES	YES	YES
	CRETE	No	---	---	---	---	---
Cs: CRETE SILT LOAM, 1 TO 3 PERCENT SLOPES	CRETE	No	hillslope	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2A,2B3,4	YES	YES	NO
Ct: CRETE SILTY CLAY LOAM, 1 TO 3 PERCENT SLOPES, ERODED	CRETE	No	---	---	---	---	---
De: DETROIT SILTY CLAY LOAM, RARELY FLOODED	DETROIT	No	flood plain	---	---	---	---
	UNNAMED HYDRIC SOILS	Yes	depression, terrace	2B3,3	YES	NO	YES
Dr: DRUMMOND LOAM, 0 TO 1 PERCENT SLOPES	DRUMMOND	No	terrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
	Unnamed wet soils	Yes	depression	3,2B3	YES	NO	YES
Ed: EDALGO SILT LOAM, 5 TO 12 PERCENT SLOPES	EDALGO	No	hillslope	---	---	---	---
	CLIME	No	hillslope	---	---	---	---
	LANCASTER	No	---	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3	YES	NO	NO

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
McPherson County, Kansas

PAGE 5 of 6

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Ee: EDALGO SILTY CLAY LOAM, 3 TO 9 PERCENT SLOPES, ERODED	EDALGO	No	---	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3	YES	NO	NO
Fa: FARNUM LOAM, 1 TO 3 PERCENT SLOPES	FARNUM	No	paleoterrace	---	---	---	---
Ge: GEARY SILT LOAM, 1 TO 3 PERCENT SLOPES	GEARY	No	hillslope	---	---	---	---
Go: GOESSEL SILTY CLAY, 0 TO 2 PERCENT SLOPES	GOESSEL	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2A,3,2B3	YES	NO	YES
Ho: HORD SILT LOAM, RARELY FLOODED	HORD	No	hillslope	---	---	---	---
	Unnamed wet soils	Yes	depression	2A,3,2B3,4	YES	YES	YES
INT: AQUOLLS	AQUOLLS	Yes	depression, terrace	2B3,3	YES	NO	YES
Ir: IRWIN SILTY CLAY LOAM, 1 TO 3 PERCENT SLOPES	IRWIN	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3	YES	NO	NO
La: LADYSMITH SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES	LADYSMITH	No	paleoterrace	---	---	---	---
	Unnamed wet soils	Yes	depression	2A,3,2B3,4	YES	YES	YES
Ld: LADYSMITH-DRUMMOND COMPLEX, 0 TO 1 PERCENT SLOPES	LADYSMITH	No	paleoterrace	---	---	---	---
	DRUMMOND Unnamed wet soils	No Yes	terrace depression	--- 2A,3,2B3	--- YES	--- NO	--- YES
Le: LANCASTER LOAM, 2 TO 6 PERCENT SLOPES	LANCASTER	No	hillslope	---	---	---	---
Lh: LANCASTER-HEDVILLE LOAMS, 6 TO 12 PERCENT SLOPES	LANCASTER	No	hillslope	---	---	---	---
	HEDVILLE	No	hillslope	---	---	---	---
LHH: LANCASTER-HEDVILLE COMPLEX, 3 TO 20 PERCENT SLOPES	LANCASTER	No	hillslope	---	---	---	---
	HEDVILLE SIDEHILL SEEP	No Yes	hillslope hillslope	--- 2B2	--- YES	--- NO	--- NO
Ln: LONGFORD SILTY CLAY LOAM, 3 TO 6 PERCENT SLOPES	LONGFORD	No	hillslope	---	---	---	---
	Unnamed wet soils	Yes	drainageway	4,2B3	YES	YES	NO
Lo: LONGFORD SILTY CLAY LOAM, 2 TO 6 PERCENT SLOPES, ERODED	LONGFORD	No	hillslope	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3,4	YES	YES	NO
Mc: MCCOOK FINE SANDY LOAM, RARELY FLOODED	MCCOOK	No	flood plain	---	---	---	---
Ns: NESS SILTY CLAY	NESS	Yes	playa	2B3,3	YES	NO	YES
	Unnamed wet soils	Yes	depression	2A,3,2B3	YES	NO	YES
Nw: NEW CAMBRIA SILTY CLAY, RARELY FLOODED	NEW CAMBRIA	No	stream terrace	---	---	---	---
	UNNAMED HYDRIC SOILS	Yes	oxbow lake	3,2B3	YES	NO	YES

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
McPherson County, Kansas

PAGE 6 of 6

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

Map symbol and map unit name	Component	Hydric	Local landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
Pa: PLEVNA FINE SANDY LOAM, FREQUENTLY FLOODED	PLEVNA	Yes	flood plain	4,2B3	YES	YES	NO
Pr: PRATT LOAMY FINE SAND, 6 TO 12 PERCENT SLOPES	PRATT	No	dune, paleoterrace	---	---	---	---
	CARWILE	Yes	depression, paleoterrace	2A	YES	NO	NO
	PLEVNA	Yes	flood plain	3,2B3	YES	NO	YES
	Unnamed wet soils	Yes	depression	3,2B3,2A	YES	NO	YES
Ro: ROXBURY SILTY CLAY LOAM, RARELY FLOODED	ROXBURY	No	flood plain	---	---	---	---
Sm: SMOLAN SILTY CLAY LOAM, 1 TO 3 PERCENT SLOPES	SMOLAN	No	hillslope	---	---	---	---
To: TOBIN SILT LOAM, OCCASIONALLY FLOODED	TOBIN	No	flood plain	---	---	---	---
	UNNAMED HYDRIC SOIL	Yes	flood plain, marsh	2B3	YES	NO	NO
	UNNAMED HYDRIC SOILS	Yes	depression, flood plain	3	NO	NO	YES
	Unnamed wet soils	Yes	drainageway	4,2B3,2A,3	YES	YES	YES
W: WATER	WATER	Yes	---	3,4	NO	YES	YES
Wb: WELLS LOAM, 1 TO 3 PERCENT SLOPES	WELLS	No	hillslope	---	---	---	---
	CLIME	No	hillside	---	---	---	---
	IRWIN	No	hillside	---	---	---	---
Wc: WELLS LOAM, 3 TO 6 PERCENT SLOPES	WELLS	No	hillslope	---	---	---	---
	Unnamed wet soils	Yes	drainageway	2B3	YES	NO	NO

FOOTNOTE: There may be small areas of included soils or miscellaneous areas that are significant to use and management of the soil; yet are too small to delineate on the soil map at the map's original scale. These may be designated as spot symbols and are defined in the published Soil Survey Report or the USDA-NRCS Technical Guide, Part II. Areas mapped as water or any map unit that contains one of the following conventional symbols is considered a hydric soil map unit: marshes or swamps; wet spots; depressions; streams, lakes and ponds.

1. All Histosols except Folists, or
2. Soils in Aquic suborders, great groups, or subgroups, Albolls suborder, Aquisalids, Pachic subgroups, or Cumulic subgroups that are:
 - a. Somewhat poorly drained with a water table equal to 0.0 foot (ft) from the surface during the growing season, or
 - b. poorly drained or very poorly drained and have either:
 - (1) water table equal to 0.0 ft during the growing season if textures are coarse sand, sand, or fine sand in all layers within 20 inches (in), or for other soils
 - (2) water table at less than or equal to 0.5 ft from the surface during the growing season if permeability is equal to or greater than 6.0 in/hour (h) in all layers within 20 in, or
 - (3) water table at less than or equal to 1.0 ft from the surface during the growing season if permeability is less than 6.0 in/h in any layer within 20 in, or
3. Soils that are frequently ponded for long duration or very long duration during the growing season, or
4. Soils that are frequently flooded for long duration or very long duration during the growing season.