

RANGELAND PRODUCTIVITY
Stevens County, Kansas

Use and Explanation of Rangeland, Grazed Forest Land, Native Pastureland Interpretations

Information in this subsection can be used to plan the use and management of soils for rangeland, grazed forest land, and native pasture. Different kinds of soils vary in their capacity to produce native grasses and other plants suitable for grazing. Information in this subsection provides groupings of similar soils and estimates of potential forage production, which can be used to determine livestock stocking rates.

Rangeland. Range is land on which the native vegetation (climax or natural potential plant community) is predominantly grasses, grasslike plants, forbs, and shrubs suitable for grazing and browsing. Range includes natural grasslands, savannas, many wetlands, some deserts, tundra, and certain shrub and forb communities. Rangeland receives no regular or frequent cultural treatment. The composition and production of the plant community are determined by soil, climate, topography, overstory canopy, and grazing management.

Grazed Forest Land. Includes land on which the understory includes, as an integral part of the forest plant community, plants that can be grazed without significantly impairing other forest values.

Native Pasture. Includes land on which the native vegetation (climax or natural potential plant community) is forest but which is used and managed primarily for production of native plants for forage. Native pasture includes cut-over forest land and forest land cleared and now managed for native or naturalized forage plants.

Rangeland

In areas that have similar climate and topography, differences in the kind and amount of vegetation produced on rangeland are closely related to the kind of soil. Effective management based on the relationship between the soils and vegetation and water.

The Rangeland, Grazed Forest land, Native Pastureland Interpretations shows, for each soil that supports rangeland vegetation, the ecological site and the potential annual production of vegetation in favorable, normal, unfavorable years. An explanation of the column headings in this table follows.

An ecological site is the product of all the environmental factors responsible for its development. It has characteristic soils that have developed over time throughout the soil development process; a characteristic hydrology, particularly infiltration and runoff, that has developed over time; and a characteristic plant community (kind and amount of vegetation). The hydrology of a site is influenced by development of the soil and plant community. The vegetation, soils, and hydrology are all interrelated. Each is influenced by the others and influences the development of the others. The plant community on an ecological site is typified by an association of species that differs from that of other ecological sites in the kind and/or proportion of species or in total production. Descriptions of ecological sites are provided in the Field Office Technical Guide, which is available in local offices of the Natural Resources Conservation Service.

Total dry-weight production is the amount of vegetation that can be expected to grow annually on well managed rangeland that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, average, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are about average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Yields are adjusted to a common percent of air-dry moisture content.

Range management requires a knowledge of the kinds of soil and of the potential natural plant community. It also requires an evaluation of the present range similarity index and rangeland trend. Range similarity index is determined by comparing the present plant community with the potential natural plant community on a particular rangeland ecological site. The more closely the existing community resembles the potential community, the higher the range similarity index. Rangeland trend is defined as the direction of change in an existing plant community relative to the potential natural plant community. Further information about the range similarity index and rangeland trend is available in chapter 4 of the National Range and Pasture Handbook, which is available in local offices of the Natural Resources Conservation Service. The objective in range management is to control grazing so that the plants growing on a site are about the same in kind and amount as the potential natural plant community for that site. Such management generally results in the optimum production of vegetation, control of undesirable brush species, conservation of water, and control of erosion. Sometimes, however, an area with a range similarity index somewhat below the potential meets grazing needs, provides wildlife habitat, and protects soil and water resources.

RANGELAND PRODUCTIVITY--Continued
Stevens County, Kansas

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
067LO: Pleasant-----	Clay Upland (pe16-20)	2,400	1,800	1,000
067OF: Otero-----	Sandy (pe17-20)	1,800	1,500	1,000
067OG: Otero-----	Sandy (pe17-20)	1,800	1,500	1,000
Schamber-----	Gravelly Hills (pe17-20)	1,800	1,300	700
067SA: Satanta-----	Sandy (pe16-20)	3,000	2,200	1,200
067SB: Satanta-----	Sandy (pe16-20)	3,000	2,200	1,200
067TF: Valent-----	Choppy Sands (pe16-20)	2,500	2,000	1,200
067TV: Valent-----	Choppy Sands (pe16-20)	2,500	2,000	1,200
Vona-----	Sands (pe16-20)	2,200	1,700	900
067UE: Ulysses-----	Loamy Upland (pe16-20)	2,400	1,800	1,000
Colby-----	Limy Upland (pe16-20)	1,600	1,200	800
067VO: Vona-----	Sands (pe16-20)	2,200	1,700	900
081OG: Otero-----	Sandy (pe17-20)	1,800	1,500	1,000
Schamber-----	Gravelly Hills (pe17-20)	1,800	1,300	700
1044: Atchison-----	Limy Upland (pe16-20)	2,700	1,900	1,200
1046: Atchison-----	Limy Upland (pe16-20)	2,700	1,900	1,200
1182: Belfon-----	Loamy Upland (pe17-20)	3,000	2,000	1,250
1184: Bigbow-----	Loamy Upland (pe17-20)	3,000	2,000	1,250
1185: Bigbow-----	Loamy Upland (pe17-20)	3,000	2,000	1,250
1504: Dalhart-----	Loamy Upland (pe17-20)	3,000	2,000	1,250
1505: Dalhart-----	Loamy Upland (pe17-20)	3,000	2,000	1,250
1506: Dalhart-----	Sandy (pe17-20)	2,300	1,750	1,250
1558: Dalhart-----	Sandy (pe17-20)	2,300	1,750	1,250
1559: Dalhart-----	Sandy (pe17-20)	2,300	1,750	1,250
Eva-----	Sands (pe17-20)	2,400	1,850	1,250
1670: Eva-----	Sands (pe17-20)	2,400	1,850	1,250
1671: Eva-----	Sands (pe17-20)	2,400	1,850	1,250
Optima-----	Choppy Sands (pe17-20)	2,250	1,250	900
1672: Eva-----	Sands (pe17-20)	2,400	1,850	1,250
1723: Feterita-----	Lakebed (pe16-20)	3,500	2,350	1,300
1979: Haverson-----	Sandy Lowland (pe16-20)	3,750	2,750	1,750
1980: Happyditch-----	Sandy Lowland (pe17-20)	3,750	2,750	1,750
1981: Happyditch-----	Sandy Lowland (pe17-20)	3,750	2,750	1,750
1984: Happyditch-----	Sandy Lowland (pe17-20)	3,750	2,750	1,750
3047: Optima-----	Choppy Sands (pe17-20)	2,250	1,250	900
3048: Optima-----	Choppy Sands (pe17-20)	2,250	1,250	900
3415: Satanta-----	Loamy Upland (pe16-20)	3,000	2,000	1,250
3506: Shore-----	Loamy Lowland (pe16-20)	4,500	3,500	2,000
3725: Ulysses-----	Loamy Upland (pe16-20)	3,000	2,000	1,250
3969: Wagonbed-----	Limy Upland (pe16-20)	2,700	1,900	1,200
Bo: Valent-----	Choppy Sands (pe17-20)	2,500	2,000	1,200
CIR: Cimarron River-----	---	---	---	---
Cm: Colby-----	Limy Upland (pe17-20)	1,600	1,200	800
Da: Dalhart-----	Sandy (pe17-20)	2,300	1,600	1,000
Db: Dalhart-----	Sandy (pe17-20)	2,300	1,600	1,000
Df:				

RANGELAND PRODUCTIVITY--Continued
Stevens County, Kansas

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

Map symbol and soil name	Ecological site	Total dry-weight production		
		Favorable year	Average year	Unfavorable year
		Lb/acre	Lb/acre	Lb/acre
Dalhart-----	Sands (pe17-20)	3,000	1,900	1,200
Dx:				
Dalhart-----	Sandy (pe17-20)	2,300	1,600	1,000
Otero-----	Sandy (pe17-20)	1,800	1,500	1,000
Go:				
Goshen-----	Loamy Terrace (pe16-20)	3,300	2,900	2,500
Lf:				
Lincoln-----	Sandy Lowland (pe17-20)	3,000	2,300	1,800
Lo:				
Pleasant-----	Lakebed (pe17-20)	2,400	1,800	1,000
Lp:				
Pleasant-----	Lakebed (pe17-20)	2,400	1,800	1,000
Ma:				
Penden-----	Limy Upland (pe17-20)	4,000	2,500	1,000
Mb:				
Penden-----	Limy Upland (pe17-20)	4,000	2,500	1,000
Mx:				
Penden-----	Limy Upland (pe17-20)	4,000	2,500	1,000
Otero-----	Sandy (pe17-20)	1,800	1,500	1,000
My:				
Manter-----	Sandy (pe17-20)	2,000	1,600	800
Ot:				
Otero-----	Sandy (pe17-20)	1,800	1,500	1,000
Ra:				
Satanta-----	Loamy Upland (pe17-20)	3,000	2,000	1,000
Rb:				
Haxtun-----	Sands (pe17-20)	2,500	2,000	1,500
Rm:				
Richfield-----	Loamy Upland (pe17-20)	2,400	1,800	800
Rx:				
Richfield-----	Loamy Upland (pe17-20)	2,400	1,800	800
Ulysses-----	Loamy Upland (pe17-20)	2,400	1,800	1,000
Tf:				
Valent-----	Choppy Sands (pe17-20)	2,500	2,000	1,200
Ua:				
Ulysses-----	Loamy Upland (pe17-20)	2,400	1,800	1,000
Ub:				
Ulysses-----	Loamy Upland (pe17-20)	2,400	1,800	1,000
Ue:				
Ulysses-----	Loamy Upland (pe17-20)	2,400	1,800	1,000
Colby-----	Limy Upland (pe17-20)	1,600	1,200	800
Vo:				
Vona-----	Sands (pe17-20)	2,200	1,700	900
Vx:				
Vona-----	Sands (pe17-20)	2,200	1,700	900
Valent-----	Choppy Sands (pe17-20)	2,500	2,000	1,200
W:				
Water-----	---	---	---	---

